

OVERSEAS ENVIRONMENTAL ASSESSMENT
For
Office of Naval Research Arctic Research Activities in the
Beaufort and Chukchi Seas 2022-2025

August 2022



This page left intentionally blank.

Abstract

Designation:	Overseas Environmental Assessment
Title of Proposed Action:	Office of Naval Research Arctic Research Activities in the Beaufort Sea 2022-2025
Project Location:	Beaufort and Chukchi Seas
Lead Agency for the OEA:	Department of the Navy
Affected Region:	Beaufort Sea, Chukchi Sea, Arctic
Action Proponent:	Office of Naval Research
Point of Contact:	Raymond Soukup Office of Naval Research Program Officer Email address:raymond.j.soukup.civ@us.navy.mil
Date:	August 2022

The Office of Naval Research (ONR) prepared this Overseas Environmental Assessment (OEA) in compliance with the Executive Order (E.O.) 12114, Department of Defense regulations found at 32 Code of Federal Regulations Part 187, Department of Defense Directive 6050.7, and the Chief of Naval Operations Instruction 5090.1E and its accompanying manual (M-5090.1).

This OEA evaluates the potential harm to the environment from ONR Arctic Research Activities during the period September 2022-September 2025. The Naval need for this scientific research relates to environmental characterization in support of combat capable forces ready to deploy worldwide in accordance with Title 10 United States Code (U.S.C.) section 8062, and to support the aims of the Arctic Research and Policy Act (15 U.S.C. sections 4101 et seq.). For the Arctic this consists of potential submarine and surface ship operations with active sonar for anti-submarine warfare and submarine/surface ship force protection. The characterization of the potential Arctic battlespace, given the changes in water properties and ice cover, is critical to performance predictions for active and passive acoustic systems. The year-round characterization of the arctic environment requires characterization of the environment by leave-behind sources and autonomous vehicles, and the research projects are geared toward building multiple sources transmitting intermittently to allow vehicles to transmit under the ice. The purpose of the Proposed Action is to test the feasibility of using a field of active acoustic sources as navigation aids to unmanned vehicles collecting oceanographic and ice data under ice-covered conditions. This OEA evaluates three alternatives: the No Action Alternative and two Action Alternatives. Alternative 1 would conduct all the scientific research described in the Proposed Action, excluding the use of the very low frequency (VLF) sources. Under Alternative 2 (Preferred Alternative), all of the scientific research described in the Proposed Action would occur, including year-round use of the VLF sources. Sources would typically be recovered and reactivated on a yearly basis.

In this OEA, the Navy analyzes potential harm to the environment that could result from the No Action Alternative and two Action Alternatives. The resources evaluated include physical resources (i.e., marine habitats), marine invertebrates, marine birds, fish, Essential Fish Habitat, and marine mammals.

This page intentionally left blank

EXECUTIVE SUMMARY

Proposed Action

Office of Naval Research's (ONR) Proposed Action, called Arctic Research Activities (ARA), is to conduct scientific research in the Beaufort and Chukchi Seas from September 2022 to September 2025. This research comprises cruises that would occur annually beginning in September 2022; acoustic testing would take place during the cruises, and a multi-frequency navigation system concept test would employ acoustic sources that are left behind. The first cruise would begin on September 14, 2022, and subsequent cruises would occur in the summer and/or fall of 2023, 2024, and 2025. The Proposed Action includes multiple scientific objectives that support the Arctic and Global Prediction Program. The Proposed Action constitutes the development of a new system under the ONR Arctic Mobile Observing System (AMOS) project, involving very-low-, low-, and mid-frequency transmissions (35 Hertz [Hz], 900 Hz, and 10 kilohertz [kHz] respectively). The AMOS project would utilize acoustic sources and receivers to provide a means of performing under-ice navigation for gliders and unmanned undersea vehicles (UUVs). This would allow for the possibility of year-round scientific observations of Arctic environmental phenomena. As an environment particularly affected by climate change, year-round observations under a variety of ice conditions are required to study the effects of this changing environment for military readiness, as well as the implications of environmental change to humans and animals. Very-low frequency technology allows for a larger range for the navigation system. The technology also has the potential to allow for development and use of navigational systems that would not be heard by some marine mammal species, and therefore would be less impactful overall. The ARA Proposed Action now consists primarily of activities under the AMOS project, but as future research could take place under different funding sources with different project names, ONR applies the term ARA to all activities described herein.

The Proposed Action would occur within the Study Area, which includes both the U.S. Exclusive Economic Zone (EEZ), the global commons, and the Canadian EEZ. The Proposed Action would primarily occur in the Beaufort Sea, but the analysis considers the drifting of active sources on buoys into the eastern portion of the Chukchi Sea. The closest point of the Study Area to the Alaska coast is 110 nautical miles (nm; 204 kilometers [km]). To allow for the equipment drift or the need to navigate around ice, small areas of the Canadian EEZ are also included in the Study Area; the appropriate permission for conducting scientific research in the Canadian EEZ would be obtained from Canada in the form of a Marine Scientific Research (MSR) permit. The AMOS project would include use of acoustic sources moored into fixed locations. The anticipated movement of any drifting sources is included in the analysis.

Purpose of and Need for the Proposed Action

The primary purpose of these activities is to test the feasibility of using a field of active acoustic sources as navigation aids to unmanned vehicles collecting oceanographic and ice data under ice-covered conditions.

The Naval need for this scientific research relates to environmental characterization in support of combat capable forces ready to deploy worldwide in accordance with Title 10 United States Code (U.S.C.) section 8062, and to support the aims of the Arctic Research and Policy Act (15 U.S.C. sections 4101 et seq.). For the Arctic, this consists of potential submarine and surface ship operations with active sonar for anti-submarine warfare and submarine/surface ship force protection. The characterization of the potential Arctic battlespace, given the changes in water properties and ice cover, is critical to performance predictions for active and passive acoustic systems. The year-round characterization of the arctic environment by use of leave-behind sources and autonomous vehicles, and the research projects

are geared toward building multiple sources transmitting intermittently to allow vehicles to transmit under the ice.

Alternatives Considered

Alternatives were developed for analysis based upon the following reasonable alternative screening factors: operation of the navigation system in areas where there will be total ice coverage during a portion of the year; acoustic source transmissions to allow for navigation of unmanned vehicles in ice-covered areas; and waters of appropriate depths to meet the scientific objectives of the Proposed Action and the need of the Navy to have the navigation systems needed to support operations far from shore. The experiment design needed to be in a location accessible during research cruises of reasonable length (5-6 weeks), while maintaining a distance that would ensure that there were no effects in subsistence hunting areas. The determination of no effect on subsistence hunting is included in the Incidental Harassment Authorization (IHA) application. The Navy is considering two action alternatives that meet the purpose and need for the Proposed Action and a No Action Alternative. Alternative 1 would be to conduct all the scientific research described in the Proposed Action, excluding the use of the VLF sources. This meets the core scientific objectives of the research projects described in Section 1.3 (Purpose and Need), particularly the use of acoustic sources as navigation aids to unmanned vehicles in the basin. Under Alternative 2 (Preferred Alternative), all of the scientific research described in the Proposed Action would occur, including year-round use of the VLF sources. This would allow for the navigation system to operate over a larger area. All acoustic sources would typically be recovered and reactivated on a yearly basis.

Under the No Action Alternative, ONR sponsored research as detailed above would not occur between 2022 and 2025. ONR would retrieve currently deployed acoustic sources to the extent this can be done under the existing supplemental OEA and regulatory authorizations, covering Arctic research from 2018 to present (U.S. Department of the Navy 2017c). Acoustic sources that cannot be recovered would cease transmitting at the end of the period of the current supplemental OEA and regulatory documents, which end October 2022.

Summary of Environmental Resources Evaluated in the OEA

Executive Order (E.O.) 12114 and Navy instructions for implementing E.O. 12114, specify that an Overseas Environmental Assessment (OEA) should address those resource areas potentially subject to harm. In addition, the level of analysis should be commensurate with the anticipated level of environmental harm.

The following resource areas are addressed in this OEA: physical resources (atmospheric temperature; bathymetry and seafloor sediments; currents, circulation, and water masses; water quality; and sea ice) and biological resources (invertebrates, marine birds, fish, Essential Fish Habitat, and marine mammals). Because potential impacts were considered to be negligible or nonexistent, the following resources were not evaluated in this OEA: air quality, cultural resources, land use, visual resources, airspace, water quality, deep sea corals and coral reefs, marine vegetation, and sea turtles.

Summary of Potential Environmental Consequences of the Action Alternatives and Major Mitigating Actions

The results of the analysis indicate that none of the alternatives considered would significantly harm physical or biological resources. The Navy consulted with the National Marine Fisheries Service (NMFS) and the United States Fish and Wildlife Service (USFWS) under section 7 of the Endangered Species Act (ESA) regarding the Preferred Alternative. The Proposed Action may adversely affect ringed seals, and is not likely to adversely affect bowhead whales, bearded seals, or polar bears.

Under both Alternatives 1 and 2, some of the species protected under the Marine Mammal Protection Act (MMPA) were predicted to be exposed to acoustic stressors (non-impulsive acoustic sources and icebreaking noise) that equated to Level B harassment levels. The Navy will consult annually with NMFS to request IHAs, for the duration of the Proposed Action, for the predicted Level B exposures; a request for an IHA to cover the period from September 2022 – September 2023 has been submitted.

The Proposed Action has the potential to decrease the quality or quantity of Arctic cod Essential Fish Habitat. The Navy prepared an Essential Fish Habitat Assessment, and submitted it to NMFS, for concurrence that adverse effects may occur.

Table ES-1 provides a summary of the potential impacts to the resources associated with each of the alternative actions analyzed.

Table ES-1. Summary of Potential Harm to Resource Areas

<i>Resource Area</i>	<i>No Action Alternative</i>	<i>Alternative 1</i>	<i>Alternative 2 (Preferred Alternative)</i>
Physical Resources	No change to baseline.	The potential harm would be temporary and localized due to the minimal number of devices and the infrequency of testing activities, and soft sediment is expected to shift back as it would following a disturbance of tidal energy. No long-term increases in turbidity (sediment suspended in water) would be anticipated. The localized disturbances would not alter the function or habitat provided by marine substrates or sea ice. No significant harm due to changes in ambient noise levels would occur as a result of the Proposed Action.	The potential harm would be temporary and localized due to the minimal number of devices and the infrequency of testing activities, and soft sediment is expected to shift back as it would following a disturbance of tidal energy. No long-term increases in turbidity (sediment suspended in water) would be anticipated. The localized disturbances would not alter the function or habitat provided by marine substrates or sea ice. No significant harm due to changes in ambient noise levels would occur as a result of the Proposed Action.
Invertebrates	No change to baseline.	With standard operating procedures and mitigation measures, potential harm from the Proposed Action would be temporary and/or minimal. The Proposed Action is not expected to result in population-level impacts to invertebrates.	With standard operating procedures and mitigation measures, potential harm from the Proposed Action would be temporary and/or minimal. The Proposed Action is not expected to result in population-level impacts to invertebrates.
Marine Birds	No change to baseline.	With standard operating procedures and mitigation measures, potential harm from the Proposed Action would be temporary and/or minimal. The Proposed Action is not expected to result in population-level impacts to marine birds.	With standard operating procedures and mitigation measures, potential harm from the Proposed Action would be temporary and/or minimal. The Proposed Action is not expected to result in population-level impacts to marine birds.
Fish	No change to baseline.	With standard operating procedures, potential harm from the Proposed Action would be temporary and/or minimal. The Proposed Action is not expected to result in population-level impacts to fish.	With standard operating procedures, potential harm from the Proposed Action would be temporary and/or minimal. The Proposed Action is not expected to result in population-level impacts to fish.
Essential Fish Habitat	No change to baseline.	With standard operating procedures, potential adverse effects from the Proposed Action would be minimal.	With standard operating procedures, potential adverse effects from the Proposed Action would be minimal.

<i>Resource Area</i>	<i>No Action Alternative</i>	<i>Alternative 1</i>	<i>Alternative 2 (Preferred Alternative)</i>
Marine Mammals	No change to baseline.	With standard operating procedures and mitigation measures, potential harm from the Proposed Action would be temporary and/or minimal. The Proposed Action is not expected to result in population-level impacts to marine mammals.	With standard operating procedures and mitigation measures, potential harm from the Proposed Action would be temporary and/or minimal. The Proposed Action is not expected to result in population-level impacts to marine mammals.

Overseas Environmental Assessment
ONR Arctic Research Activities 2022-2025
Beaufort and Chukchi Seas, Arctic

TABLE OF CONTENTS

Abbreviations and Acronyms	vii
1	PURPOSE OF AND NEED FOR THE PROPOSED ACTION 1-1
1.1	Introduction..... 1-1
1.2	Location 1-1
1.3	Purpose of and Need for the Proposed Action..... 1-3
1.4	Scope of Environmental Analysis..... 1-3
1.5	Relevant Laws and Regulations 1-3
2	PROPOSED ACTION AND ALTERNATIVES 2-1
2.1	Proposed Action 2-1
2.2	Research Equipment and Platforms 2-1
2.2.1	Glider Surveys 2-1
2.2.2	Research Vessels: R/V Sikuliaq and CGC HEALY 2-2
2.2.3	Acoustic Sources 2-3
2.2.4	Drifting Oceanographic Sensors..... 2-4
2.2.5	Moored Oceanographic Sensors..... 2-5
2.2.6	On-Ice Measurement Systems 2-5
2.3	Screening Factors 2-5
2.4	Alternatives Carried Forward for Analysis..... 2-6
2.4.1	No Action Alternative..... 2-6
2.4.2	Alternative 1..... 2-6
2.4.3	Alternative 2 (Preferred Alternative) 2-6
2.5	Alternatives Considered but not Carried Forward for Detailed Analysis 2-6
3	AFFECTED ENVIRONMENT 3-1
3.1	Physical Resources..... 3-1
3.1.1	Affected Environment 3-1
3.2	Biological Resources 3-3
3.2.1	Regulatory Setting..... 3-3
3.2.2	Affected Environment 3-6
4	ENVIRONMENTAL CONSEQUENCES 4-1
4.1	Stressors Associated with the Proposed Action 4-1
4.1.1	Acoustic Stressors 4-1
4.1.2	Physical Stressors 4-4
4.1.3	Expended Materials 4-6

4.2	Physical Resources.....	4-7
4.2.1	No Action Alternative.....	4-7
4.2.2	Action Alternatives.....	4-7
4.3	Biological Resources.....	4-9
4.3.1	No Action Alternative.....	4-9
4.3.2	Action Alternatives.....	4-9
4.4	Summary of Potential Impacts to Resources.....	4-49
5	STANDARD OPERATING PROCEDURES AND MITIGATION MEASURES.....	5-1
5.1	Standard Operating Procedures.....	5-1
5.2	Mitigation Measures.....	5-1
5.3	Monitoring and Reporting.....	5-2
6	CONSISTENCY WITH OTHER FEDERAL, STATE, AND LOCAL LAWS, PLANS, POLICIES, AND REGULATIONS..	6-1
7	REFERENCES.....	7-1
8	LIST OF PREPARERS.....	8-1

List of Figures

Figure 1-1.	Arctic Study Area.....	1-2
Figure 2-1.	Example of Seagliders.....	2-2
Figure 3-1.	Arctic Ocean Circulation.....	3-5
Figure 3-3.	Average Arctic Sea Ice Extent for March (1979-2022).....	3-3
Figure 3-4.	Essential Fish Habitat for Arctic Cod.....	3-19
Figure 3-6.	Bowhead Whale Distribution in the Study Area.....	3-24
Figure 3-7.	Polar Bear At-Sea Distribution in Study Area.....	3-26
Figure 3-8.	Ringed Seal Distribution in Study Area.....	3-29
Figure 3-9.	Pacific Walrus Distribution Near the Study Area.....	3-36
Figure 4-1.	The Bayesian biphasic dose-response BRF for A) Odontocetes, B) Pinnipeds, and C) Mysticetes.....	4-29

List of Tables

Table 2-1.	Source Characteristics of Modeled Acoustic Sources for the Proposed Action.....	2-3
Table 2-2.	Parameters for De Minimis Acoustic Sources.....	2-4
Table 3-1.	Taxonomic Groups of Marine Invertebrates in the Study Area.....	3-7
Table 3-2.	Major Groups of Marine Fish in the Study Area during the Proposed Action.....	3-14
Table 3-3.	Mammals Found in the Study Area during the Proposed Action.....	3-20
Table 3-4.	Species in Marine Mammal Hearing Groups Potentially Within the Study Area.....	3-39

Table 4-1. Acoustic In-Water Criteria and Thresholds for Predicting Physiological and Behavioral Effects on Marine Mammals Potentially Occurring in the Study Area 4-28

Table 4-2. NAEMO-Calculated Marine Mammal Estimated Yearly Exposures 4-30

Table 4-3. Modeled Bins for 8/10 Ice Coverage (Full Power) Ice Breaking on CGC HEALY..... 4-35

Table 4-4. Modeled Bins for 3/10 Ice Coverage (Quarter Power) Ice Breaking on CGC HEALY..... 4-35

Table 4-5. Model-Calculated Yearly Acoustic Exposures for CGC HEALY Icebreaking 4-36

Table 4-6. Summary of Potential Impacts to Resource Areas 4-50

Table 6-1. Principal Federal and State Laws Applicable to the Proposed Action 6-1

Appendices

APPENDIX A STRESSOR MATRICES..... A-1

APPENDIX B NON-IMPULSIVE AND IMPULSIVE SOURCE MODELING B-1

Appendix Tables

Appendix Table A-1. Stressors by Activity..... A-2

Appendix Table A-2. Stressors by Resource..... A-3

Appendix Table B-1. Environmental Parameters for ARA..... B-2

Appendix Table B-2. Acoustic Injury (PTS) and Disturbance (TTS, Behavioral) Thresholds for Underwater Sounds¹ B-3

Appendix Table B-4. Predicted Annual Marine Mammal Exposures All Events (Acoustic and Icebreaking).. B-9

Appendix Figures

Appendix Figure B-1. A) The Bayesian biphasic dose-response BRF for Odontocetes. B) The Bayesian biphasic dose-response BRF for Pinnipeds C) The Bayesian biphasic dose-response BRF for Mysticetes. The blue solid line represents the Bayesian Posterior median values, the green dashed line represents the biphasic fit, and the grey represents the variance. [X-Axis: Received Level (dB re 1 μ Pa), Y-Axis: Probability of Response] B-6

Abbreviations and Acronyms

Acronym	Definition	Acronym	Definition
°C	degrees Celsius	in/s	inches per second
°E	degrees East	kg	kilogram
°F	degrees Fahrenheit	kHz	kiloHertz
°N	degrees North	km	kilometers
°W	degrees West	km/day	kilometers per day
ABR	auditory brainstem response	km ²	square kilometers
ADCP	Acoustic Doppler Current Profiler	lb	pound
AEP	auditory evoked potential	m	meters
AMOS	Arctic Mobile Observing System	m/s	meters per second
ARA	Arctic Research Activities	MBTA	Migratory Bird Treaty Act
BOEM	Bureau of Ocean Energy Management	mi ²	square miles
BRF	behavioral response function	MLLW	mean lower low water
CASS	Comprehensive Acoustic System Simulation	MMPA	Marine Mammal Protection Act
CFR	Code of Federal Regulations	MSA	Magnuson-Stevens Fishery Conservation and Management Reauthorization Act
CGC	Coast Guard Cutter	MSR	Marine Scientific Research
cm	centimeters	NAAQS	National Ambient Air Quality Standards
cm/s	centimeters per second	NAEMO	Navy Acoustic Effects Model
CTD	Conductivity Temperature Depth	Navy	Department of the Navy
dB	decibels	nm	nautical miles
dB re 1 μPa	decibels referenced to 1 micropascal	NMFS	National Marine Fisheries Service
DPS	distinct population segment	NMSDD	Navy Marine Species Density Database
E.O.	Executive Order	NTE	not to exceed
EEZ	Exclusive Economic Zone	OAML	Oceanographic and Atmospheric Master Library
EFH	Essential Fish Habitat	OEA	Overseas Environmental Assessment
ESA	Endangered Species Act	ONR	Office of Naval Research
ft	feet	psu	practical salinity unit
ft/s	feet per second	PTS	permanent threshold shift
GRAB	Gaussian Ray Bundle	R/V	Research Vessel
Hz	Hertz	RMS	root mean square
IGB	Ice Gateway Buoy	SEL	sound exposure level
IHA	Incidental Harassment Authorization	SPL	sound pressure level
in	inches		

Acronym	Definition	Acronym	Definition
TTS	temporary threshold shift		Institute
U.S.	United States	yd	yard
U.S.C.	United States Code		
USFWS	United States Fish and Wildlife Service		
UUV	Unmanned Underwater Vehicle		
VLF	very-low frequency		
WHOI	Woods Hole Oceanographic		

1 Purpose of and Need for the Proposed Action

1.1 Introduction

The Office of Naval Research's (ONR) Arctic Research Activities (ARA), the Proposed Action, would conduct scientific research in the Beaufort and Chukchi Seas from September 2022 to September 2025. This research comprises cruises that would occur annually beginning in September 2022; acoustic testing would take place during the cruises, and a multi-frequency navigation system concept test would employ sources that are left behind. The 2022 cruise would begin on September 14, 2022. The Proposed Action is a continuation of the research conducted under the OEA for ARA covering the time period 2018-2021 (U.S. Department of the Navy 2018a) and associated supplemental and regulatory documentation; the Proposed Action utilizes moored acoustic sources that have been deployed under the 2021 supplemental OEA (U.S. Department of the Navy 2021b). Coverage under the previous documentation ends in October 2022. Changes to the 2018 Proposed Action, as addressed in supplemental documentation in 2019, occurred as research objectives and participants evolved. Creation of a new OEA for 2022-2025 was motivated by the completion of some projects by the Naval Research Laboratory and the ONR Ocean Acoustics Program. The Proposed Action for 2022-2025 is a continuation of research activities by the ONR Arctic and Global Prediction Program. The Proposed Action includes the development of a new system under the ONR Arctic Mobile Observing System (AMOS) project, involving very-low-, low-, and mid-frequency transmissions.

The United States (U.S.) Department of the Navy (Navy) has prepared this Overseas Environmental Assessment (OEA) in accordance with Executive Order (E.O.) 12114, Environmental Effects Abroad of Major Federal Actions.

1.2 Location

The Proposed Action would occur within the Study Area (Figure 1-1), which includes the U.S. Exclusive Economic Zone (EEZ), the global commons, and the Canadian EEZ. The Proposed Action would primarily occur in the Beaufort Sea, but the analysis considers the drifting of active sources on buoys into the eastern portions of the Chukchi Sea. The closest point of the Study Area to the Alaska coast is 110 nautical miles (nm; 204 kilometers [km]). To allow for the equipment drift or the need to navigate around ice, small areas of the Canadian EEZ are included in the Study Area; the appropriate permission for conducting scientific research in the Canadian EEZ would be obtained from Canada in the form of a Marine Scientific Research (MSR) permit. The map shows the positions of fixed sources. The anticipated movement of drifting sources is included in the analysis. Additional details regarding the specific experiments, timeframes, and research are further detailed below in Section 2.1.




Figure 1-1. Arctic Study Area

1.3 Purpose of and Need for the Proposed Action

The primary purpose of these activities is to test the feasibility of using a field of active acoustic sources as navigation aids to unmanned vehicles collecting oceanographic and ice data under ice-covered conditions. The need for this scientific research relates to environmental characterization in support of combat capable forces ready to deploy worldwide in accordance with Title 10 United States Code (U.S.C.) section 8062, and to support the aims of the Arctic Research and Policy Act (15 U.S.C. sections 4101 et seq.). For the Arctic, this consists of potential submarine and surface ship operations with active sonar for anti-submarine warfare and submarine/surface ship force protection. The characterization of the potential Arctic battlespace, given the changes in water properties and ice cover, is critical to performance predictions for active and passive acoustic systems. The year-round characterization of the arctic environment requires characterization of the environment by leave-behind sources and autonomous vehicles, and the research project is geared toward building multiple sources transmitting intermittently to allow vehicles to navigate under the ice. The Navy's strategic objective for the Arctic Region, according to the U.S. Navy Strategic Blueprint for the Arctic (U.S. Department of the Navy 2021a) is to apply Naval power as the U.S. continues to prepare for a more navigable Arctic Region over the next two decades.

1.4 Scope of Environmental Analysis

This OEA includes an analysis of potential environmental harm associated with the Action Alternatives and the No Action Alternative. The environmental resource areas analyzed in this OEA include the following: physical environment (atmospheric temperature; bathymetry and seafloor sediments; currents, circulation, and water masses; water quality; and sea ice) and biological resources (invertebrates, marine birds, fish, Essential Fish Habitat [EFH], and marine mammals).

1.5 Relevant Laws and Regulations

The Navy has prepared this OEA based upon federal statutes, regulations, and policies that are pertinent to the implementation of the Proposed Action, including the following:

- Arctic Research and Policy Act (15 U.S.C. sections 4101-4111)
- Endangered Species Act (ESA) (16 U.S.C. section 1531 et seq.)
- Magnuson-Stevens Fishery Conservation and Management Reauthorization Act (MSA) (16 U.S.C. section 1801 et seq.)
- Marine Mammal Protection Act (MMPA) (16 U.S.C. section 1361 et seq.)
- Migratory Bird Treaty Act (MBTA) (16 U.S.C. sections 703-712)
- E.O. 12114, Environmental Effects Abroad of Major Federal Actions

A description of the Proposed Action's consistency with these laws, policies, and regulations, as well as the names of regulatory agencies responsible for their implementation, is presented in Table 6-1.

This page intentionally left blank

2 Proposed Action and Alternatives

2.1 Proposed Action

ONR's Proposed Action is to conduct scientific research in the Beaufort and Chukchi Seas from September 2022 to September 2025. This research comprises cruises that would occur annually beginning in September 2022. Acoustic testing would take place during the cruises, and a multi-frequency navigation system concept test would employ sources left behind. The first cruise would begin on September 14, 2022. The Proposed Action includes multiple scientific objectives that support the Arctic and Global Prediction Program. The Proposed Action constitutes the development of a new system under the ONR AMOS project, involving very-low-, low-, and mid-frequency transmissions (35 Hertz [Hz], 900 Hz, and 10 kilohertz [kHz] respectively). The AMOS project would utilize acoustic sources and receivers to provide a means of performing under-ice navigation for gliders and unmanned undersea vehicles (UUVs). This would allow for the possibility of year-round scientific observations of Arctic environmental phenomena. As an environment particularly affected by climate change, year-round observations under a variety of ice conditions are required to study the effects of this changing environment for military readiness, as well as the implications of environmental change to humans and animals. Very-low frequency technology is an important method of observing ocean warming, and the continued development of these types of acoustic sources would allow for characterization of larger areas. The technology also has the potential to allow for development and use of navigational systems that would not be heard by some marine mammal species, and therefore would be less impactful overall. The ARA Proposed Action consists primarily of activities under the AMOS project, but future research could take place under different funding sources with different project names under the umbrella of ARA.

The Proposed Action would occur within the Beaufort and Chukchi Seas (Figure 1-1). The Proposed Action would primarily occur in the Beaufort Sea, but the analysis considers the drifting of active sources on buoys into the eastern portion of the Chukchi Sea. The closest point of the Study Area to the Alaska coast is 110 nm (204 km).

2.2 Research Equipment and Platforms

Below are the descriptions of the equipment and platforms that would be deployed at different times during the Proposed Action.

2.2.1 Glider Surveys

Glider surveys are proposed for the September 2022 research cruise. All gliders would be recovered. Some may be recovered by Research Vessel (R/V) Sikuliaq during the September 2022 cruise, but all of the remainder would be recovered during the later cruises.

Long-endurance, autonomous Seagliders (Figure 2-1) are intended for use in extended missions in ice-covered waters. Gliders are buoyancy-driven, equipped with satellite modems providing two-way communication, and are capable of transiting to depths of up to 3,280 feet (ft; 1,000 meters [m]). Gliders would collect data in the area of the shallow water sources and moored sources, moving at a speed of 0.25 meters per second (m/s; 23 kilometers per day [km/day]). A combination of recent advances in Seaglider technology would provide full-year endurance. When operating in ice-covered waters, gliders navigate by trilateration (the process of determining location by measurement of distances, using the geometry of circles, spheres or triangles) from moored acoustic sound sources (or dead reckoning should navigation signals be unavailable). Hibernating gliders would continue to track

their position, waking to reposition should they drift too far from their target region. Gliders would measure temperature, salinity, dissolved oxygen, rates of dissipation of temperature variance (and vertical turbulent diffusivity), and multi-spectral downwelling irradiance.



Figure 2-1. Example of Seagliders

2.2.2 Research Vessels: R/V Sikuliaq and CGC HEALY

The R/V Sikuliaq would perform the research cruise in September 2022. The vessel would conduct testing of acoustic sources during a cruise, as well as leave sources behind to operate as a year-round navigation system observation. The ships to be used for future cruises are yet to be determined. The most probable additional option would be the Coast Guard Cutter (CGC) HEALY, so that ship is described here.

The R/V Sikuliaq has a maximum speed of approximately 12 knots with a cruising speed of 11 knots (University of Alaska Fairbanks 2014). The R/V Sikuliaq is not an ice breaking ship, but it is an ice-strengthened ship. CGC HEALY is an icebreaking ship; it travels at a maximum speed of 17 knots with a cruising speed of 12 knots (United States Coast Guard 2013), and a maximum speed of 3 knots when traveling through 3.5 ft (1.07 m) of sea ice (Murphy 2010).

The R/V Sikuliaq, CGC HEALY, or any other vessel operating a research cruise associated with the Proposed Action may perform the following activities during their research cruises:

- Deployment of moored and/or ice-tethered passive sensors (oceanographic measurement devices, acoustic receivers);
- Deployment of moored and/or ice-tethered active acoustic sources to transmit acoustic signals;
- Deployment of UUVs;
- Deployment of drifting buoys, with or without acoustic sources; or,
- Recovery of equipment.

2.2.3 Acoustic Sources

2.2.3.1 Moored/Drifting Acoustic Sources

Active acoustic sources would be lowered from a cruise vessel while stationary, deployed on gliders and UUVs, or deployed on fixed moorings. During the September 2022 and subsequent cruises, acoustic sources could be deployed from the ship for intermittent testing of the system components. The total amount of active source testing for ship-deployed sources used during a cruise would be 120 hours. The testing would take place in the vicinity of the source locations in Figure 1-1, with UUVs running tracks within the designated box. During this testing, 35 Hz, 900 Hz and 10 kHz sources would be employed.

Up to seven fixed acoustic navigation sources, transmitting at 900 Hz, would remain in place for a year at the locations given in Figure 1-1. These moorings would be anchored on the seabed and held in the water column with subsurface buoys. All sources would be deployed by shipboard winches, which would lower sources and receivers in a controlled manner. Anchors would be steel “wagon wheels” typically used for this type of deployment. Up to two very low frequency (VLF) sources, transmitting at 35 Hz, would be deployed in a similar manner. These sources would be deployed in two of the three VLF source positions in Figure 1-1. Two Ice Gateway Buoys (IGB) would also be configured with active acoustic sources at 900 Hz and 10 kHz.

Moored source transmits would be offset by 15 minutes from each other (i.e., sources would not be transmitting at the same time). All navigation sources would be recovered. Acoustic parameters for active acoustic sources are described in Table 2-1.

Table 2-1. Source Characteristics of Modeled Acoustic Sources for the Proposed Action

<i>Platform</i>	<i>Acoustic Source</i>	<i>Purpose/Function</i>	<i>Frequency</i>	<i>Signal Strength (dB re 1µPa @1 m)</i>	<i>Bandwidth</i>	<i>Pulse Width/Duty Cycle</i>
REMUS 600 UUV	WHOI Micro-modem	Acoustic communications	900-950 Hz	NTE 180 dB by system design limits	50 Hz	5 pings/hour with 30 sec pulse length.
	UUV/ WHOI Micro-modem	Acoustic communications	8-14 kHz	NTE 185 dB by system design limits	5 kHz	10% average duty cycle, with 4 sec pulse length
IGB (drifting)	WHOI Micro-modem	Acoustic communications	900-950 Hz	NTE 180 dB by system design limits	50 Hz	Transmit every 4 hours, 30 sec pulse length
	WHOI Micro-modem	Acoustic communications	8-14 kHz	NTE 185 dB by system design limits	5 kHz	Typically receive only. Transmit is very intermittent.
Mooring	WHOI Micro-modem	Acoustic Navigation	900-950 Hz	NTE 180 dB by system design limits	50 Hz	Transmit every 4 hours, 30 sec pulse length
	VLF	Acoustic Navigation	35 Hz	NTE 190 dB	6 Hz	Up to 4 times per day, 10 minutes each.

Note: dB re 1 µPa at 1 m= decibels referenced to 1 micropascal at 1 meter; NTE= not to exceed; WHOI= Woods Hole Oceanographic Institution

2.2.3.2 De minimis Sources

De minimis sources have the following parameters: low source levels, narrow beams, downward directed transmission, short pulse lengths, frequencies above (outside) known marine mammal hearing ranges, or some combination of these factors (U.S. Department of the Navy 2018b). Additionally, any sources 200 kHz or above in frequency and/or 160 dB or below in source level are automatically considered *de minimis*. Sources 200 kHz or above are considered outside of marine mammal hearing ranges. Assuming spherical spreading for a 160 decibels referenced to 1 micropascal (dB re 1 μ Pa) source, the sound would attenuate to less than 140 dB within 32 ft (10 m) and less than 120 dB within 328 ft (100 m) of the source. Ranges would be even shorter for a source less than 160 dB re 1 μ Pa source level. All of the sources described in this section are considered *de minimis*. Since they are not expected to have effects on marine mammals, *de minimis* sources are not quantitatively analyzed. Qualitative analysis is performed when special circumstances (i.e., unusual method of usage, enclosed environment) dictate.

The following are planned *de minimis* sources that would be used during the Proposed Action: Woods Hole Oceanographic Institute (WHOI) micromodem, Acoustic Doppler Current Profilers (ADCPs), ice profilers, and additional sources below 160 dB re 1 μ Pa used during towing operations. ADCPs may be used on moorings. Ice-profilers measure ice properties and roughness. The ADCPs and ice-profilers would all be above 200 kHz and, therefore, out of marine mammal hearing ranges, with the exception of the 75 kHz ADCP which has the characteristics and *de minimis* justification listed in Table 2-2. They may be employed on moorings or UUVs.

A WHOI micromodem would be employed during the leave behind period. During this period, the micromodem would be used for very intermittent communication with vehicles to communicate vehicle status for safety of navigation purposes, and it is treated as *de minimis* while employed in this manner.

Table 2-2. Parameters for De Minimis Acoustic Sources

<i>Source Name</i>	<i>Frequency Range (kHz)</i>	<i>Sound Pressure Level (dB re 1 μPa at 1 m)</i>	<i>Pulse Length (seconds)</i>	<i>Duty Cycle (Percent)</i>	<i>De minimis Justification</i>
ADCP	>200, 150, or 75	190	<0.001	<0.1	Very low pulse length, narrow beam, moderate source level
Nortek Signature 500 kHz Doppler Velocity Log	500	214	<0.1	<13	Very high frequency
CTD Attached Echosounder	5-20	160	0.004	2	Very low source level

Note: CTD = Conductivity Temperature Depth

2.2.4 Drifting Oceanographic Sensors

Observations of ocean-ice interactions require the use of sensors that are moored and embedded in the ice. For the Proposed Action, it will not be required to break ice to do this, as deployments can be performed in areas of low ice coverage or free floating ice. Sensors would be deployed within a few dozen meters of each other on the same ice floe. Three types of sensors would be used: autonomous

ocean flux buoys, Integrated Autonomous Drifters, and ice-tethered profilers. The autonomous ocean flux buoys measure oceanographic properties just below the ocean-ice interface. The autonomous ocean flux buoys would have ADCPs and temperature chains attached, to measure temperature, salinity, and other ocean parameters the top 20 ft (6 m) of the water column. Integrated Autonomous Drifters would have a long temperature string extending down to 656 ft (200 m) depth and would incorporate meteorological sensors, and a temperature string to estimate ice thickness. The ice-tethered profilers would collect information on ocean temperature, salinity and velocity down to 820 ft (250 m) depth.

Up to 20 Argo-type autonomous profiling floats may be deployed in the central Beaufort Sea. Argo floats drift at 4,921 ft (1,500 m) depth, profiling from 6,562 ft (2,000 m) to the sea surface once every 10 days to collect profiles of temperature and salinity.

2.2.5 Moored Oceanographic Sensors

Moored sensors would capture a range of ice, ocean, and atmospheric conditions on a year-round basis. These would be bottom anchored, sub-surface moorings measuring velocity, temperature, and salinity in the upper 1,640 ft (500 m) of the water column. The moorings also collect high-resolution acoustic measurements of the ice using the ice profilers described above. Ice velocity and surface waves would be measured by 500 kHz multibeam sonars from Nortek Signatures.

2.2.6 On-Ice Measurement Systems

On-ice measurement systems would be used to collect weather data. These would include an Autonomous Weather Station and an Ice Mass Balance Buoy. The Autonomous Weather Station would be deployed on a tripod; the tripod has insulated foot platforms that would be frozen into the ice. The system would consist of an anemometer, humidity sensor, and pressure sensor. The Autonomous Weather Station also includes an altimeter that is *de minimis* due to its very high frequency (200 kHz). The Ice Mass Balance Buoy is a 20 ft (6 m) sensor string, which is deployed through a 2 inch (in; 5 centimeters [cm]) hole drilled into the ice. The string is weighted by a 2.2 pound (lb) (1 kilogram [kg]) lead weight, and is supported by a tripod. The buoy contains a *de minimis* 200 kHz altimeter and snow depth sensor. Autonomous Weather Stations and Ice Mass Balance Buoys will be deployed, and will drift with the ice, making measurements, until their host ice floes melt, thus destroying the instruments (likely in summer, roughly one year after deployment). After the on-ice instruments are destroyed they cannot be recovered, and would sink to the seafloor as their host ice floes melted.

2.3 Screening Factors

E.O. 12114's implementing regulations provide guidance on the consideration of alternatives to a federally proposed action and require rigorous exploration and objective evaluation of reasonable alternatives. Only those alternatives determined to be reasonable and meet the purpose and need require detailed analysis.

Potential alternatives that meet the purpose and need were evaluated against the following screening factors:

- Geographic sampling in deep water areas where there will be total ice coverage during a portion of the year.
- Acoustic source transmissions in deep water to allow for navigation of unmanned vehicles in ice-covered areas.
- Waters of appropriate depths to meet the scientific objectives of the Proposed Action (e.g., deep water sources require specific depths in order to appropriately measure duct propagation).

2.4 Alternatives Carried Forward for Analysis

Based on the reasonable alternative screening factors and meeting the purpose and need for the proposed action, two Action Alternatives (plus a No Action Alternative) were identified and will be analyzed within this OEA.

2.4.1 No Action Alternative

Under the No Action Alternative, ARA would not occur. ONR would retrieve currently deployed acoustic sources to the extent this can be done under the existing supplemented 2018 OEA and regulatory authorizations (U.S. Department of the Navy 2018a). Without icebreaking, it is possible that not all sources will be recovered, but the sources will no longer be active after the end period of the previous supplemental OEA, which ends October 2022. The No Action Alternative would not meet the purpose and need for the Proposed Action; however, the No Action Alternative is carried forward for analysis in this OEA and provides a baseline for measuring the environmental consequences of the action alternatives.

2.4.2 Alternative 1

Alternative 1 would be to conduct all the scientific research described in the Proposed Action, excluding the use of the VLF sources. This meets the core scientific objectives of the research projects described in Section 1.3 (Purpose and Need), particularly the use of acoustic sources as navigation aids to unmanned vehicles.

2.4.3 Alternative 2 (Preferred Alternative)

Under Alternative 2 (Preferred Alternative), all of the scientific research described in the Proposed Action would occur, including year-round use of the VLF sources. Sources would typically be recovered and reactivated on a yearly basis. This alternative would meet the core scientific objectives of the research projects described in Section 1.3 (Purpose and Need), particularly in the use of acoustic sources as navigation aids to unmanned vehicles, and also allow the system to operate over long ranges and have some acoustic transmissions that would be inaudible to certain marine species.

2.5 Alternatives Considered but not Carried Forward for Detailed Analysis

Other locations were considered, but specific water depths are required to meet the goals of creating an underwater navigation system. The experiments must be left out long term to collect the data necessary for proper acoustic propagation analysis under open-water, marginal ice, and ice-covered conditions. The environment is complex and variable, and models to successfully simulate acoustic conditions need to be developed – hence the need for at-sea observations. There are no reasonable surrogate

environments that can be used to observe the various phenomena associated with unmanned vehicle navigation and acoustic propagation in the Arctic. The proposed location was selected due to the substantial distance from areas in which marine mammal hunting by Alaska Natives takes place.

3 Affected Environment

This chapter presents a description of the environmental resources and baseline conditions that could be affected from implementing any of the alternatives.

All potentially relevant environmental resource areas were initially considered for analysis in this OEA. In compliance with E.O. 12114, the discussion of the affected environment (i.e., existing conditions) focuses only on those resource areas potentially subject to harm. Additionally, the level of detail used in describing a resource is commensurate with the anticipated level of potential environmental harm. This section includes physical resources and biological resources.

The potential harm to the following resource areas are considered to be negligible or non-existent, so they will not be analyzed in this OEA:

Air Quality: The Proposed Action is substantially outside of 12 nm, attainment status is not applicable, and the Clean Air Act National Ambient Air Quality Standards (NAAQS) do not apply. Additionally, all coastal Alaska boroughs and counties are classified as attainment areas for criteria pollutants (40 Code of Federal Regulations [CFR] § 81.302). Attainment areas are areas that meet the NAAQS for specific pollutants. Under the Clean Air Act, only nonattainment areas are required to limit and act to decrease emissions below the NAAQS.

Cultural Resources: There are no known cultural resources within the Study Area.

Land Use: There would be no land use as part of the Proposed Action.

Visual Resources: The use of research vessels in the Arctic is common, and the limited use of the vessels would not harm visual resources. The general project area is outside of the viewshed of anyone on land.

Airspace: There would be no use of airspace as part of the Proposed Action.

Water Quality: The Proposed Action would not have any discharges or chemical interaction with the water.

Deep Sea Corals and Coral Reefs: No deep sea corals or coral reefs are present in the Study Area.

Marine Vegetation: The marine vegetation in the Study Area is made up of free-floating diatoms and plankton, which would not be harmed by objects deployed on the sea ice, in the water column, or by acoustic stressors.

Sea Turtles: No sea turtles would be present in the Study Area.

3.1 Physical Resources

This discussion of physical resources includes atmospheric temperature, bathymetry, currents, circulation and water masses, water quality, and sea ice. This section discusses the physical characteristics of the Study Area; biological resources are addressed in Section 3.2. There are no specific regulations that apply to these resources.

3.1.1 Affected Environment

The following discussions provide a description of the existing conditions for the physical environment of the Arctic in the Study Area.

3.1.1.1 Atmospheric Temperature

The Earth's climate has warmed approximately 1.1 degrees Fahrenheit (°F; 0.6 degrees Celsius [°C]) over the past 100 years with two main periods of warming occurring from 1910 to 1945 and from 1976 to present day (Overland et al. 2014; Walther et al. 2002). The period from October 2020-September 2021 was the seventh warmest on record over Arctic lands (beginning in 1900 with surface air temperature anomalies 34 °F [1.1°C] above the 1981-2010 mean); the warmest autumn (October-December) and fourth warmest spring (April-June) in the Arctic strongly contributed to the annual temperature anomaly (Ballinger et al. 2021). Temperature trends in the Arctic exhibit regional and annual variability (Maxwell 1997; Symon et al. 2005); however, a general warming trend has been observed since the late 1970s. Warming air temperatures have played a major role in the observed increase in permafrost temperatures around the Arctic rim, earlier spring snowmelt, reduced sea ice, widespread glacial retreat, increases in river discharge into the Arctic Ocean, and an increase in greenness of Arctic vegetation (Overland et al. 2014). Arctic atmospheric circulation is a complicated system, though air moves west to east across the Study Area and into the Canadian Archipelago and mainland (Hudson et al. 2001). Based on approximately nine months of data (including those months during which the Proposed Action would occur) from a 2014 model, the wind speed measured at a point in the Beaufort Sea south of the Study Area averaged 14.6 feet per second (ft/s; 6.83 m/s) (Naval Oceanographic Office 2014). The climatologic, hydrologic, and biologic subsystems of the Arctic are highly interconnected, and, thus, they cannot be easily isolated for discussion (Hinzman et al. 2005).

3.1.1.2 Bathymetry and Seafloor Sediments

The Beaufort Sea has a narrow, shallow shelf along the north coast of Alaska, with a width of less than 80 nm (148 km) at any given point (Dome Petroleum Ltd. et al. 1982). Off the coast of Canada, the shelf is broader, and depths of 33 feet (ft; 10 meters [m]) or less can be found up to 16 nm (30 km) from shore (Wilkinson et al. 2009). The average depth within the shelf of the Beaufort Sea is less than 213 ft (65 m) (Dome Petroleum Ltd. et al. 1982). The continental slope in this area drops steeply to the Canada Basin. The Canada Basin, which extends north into the Arctic Ocean and is bordered to the west by the Mendeleev Ridge, averages a depth of about 11,811 ft (3,600 m) (Wilkinson et al. 2009). Seafloor sediments in this deep water basin are typically muddy (Bluhm et al. 2011). Based on visual evaluation by Bluhm et al. (2005), the seafloor within the Canada Basin is composed of very fine, silty sediment over a thick clay layer. Coastal erosion supplies an estimated 7 million tons of sediment each year near shoreline areas of the Beaufort Sea. While erosion is an important local source of sediments, the relative contribution of coastal erosion to sediment loading in the Beaufort Sea is minor compared to sediments originating from the Mackenzie River, which reaches approximately 130 million tons of sediment each year (Carmack and Macdonald 2002).

The Study Area also encompasses the majority of the Chukchi Plateau, which lies to the west of the Canada Basin in the Chukchi Sea. The eastern margin of the Chukchi Plateau, the Northwind Ridge, is also contained in the Study Area. It runs parallel to the northward trend of the plateau and is separated from the rest of the plateau by the Northwind Basin, an abyssal plain that reaches depths of 11,482 ft (3,500 m) (Nuttall 2005). The Northwind Ridge is bounded on the eastern side to the Canada Basin by a steep, downward slope. Due to the escarpment, the slope contains a large amount of rock substrate, but clayey mud forms the predominant sediments (Mayer and Armstrong 2012).

The benthic communities of the Beaufort Sea are comprised of benthic macroalgae, macrophytic algae, infaunal invertebrates (living within the sediment), and epifaunal fish and invertebrates (living on the seafloor) (Minerals Management Service 1991). The biomass and diversity of benthic communities generally increase with depth within the inshore or intermediate zone, except from 49 to 82 ft (15 to

25 m) depth, which is an area where the most intensive ice-gouging occurs (Minerals Management Service 1991). Soft sediments dominate the continental shelves of the Beaufort and Chukchi Seas. This sediment is largely a combination of muds, sands, and gravels—substrates that support high densities of invertebrates (Holland-Bartels et al. 2011). Benthic macroalgae requires rocky substrate for attachment, which is rare within the Study Area. Sediments in the Study Area of the Beaufort Sea consist mostly of gravel and sand, and those in the deep Canada Basin, in particular, are mainly fine-grained clay and silt (Hong et al. 2012). There are no known areas with hard substrate suitable for attachment by kelps and macroalgae within the Study Area.

3.1.1.3 Currents, Circulation, and Water Masses

The processes governing water currents and circulation into and out of the Beaufort and Chukchi Seas are complex. Cold, saline water (averaging about 32.5 practical salinity units [psu]) enters the Bering Strait from the Pacific Ocean (Woodgate et al. 2005). Because the Bering Strait is a narrow, shallow passageway that measures only 46 nm (85 km) wide and 164 ft (50 m) deep, it is only an inflow point (Woodgate 2012). On the Atlantic side, both inflow (through the Barents Sea and Fram Strait) and outflow movement of water occurs (Woodgate 2012).

Currents within the Beaufort Gyre are variable and depend on multiple factors, including wind speed, presence of eddies, and the value of the Arctic Oscillation. These factors come together to affect the overall velocity of the waters as they move throughout the Arctic Ocean, and they can make predicting the velocity of the currents difficult. While subsurface velocities have been measured from ice camps historically, the most comprehensive studies are often of short duration (Plueddemann et al. 1998). Plueddemann et al. (1998) used an Ice-Ocean Environmental Buoy frozen into Arctic pack ice approximately 130 nm (241 km) north of Prudhoe Bay, Alaska, to take long-term measurements of meteorological and oceanographic variables in the Arctic. That study concluded that the ice drift within the Beaufort Gyre ranged from approximately 0.4 to 2 inches per second (in/s; 1 to 5 centimeters per second [cm/s]) (Plueddemann et al. 1998). Ice Ocean Environmental Buoy deployment within the Beaufort Gyre also has been used to study various physical properties of Arctic eddies. O'Brien et al. (2013) used moorings with sequential sediment traps to investigate downward sediment flux in the Canada Basin. The sediment traps measured water current speed at multiple depths. From the surface to a depth of 272 ft (83 m), observed velocities were typically between 0 and 4 in/s (0 and 10 cm/s), although they could climb to 16 in/s (40 cm/s) in the event of encounter with an eddy. The Beaufort Gyre expands and contracts based on the state of Arctic Oscillation; under high Arctic Oscillation conditions, the Beaufort Gyre will contract (Woodgate 2012).

In the Arctic, areas of ice-cover usually have a surface mixed layer 16 to 32 ft (5 to 10 m) deep. In ice-free regions, which have increased over time, this mixed layer, driven by winds, can be more than twice as deep (Rainville et al. 2011). In most of the western Arctic (also referred to as the Canada Basin), Pacific Waters are found below this mixed layer. Pacific Winter Waters are indicated by a deep minimum temperature around freezing at depths of about 320 to 492 ft (100 to 150 m) (Woodgate 2012). Shallower temperature maxima, probably formed locally by solar heating, are observed in some regions (Jackson et al. 2010; Shimada et al. 2001). Below the Pacific Water, Atlantic Water forms a temperature maximum (up to 33.8 °F [1 °C]) at depths of around 640 to 1,312 ft (200 to 400 m). These are called Fram Strait Branch Waters since they come mainly from the Fram Strait inflow (Rudels et al. 1994), although some influence is likely from the Barents Sea (Rudels et al. 2000; Woodgate et al. 2001). Below the Fram Strait Branch Waters, temperatures decrease, and an inflexion point in temperature-salinity marks waters of mainly Barents Sea origin (Rudels et al. 1994; Smith et al. 1999). Throughout the Arctic, a cold halocline layer provides a density barrier, trapping Atlantic Water heat at depth away from the

ice. Arctic Bottom Water occurs at depths greater than 2,953 ft (900 m), and this water has temperatures from 30.6 to 30.4 °F (-0.8 to -0.9 °C) and salinities from 34 to 35 psu (Woodgate 2012). Upwelling and eddies allow for increased mixing of water both by currents and by mixing of water layers containing different temperatures and salinities (Weingartner et al. 2008).

In the Beaufort Sea, the Alaska Shelf-Slope Front stretches along the north coast of Alaska from Point Barrow to the Mackenzie Delta by the Canadian Border. This front is a “hot spot” of activity where marine life gathers, including mammals and sea birds. Additionally, this is the site of the Cape Bathurst Polynya (an area of open sea surrounded by pack ice) (Belkin and Cornillon 2007). In the Arctic Ocean, the observation of fronts is hampered by perennial ice cover that prevents satellite remote sensing in the Arctic Basin. Data collected from drifting stations and submarines has revealed a major front separating Atlantic waters from Pacific waters. Until the mid-1990s, this front was located over the Lomonosov Ridge, but is now along the Mendeleev-Alpha Ridge (Belkin and Cornillon 2007).




Figure 3-1. Arctic Ocean Circulation

3.1.1.4 Water Quality

The high Arctic waters (a term used to describe barren polar areas) consist of water with relatively low nutrient loads. At the end of the winter, a burst of primary productivity occurs under the ice when light levels become sufficiently high and nutrients are released from the ice. This surge of nutrients includes nitrogen (as ammonium, nitrite, and nitrate), phosphorus (as phosphate), iron, and other elements, which would be either grazed upon and move through the food chain, or sink to the bottom and incorporate into bottom sediments (Vancoppenolle et al. 2013). In polar waters, nutrient concentrations undergo seasonal depletion in surface waters due to photosynthesis during spring/summer and renewal during winter when photosynthesis stops (Whitledge et al. 2008).

3.1.1.5 Sea Ice

3.1.1.5.1 Arctic Sea Ice Regime

Sea ice is frozen seawater that floats on the surface of the ocean, covering millions of square miles. Sea ice that persists year after year, surviving at least one summer melt season, is known as multiyear ice. Sea ice forms and melts with polar seasons and affects both human activity and biological habitat (Jeffries et al. 2014). Arctic sea ice plays a crucial role in Northern Hemisphere climate and ocean circulation, and it is thought to play an even more crucial role in regulating climate than Antarctic sea ice (National Snow and Ice Data Center 2007; Serreze et al. 2003).

Sea ice directly impacts coastal areas and broadly affects surface reflectivity, ocean currents, water cloudiness, humidity, and the exchange of heat and moisture at the ocean's surface. Since sea ice reflects the sun's heat, when ice retreat is greater and there is more open ocean, more of the sun's heat is absorbed, increasing the warming of the water (Timmermans and Proshutinsky 2014).

3.1.1.5.2 Sea Ice Extent

Though the record of sea ice extent dates as far back as 1900 in the Northern Hemisphere, the most complete record of sea ice is provided by microwave satellites, which have routinely and accurately monitored sea ice extent since 1979 (Jeffries et al. 2014; Timmermans and Proshutinsky 2014). Annually, sea ice extent is at its maximum in March, representing the end of winter, and is at its minimum in September (Jeffries et al. 2014). Figure 3-2 demonstrates minimum and maximum 2021 ice extent in comparison to the median minimum and maximum extents from 1981 to 2010.

September 2012 remains the record low minimum ice extent of 1.3 million square miles (mi^2 ; 3.4 million square kilometers [km^2]) (National Snow and Ice Data Center 2022). Data from 2016 reveals a minimum extent of 1.6 million mi^2 (4.14 million km^2), tied for the second lowest minimum extent with September 2007. In September 2007, the sea ice recession was so vast that the Northwest Passage completely opened up for the first time on record (National Snow and Ice Data Center 2007). Data from 2021 reveals a minimum extent of 2.93 million mi^2 (4.72 million km^2), tied for the seventh lowest extent on record.




Figure 3-2. Average Arctic Sea Ice Extent in March and September

The age of the sea ice is an indicator for its physical properties, including surface roughness, melt pond coverage, and ice thickness. Older ice tends to be thicker and, thus, more resilient to changes in atmospheric and oceanic forcing than younger ice. The distribution of ice of different ages illustrates the extensive loss in recent years of the older ice types (Maslanik et al. 2011). Current ice coverage favors first-year ice, or ice that has not survived a melt season. This is the thinnest type of ice. Sea ice has also been experiencing later freeze-up than usual and earlier ice melt, leading to a declining trend in multiyear ice (Overland and Wang 2013). The age of the ice can be determined using satellite observations and drifting buoy records to track ice parcels over several years (Tschudi et al. 2010). Satellite data has revealed that the Arctic has lost approximately one-third of its winter ice volume over the past 18 years, primarily as a thinning of multiyear ice (Kacimi and Kwok 2022). Although sea ice thickness is variable, perennial ice cover has been observed to be in rapid decline for decades (Comiso et al. 2008). When satellite data from 2002-2018 was compared with new dynamic snow coverage data, ice thickness of the Chukchi Sea in April was found to be declining at a rate 210 percent faster than with conventional climatology (Mallett et al. 2021). With the additional snow data, Chukchi Sea ice thickness was found to be in faster decline than previously modelled for the months of November, December, January, and April.

Sea ice extent fluctuates annually and is influenced by natural variations in atmospheric pressure and wind patterns, but clear linkages have been made to decreased Arctic sea ice extent and rising greenhouse gas concentrations dating back to the early 1990s (Timmermans and Proshutinsky 2014). A general downward trend in Arctic sea ice has occurred during the last few decades (Serreze et al. 2003). The maximum ice extent in March 2021 was tied with 2007 for the seventh lowest in the 43-year satellite record (5.7 million mi² [14.77 million km²]). The lowest maximum ice extents in the record occurred in March 2014 and March 2016 (2.9 million mi² [7.6 million km²]). This lowest maximum extent is five percent below the 1981 through 2010 average, though fairly typical of measurements taken in the last decade (Perovich et al. 2013). The ice is declining faster than previous computer models had projected, and this downward trend is predicted to continue (National Snow and Ice Data Center 2022; Timmermans and Proshutinsky 2014). A recent study from 21 research institutes found that under most climate simulations, including scenarios with significant greenhouse gas reductions, the Arctic Ocean is predicted to be ice-free (sea ice area less than 1 million km²) in summers by 2050 (Notz and Community 2020). The decrease in sea ice extent can be seen in Figure 3-3 below, illustrating yearly sea ice extent over various years (National Snow and Ice Data Center 2022).




Figure 3-3. Average Arctic Sea Ice Extent for March (1979-2022)

3.2 Biological Resources

Biological resources include living, native, or naturalized plant and animal species and the habitats within which they occur. Plant associations are referred to generally as vegetation, and animal species are referred to generally as wildlife. Habitat can be defined as the resources and conditions present in an area that support a plant or animal.

Within this OEA, biological resources are divided into five major categories: (1) invertebrates, (2) marine birds, (3) fish, (4) EFH, and (5) marine mammals. Threatened, endangered, and other special status species are discussed in their respective categories.

3.2.1 Regulatory Setting

For the purposes of this OEA, special status species are those species listed as threatened or endangered under the Endangered Species Act (ESA) and species afforded federal protection under the Marine Mammal Protection Act (MMPA) or the Migratory Bird Treaty Act (MBTA). Habitat may be protected as critical habitat for threatened or endangered species under the ESA or as EFH under the Magnuson-Stevens Fishery Conservation and Management Act (MSA).

3.2.1.1 Endangered Species Act

The purpose of the ESA (16 U.S.C. §§ 1531-1544) is to conserve the ecosystems upon which threatened and endangered species depend and to conserve and recover listed species. Section 7 of the ESA requires Federal agencies to consult with the responsible wildlife agency (i.e., U.S. Fish and Wildlife Service [USFWS] and/or National Marine Fisheries Service [NMFS]) to ensure that their actions are not likely to jeopardize the continued existence of federally listed threatened and endangered species, or result in the destruction or adverse modification of designated critical habitat (16 U.S.C. § 1536 (a)(2)). Regulations implementing the ESA include a requirement for consultation on those actions that “may affect” a listed species or adversely modify critical habitat.

If an agency’s proposed action would “take” a listed species, then the agency must obtain an incidental take authorization from the responsible wildlife agency. The ESA defines the term “take” to mean “harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or attempt any such conduct” (16 U.S.C. §1532(19)). The regulatory definitions of “harm” and “harass” are relevant to the Navy’s determination as to whether the Proposed Action would result in adverse effects on listed species:

Harm is defined by regulation as “an act which actually kills or injures” fish or wildlife (50 CFR § 222.102, 50 CFR § 17.3; 64 FR 60727, Nov 8 1999).

Harass is defined by USFWS regulation to mean an “intentional or negligent act or omission which creates the likelihood of injury to wildlife by annoying it to such an extent as to significantly disrupt normal behavioral patterns which include, but are not limited to, breeding, feeding, or sheltering” (50 CFR § 17.3). NMFS has not defined the term in its regulations.

Designation of critical habitat for listed species also falls under the ESA. NMFS regulations (50 CFR § 424.12(b)) state that, in determining what areas qualify as critical habitat, the agencies “shall consider those physical and biological features that are essential to the conservation of a given species and that may require special management considerations or protection.” These principal biological or physical constituent elements are referred to as “essential features” and “may include, but are not limited to, the following: spawning sites, feeding sites, seasonal wetland or dryland, water quality or quantity, geological formation, vegetation type, tide, and specific soil types.”

3.2.1.2 Marine Mammal Protection Act

All marine mammals are protected under the provisions of the MMPA (16 U.S.C. §§ 1361-1407). The MMPA prohibits any person or vessel from “taking” marine mammals in the United States or the high seas without authorization. The act further regulates “takes” of marine mammals in U.S. waters and by U.S. citizens on the high seas. The term “take,” as defined in Section 3 (16 U.S.C. § 1362) of the MMPA, means “to harass, hunt, capture, or kill, or attempt to harass, hunt, capture, or kill any marine mammal.”

The Proposed Action constitutes a military readiness activity as defined in Public Law 107-314 (MBTA (as amended) at 16 U.S.C. § 703 note) because these proposed scientific research activities directly support the “adequate and realistic testing of military equipment, vehicles, weapons, and sensors for proper operation and suitability for combat use” by providing critical data on the changing natural and physical environment in which such materiel will be assessed and deployed. The proposed scientific research enables navigation of vehicles that can be used to collect year-round information on the potential arctic battlespace. For military readiness activities, such as the Proposed Action, the relevant definition of harassment is any act that:

- Injures or has the significant potential to injure a marine mammal or marine mammal stock in the wild (“Level A harassment”); or
- Disturbs or is likely to disturb a marine mammal or marine mammal stock in the wild by causing disruption of natural behavioral patterns including, but not limited to, migration, surfacing, nursing, breeding, feeding, or sheltering to a point where such behavioral patterns are abandoned or significantly altered (“Level B harassment”) [16 U.S.C. § 1362(18)(B)(i) and (ii)].

3.2.1.3 Migratory Bird Treaty Act

The MBTA (16 U.S.C. §§ 703-712) and the Migratory Bird Conservation Act (16 U.S.C. §§ 715-715d, 715e, 715f-715r) of February 18, 1929, are the primary laws in the U.S. established to conserve migratory birds. The MBTA prohibits the taking, killing, or possessing of any migratory bird or their parts, nests, or eggs of such birds, unless permitted by regulation.

On February 28, 2007, the USFWS issued a final military readiness rule authorizing incidental takes of migratory birds resulting from military readiness activities. The definition of military readiness activities applies to the MBTA in the same way that it applies to the MMPA, and the Proposed Action is considered a military readiness activity for the purposes of this act. Under this regulation, the Navy must consider the potential environmental effects of its actions and assess the adverse effects of military readiness activities on migratory birds. If a Proposed Action may result in a significant adverse effect on a population of migratory bird species, the Navy shall consult with the USFWS to develop and implement appropriate conservation measures to minimize or mitigate these effects. A significant adverse effect on a population is defined as an effect that could, within a reasonable period of time, diminish the capacity of a population of a migratory bird species to sustain itself at a biologically viable level (50 CFR § 21.3). Conservation measures, as defined in 50 CFR § 21.3, include project designs or mitigation activities that are reasonable from a scientific, technological, and economic standpoint and are necessary to avoid, minimize, or mitigate the take of migratory birds or other potentially adverse impacts.

3.2.1.4 Magnuson-Stevens Fishery Conservation and Management Act

The MSA (16 U.S.C. §§ 1801-1822), enacted to conserve and restore the nation’s fisheries, includes a requirement for NMFS and regional fishery management councils to describe and identify EFH for all species that are federally managed. EFH is defined as those waters and substrate necessary to fish for spawning, breeding, feeding, or growth to maturity. Under the MSA, federal agencies must consult with the Secretary of Commerce regarding any activity or proposed activity that is authorized, funded, or undertaken by the agency that may adversely affect EFH. An adverse effect is any effect that may reduce the quantity or quality of EFH. Adverse effects may include direct or indirect physical, chemical, or biological alterations of the waters or substrate and loss of, or injury to, benthic organisms, prey species and their habitat, and other ecosystem components, if such modifications reduce the quality and/or quantity of EFH.

3.2.2 Affected Environment

The following discussions provide a description of the existing conditions for each of the categories of biological resources in the Study Area.

3.2.2.1 Invertebrates

Marine invertebrates are a large, diverse group containing tens of thousands of species distributed from warm shallow waters to cold deep waters throughout the global marine environment (Kohlbach et al. 2016). Invertebrates are the dominant animals in all habitats of the Study Area. Excluding microbes, approximately 5,000 known marine invertebrates have been documented in the Arctic; the number of species is likely higher, though, since this area is not well studied (Josefson et al. 2013). Although most species are found within the benthic (bottom) zone, marine invertebrates can be found in all zones (sympagic [within the sea ice], pelagic [water column], or benthic) of the Beaufort Sea (Josefson et al. 2013). Marine invertebrate distribution in the Beaufort Sea is influenced by habitat and oceanographic conditions (e.g., depth, temperature, salinity, nutrient concentrations, and ocean currents) (Levinton 2009). The cold water of the Arctic generally results in slow growth and high longevity among invertebrates and food sources, which are only seasonally abundant.

Major taxonomic groups found within the Beaufort Sea are listed and described in Table 3-1. No endangered, threatened, candidate, species of concern, or proposed species for listing under the ESA exist within the Study Area.

3.2.2.1.1 Canada Basin and Chukchi Sea Species

MacDonald et al. (2010) conducted an invertebrate sampling cruise in the summer of 2005 within the Canada Basin and Chukchi Borderland areas. MacDonald et al. (2010), as well as Ravelo et al. (2020), observed that abundance and biomass of invertebrates decreased with increasing depth and were lowest in the Canada Basin compared to the Chukchi Sea. However, biodiversity increased with increasing depth. Taxon inhabiting the Canada Basin ranged from 8 to 10 for macrofauna assemblages and 11 to 15 for megafauna assemblages, depending on where the sample was collected (MacDonald et al. 2010). Macrofauna assemblages were mainly composed of polychaetes, crustaceans (copepods, tanaids, isopods, cumaceans, amphipods, and ostracods), and mollusks, but also included nematodes, sipunculids, nemertean, pogonophorans, turbellarians, sponges, bryozoans, cnidarians, ascideans, holothurians, and ophiuroids. Megafauna assemblages within the Canada Basin were dominated by lebensspuren, Actiniaria, and holothuroid (MacDonald et al. 2010).

Table 3-1. Taxonomic Groups of Marine Invertebrates in the Study Area

<i>Invertebrate Group</i>		<i>Presence in Study Area</i>		
<i>Common Name (Taxonomic Group)</i>	<i>Description</i>	<i>Sympagic</i>	<i>Pelagic</i>	<i>Benthic</i>
Flatworms (Phylum Platyhelminthes) ¹	Simplest form of marine worm with a flattened body.	✓		✓
Ribbon worms (Phylum Nemertea) ¹	Worms with a long extension from the mouth (proboscis) that helps capture food.		✓	✓
Roundworms (Phylum Nematoda) ¹	Small worms; many live in close association with other animals (typically as parasites).	✓	✓	✓
Arrow worms (Phylum Chaetognatha) ¹	Predatory worms with bristle-like jaws.			✓
Sponges (Phylum Porifera) ²	Large species have calcium carbonate or silica structures embedded in cells to provide structural support.			✓
Segmented worms (Phylum Annelida) ²	Highly mobile marine worms; many tube-dwelling species.	✓	✓	✓
Bryozoans (Phylum Bryozoa) ³	Lace-like animals that exist as filter feeding colonies. Form either encrusting or bushy-tuftlike lacy colonies.			✓
Hydroids and jellyfish (Phylum Cnidaria) ²	Animals with stinging cells.	✓	✓	✓
Comb jellies (Phylum Ctenophora) ²	Jelly-like animals that swim with the use of cilia. They lack stinging cells.		✓	✓
Cephalopods, bivalves, sea snails, chitons (Phylum Mollusca) ²	Mollusks are a diverse group of soft-bodied invertebrates with a specialized layer of tissue called a mantle. Mollusks such as squid are active swimmers and predators, while others such as sea snails are predators or grazers and clams are filter feeders.		✓	✓
Shrimp, crab, barnacles, copepods, amphipods, ostracods (Phylum Arthropoda – Crustacea) ²	Diverse group of animals, some of which are immobile. Most have an external skeleton. All feeding modes from predator to filter feeder.	✓	✓	✓
Sea stars, sea urchins, sea cucumbers (Phylum Echinodermata) ²	Predators and filter feeders with tube feet.			✓
Tunicates, ascidians, larvaceans, and sea squirts (Phylum Chordata; Subphylum Tunicata) ²	Filter feeders that siphon water for feeding and respiration. Adults may be sessile or free-swimming. Some species are solitary; others live in clonal colonies.		✓	✓

¹Based on Arctic Ocean biodiversity (Bluhm 2008), and confirmed within the Study Area by (Kosobokova et al. 2011) and (Grebmeier et al. 2015).

²Invertebrate phyla are based on the World Register of Marine Species (Appeltans et al. 2010), Catalogue of Life (Bisby et al. 2014), and Ecological Atlas of the Bering, Beaufort, and Chukchi Seas (Smith et al. 2017).

³Phyla not extracted when searched the distribution of the Beaufort Sea on the World Register of Marine Species. Individual species found on Arctic Ocean biodiversity, and verified via the distribution maps on the World Register of Marine Species (Appeltans et al. 2010).

Another survey of zooplankton by Kosobokova and Hopcroft (2010) was conducted in the summer of 2005 in the Canada Basin. This study identified 14 taxonomic groups of zooplankton including 111 species in the area. These taxonomic groups included Copepoda, Amphipoda, Euphausiacea, Decapoda, ostracoda, Cnidaria, Ctenophora, Polychaeta, Pteropoda, Chaetognatha, Larvacea, Forminifera, Radiolaria, and Tintinnina. Of the 111 species identified, 74 were crustaceans (copepods, euphausiids, amphipods, decapods, and ostracods), 17 were cnidarians (hydromedusae, scyphomedusae, siphonophora), 1 was foraminifera, 4 were ctenophores, 2 were pteropods, 4 were larvaceans, 4 were chaetognaths, and 5 were polychaetes. Copepods were the most dominant invertebrate species in the area, making up approximately 91 percent of the species' abundance. Similar to MacDonald et al. (2010), Kosobokova and Hopcroft (2010) observed abundance and biomass of invertebrates decreasing with an increased depth whereas biodiversity increased with an increase in depth. Specifically, they noted a progressive decrease in zooplankton abundance and biomass below 164 ft (50 m), followed by a slight increase in the 656 ft to 984 ft (200 to 300 m) layer, and a slow decrease below 984 ft (300 m). The increase at 656 ft (200 m) is thought to be attributed to the transition between the Pacific halocline and Atlantic waters (Kosobokova and Hopcroft 2010). It is important to note that both of these studies only surveyed species in the water column, at a limited number of locations, and during the summer months (Kosobokova and Hopcroft 2010; MacDonald et al. 2010). Therefore, not all species (i.e., benthic invertebrates) are represented in these surveys.

A large data set of benthic invertebrate species abundance and biomass in the Chukchi Sea was developed by Grebmeier et al. (Grebmeier et al. 2015) using time-series data taken from sampling stations during long-term census cruises that are part of the Distributed Biological Observatory Network (DBO). Macroinfauna were sampled at 114 stations during Russian-American Long-term Census of the Arctic (RUSALCA) cruises in 2004, 2009, and 2012. Epibenthic data was also collected at most of the stations. The dominant taxa of macrofauna identified in Alaskan coastal waters and the Bering Sea were Ascidiacea (tunicates and sea squirts), Polychaeta, Bivalvia, and Sipuncula (annelid worms). The dominant epifauna were Echinodermata, Gastropoda, Crustacea, and Bivalvia (Grebmeier et al. 2015). Community compositions at time-series sites remained relatively constant over the study period, although estimates of biomass were variable and indicated a declining trend since 2009 (Grebmeier et al. 2015).

Cooper et al. (2019) conducted a two-phase video survey (2008 and 2016-2017) of the seafloor in the northern Bering and Chukchi seas as part of the DBO. While sampling was not extensive enough to directly compare epifaunal density with earlier trawling surveys, observed community assemblages were similar. The presence of specific organismal community assemblages were associated with environmental data available from other DBO projects, such as sediment grain size and water mass identity (Cooper et al. 2019). In muddier sediments, like those found within the Study Area, deposit feeders (e.g., brittle stars) dominated.

Based on previous studies (Harding 1966; Virketis 1957), the overall species assemblages in the Canada Basin had not changed significantly in the 60 years prior to the research of Kosobokova and Hopcroft (2010). However, given major changes to the Arctic environment in the last decade, it is predicted that subarctic taxon will expand their ranges northward from both the northeast Atlantic and northwest Pacific into the Arctic region (Chan et al. 2019; Renaud et al. 2015; Waga et al. 2020). Recent research has provided initial evidence of this trend toward species range shifts in the Arctic. Grebmeier et al. (2018) identified decline in seasonal ice cover and changes to the timing of spring ice melt as an influence to nutrient distribution in seafloor communities. In southern Arctic waters, key saltwater clams were observed to be declining in biomass; however, bivalve biomass expanded northward (Grebmeier et al. 2018; Lovvorn et al. 2016).

Environmental changes also are expected to increase the potential for non-indigenous species to be introduced to the Arctic. Between the years 1960 and 2015, Chan et al. (2019) recorded 54 instances where 34 unique non-indigenous species were identified in the Arctic. The majority of species with known origins were from the northeast Atlantic and the northwest Pacific. Routes of transfer included vessels and aquaculture activities, although this research also included species that arrived via natural spread (Chan et al. 2019). The phyla Arthropoda and Ochrophyta were most frequently introduced.

Because of the large number of species, a general discussion of each ecologic zone (sympagic, pelagic, and benthic) is provided below.

3.2.2.1.2 Sympagic Zone

Sea ice provides a habitat for algae and a nursery ground for invertebrates during times when the water column does not support phytoplankton growth (Michel et al. 2002; Winfree 2005). Sympagic zone invertebrates live within the pores and brine channels of the ice (small spaces within the sea ice which are filled with a salty solution called brine) or at the ice-water interface. Biodiversity of species is low within the sympagic zone due to the extreme conditions of the sea ice (Nuttall 2005). Species abundance within the ice has been found to be highly variable with most species occurring within the bottom 4 inches (in; 10 centimeters [cm]) of ice core samples. Species are also found in greater densities in coastal fast ice compared to offshore pack ice. Additionally, abundance is 1 to 4 orders of magnitude greater in spring and early summer (compared to winter) in coastal fast ice (Bluhm et al. 2010). Within the Study Area, many sympagic species also exist in and along the edges of ice coverage, feeding on blooms of phytoplankton and other algae which grow in, on, or adjacent to the ice (Wyllie-Echeverria and Ackerman 2003).

The most dominant sympagic species are nematodes, harpacticoid copepods, and rotifers (Josefson et al. 2013). At the ice-water interface, *Apherusa glacialis*, *Onisimus glacialis*, *O. nanseni*, and *Gammarus wilkitzkii* are common amphipods (Gradinger et al. 2010). Although the sympagic environment is spatially limited, recent research indicates that large pelagic copepod species such as *Calanus glacialis* and *C. hyperboreus*, which are a primary food source for higher trophic levels, are substantially dependent on sea ice synthesized carbon, illustrating the importance of this unique environment to the broader Arctic food web (Sheffield-Guy et al. 2014).

3.2.2.1.3 Pelagic Zone

Pelagic habitats include downwelling and upwelling areas and frontal zones. Dominant species groups within the pelagic zone are highly stratified by depth. In a zooplankton survey from the Arctic Canadian Basin (east of the Study Area) within the pelagic zone, 50 percent of the biomass was concentrated in the upper layer from 0 to 328 ft (0 to 100 m) in depth (Hopcroft et al. 2005). The pelagic zone invertebrate fauna is dominated by large copepods, such as *Calanus glacialis* and *C. hyperboreus*. Copepods in the pelagic zone of the Beaufort Sea have longer life cycles (2–4 years) and are larger than copepod species living in warmer water (Hopcroft et al. 2008). Sirenko (2001) and Sirenko et al (2010) found that cnidarians are second to copepods in diversity and numbers. Jellyfish are likely important invertebrate predators within this zone (Josefson et al. 2013). Due to warming in the regional water, the distribution ranges of pelagic zooplankton species in the Chukchi Sea have been shifting in recent years, especially with copepods (Ershova et al. 2015).

3.2.2.1.4 Benthic Zone

The benthic zone is the most diverse and species-rich habitat, where the majority of the species within the Study Area can be found. In a compilation of 35 datasets, the Beaufort Sea was found to be a hotspot of benthic diversity in Canada's oceans (Wei et al. 2020). Generally, benthic marine

invertebrates play an important role in the food web as scavengers, recyclers of nutrients, habitat-forming organisms, and prey to predators (e.g., fish and whales). The highest epibenthic (i.e., living on the surface of the seafloor) biomass is found in the Polar Mixed Layer and the Arctic Halocline (outer shelf and upper slope) and the least biomass is found in deep waters of the Canada Basin (Ravelo et al. 2020).

Within the Arctic region, major species groups within the benthic zone that have the highest diversity and abundance are Arthropoda (e.g., crabs and barnacles), Bryozoa (moss animals), Mollusca (e.g., snails and clams), and Nematoda (Josefson et al. 2013). In a 2010 Beaufort Sea trawl, the invertebrates with the highest densities in descending order of abundance were the notched brittle star (*Ophiura sarsi*), snow crab (*Chionoecetes opilio*), mussels (*Musculus* spp.), and the mud star (*Ctenodiscus crispatus*) (Rand and Logerwell 2010). Within the sediment, roundworms are one of the most widespread marine invertebrates with population densities of one million organisms per 10.8 square feet (1 square meter) of mud (Levinton 2009). The principal habitat-forming invertebrates of the benthos are Porifera (e.g., sponges), Annelida (e.g., tube worms), and Mollusca (e.g., spiral margarite).

3.2.2.1.5 Invertebrate Hearing

Limited data is available on the hearing capabilities of marine invertebrates (Lovell et al. 2005; Popper and Schilt 2008). While data are limited, research suggests that some of the major cephalopods and decapods have limited hearing capabilities (Hanlon 1987; Offutt 1970). They may hear only low-frequency (less than 1 kHz) sources, with best sensitivities at lower frequencies (Budelmann 2010; Mooney et al. 2010; Offutt 1970; Packard et al. 1990), which is most likely within the frequency band of biological signals (Hill 2009). A few cephalopods may sense higher frequencies up to 1,500 Hz (Hu et al. 2009). Both behavioral and auditory brainstem response studies suggest that crustaceans may sense frequencies up to 3 kHz, but best sensitivity is likely below 200 Hz (Goodall et al. 1990; Lovell et al. 2005; Lovell et al. 2006). Based on a review of crustacean sensitivity of high amplitude underwater noise by Edmonds et al. (2016), crustaceans may be able to hear the frequencies at which they produce sound, but it is unclear which noises are incidentally produced and if there are any negative effects from masking those frequencies. Acoustic signals produced by crustaceans range from low frequency rumbles (20-60 Hz) to high frequency signals (20-55 kHz) (Celi et al. 2014; Henninger and Watson 2005; Patek and Caldwell 2006; Staaterman et al. 2011). Another review by Tidau and Briffa (2016) recognized that noise from low-frequency pile driving, airguns, seismic surveys, vessel noise, and ambient white noise had a variety of impacts on crustaceans, including increased locomotion and stress, reduced antipredator behavior, changes in foraging, and reduced actions with ecological functions (e.g., bioirrigation).

Aquatic invertebrates that can sense local water movements with ciliated cells include cnidarians, flatworms, segmented worms, urochordates (tunicates), mollusks, and arthropods (Budelmann 1992a, 1992b; Popper et al. 2001). Some aquatic invertebrates have specialized organs called statocysts for determination of equilibrium and, in some cases, linear or angular acceleration. Statocysts allow an animal to sense movement and may enable some species, such as cephalopods and crustaceans, to be sensitive to water particle movements associated with sound (Hu et al. 2009; Kaifu et al. 2008; Montgomery et al. 2006; Popper et al. 2001). Because any acoustic sensory capabilities, if present at all, are limited to detecting water motion, and water particle motion near a sound source falls off rapidly with distance, aquatic invertebrates are probably limited to detecting nearby sound sources rather than sound caused by pressure waves from distant sources. However, sound propagation models used to assess the distance over which impacts might occur have rarely been validated by actual measurements and are ineffective at modelling transmission under shallow water conditions, close to or within the seabed, or at the surface (Hawkins and Popper 2017; Popper and Hawkins 2018).

Outside of studies conducted to test the sensitivity of invertebrates to vibrations, very little is known on the effects of anthropogenic underwater noise on invertebrates (Edmonds et al. 2016). Many studies that have assessed the impacts of noise on invertebrates have overlooked the sensitivity of these species to particle motion rather than sound pressure (Hawkins and Popper 2017). However, a growing research body offers insights on the effects of anthropogenic noise on marine invertebrate behavior and physiology. A meta-analysis by Murchy et al. (2019) of 11 studies found that shipping noise had negative behavioral and physiological consequences for bivalves, cephalopods, and gastropods. Shipping noise altered the foraging and predator avoidance behaviors across taxa in various ecosystems (Murchy et al. 2019). Additional studies also have revealed negative consequences of sound. Blue mussels exhibited DNA damage, oxidative stress, and reduced filter feeding and oxygen consumption when exposed to ship noise (approximately 656 to 984 ft [200 to 300 m] from the source) (Wale et al. 2019). Cuttlefish experienced damage to their statocysts when exposed to low intensity, low frequency vessel sounds (139 to 142 dB at band centered at 315 Hz and 400 Hz) (Solé et al. 2017). Cnidarians were susceptible to statocyst damage from noise pollution at 50-400 Hz and 157-175 dB (Solé et al. 2016). When eggs of sea hares were exposed to additional boat noise during embryonic development, a link to developmental failure at the embryonic stage and increased mortality of larvae was noted (Nedelec et al. 2014b).

Marine invertebrates sometimes respond to noise in a similar way to their response to predators, which costs energy and can have a negative effect on their long-term survivability. In response to sounds from pile driving, squid exhibited antipredator defense behaviors, altered feeding behaviors, and suggested a disruption to essential communication behaviors (Jones et al. 2021; Jones et al. 2020). Vibrations directly connected to the substrate also reduced a chemically guided shell-searching behavior in hermit crabs (Roberts and Laidre 2019). In contrast, the meta-analysis by Murchy et al. (2019) found a net positive effect on the overall health of invertebrates sampled when exposed to vibrations from low-frequency (10 – 300 Hz) impulsive sounds associated with seismic surveys (Carroll et al. 2017), although additional data is still needed to accurately assess the effects of seismic surveys on marine invertebrates.

3.2.2.2 Marine Birds

For the purpose of this document, “marine birds” refers to shoreline, coastal, and pelagic bird species. A description is provided below for species of marine birds that would likely occur in the Study Area and includes species protected under the MBTA. Although ESA-listed Steller’s eider (*Polysticta stelleri*) and spectacled eider (*Somateria fischeri*) may occur within the Study Area, it is unlikely that they would overlap with any of the elements of the Proposed Action; therefore, they are not considered further herein.

A combination of short-distance migrants, long-distance migrants, and year-round resident marine bird species occur within the Study Area. Typical behaviors that would be encountered within the Study Area predominantly include foraging and migrating, among others. Black-legged kittiwakes (*Rissa tridactyla*) and ivory gulls (*Pagophila aburnea*) are associated with sea ice and inhabit waters along the continental shelf about 90 nm (167 km) from the shore. Other species found near or over the Canada Basin include glaucous gull (*Larus hyperboreus*), herring gull (*Larus argentatus*), Ross’s gull (*Rhodostethia rosea*), northern fulmar (*Fulmarus glacialis*), and thick-billed murre (*Uria lomvia*) (Harwood et al. 2005). Of all the marine birds that occur in the vicinity of the Study Area, only the thick-billed murre exhibits foraging diving behaviors at distances greater than 90 nm (167 km) from the shoreline during the timeframe of the Proposed Action. However, the thick billed murre is not expected to forage in the deep waters of the Study Area. Therefore, no birds are expected to be foraging within the Study Area.

Some bird species may be migrating through the Study Area. These species include black guillemot (*Cepphus grylle*), ivory gull, Ross's gull, short-tailed shearwater (*Puffinus tenuirostris*), king eider (*Somateria spectabilis*), and long-tailed duck (*Clangula hyemalis*) (National Audubon Society 2015). None of these species are listed under the ESA, but all are protected under the MBTA. Species in the orders Charadriiformes (i.e., ivory gull, Ross's gull, thick-billed murre, black guillemot) and Procellariiformes (i.e., northern fulmar, short-tailed shearwater) are expected.

Within the Study Area, two species from the family Laridae (ivory gull [*Pagophila eburnea*] and Ross's gull [*Rhodostethia rosea*]) may be present during the timeframe of the Proposed Action. These species winter in the Arctic Ocean, and will spend time at edges of pack ice. Outside of the breeding season, ivory gulls occur singly or in flocks of up to 20 individuals (BirdLife International 2016; International Union for the Conservation of Nature 2016). These species consume fish, surface-dwelling marine invertebrates, and algae, though ivory gulls also will scavenge on marine mammal remains on the sea ice (International Union for the Conservation of Nature 2016; Kaufman 1996). Ross's gull will forage solitarily or in small, loose flocks.

Thick-billed murre may be encountered in the shallower water near the Study Area year-round, but more commonly in the summer. They have a circumpolar distribution in the arctic region (BirdLife International 2012). Known breeding sites occur in coastal areas and islands of the Beaufort Sea (Gaston and Hipfner 2000). The thick-billed murre is one of the most numerous marine birds in the Northern Hemisphere. In the summer months, it inhabits continental-shelf waters of the Arctic Ocean and adjacent seas, including the Beaufort Sea. Their range shifts a bit more to the south in the winter, occurring in greater number in the Bering Sea and coastal areas of southern Alaska (Gaston and Hipfner 2000). Thick-billed murre's diet consists of mid-water school fish (cod, smelt, sandlance), pelagic amphipods and euphausiids, benthic fish (sculpins, blennies, lumpsuckers), deep water fish (lanternfish), shrimp, squid, and annelids (Gaston and Hipfner 2000). Thick-billed murre may travel up to 92 nm (170 km) from their breed colonies to forage (Gaston and Hipfner 2000). They travel in V-formation flocks to foraging sites, but are mainly solitary feeders. However, they can aggregate in large groups where prey is concentrated (Cairns and Schneider 1990; Schneider et al. 1990). They capture prey underwater with maximum diving depths of 690 ft (210 m) and more typical depths around 33 ft (10 m) (Croll et al. 1992).

Procellariiformes is a large order of pelagic marine birds. Fulmars are medium to large birds, and are typically scavengers. Shearwaters obtain their food at or close to the water's surface (Brooke 2001). Typically only non-breeding short-tailed shearwaters will stay within the Study Area during the winter, though most of this species migrates south and will return to the Arctic in May (U.S. Fish and Wildlife Service 2006). This order includes species that are generally long lived, breed once per year, and lay only one egg; thus, they have a low reproductive output.

3.2.2.1 Marine Bird Hearing

Although hearing range and sensitivity has been measured for many terrestrial birds, little research has been conducted on the hearing capabilities of marine birds. The majority of published literature on bird hearing focuses on terrestrial birds, particularly songbirds, and their ability to hear in the air. A review of 32 terrestrial and marine species reveals that birds generally have greatest hearing sensitivity between 1 and 4 kHz (Beason 2004; Dooling 2002). Research shows that very few birds can hear below 20 Hz, most have an upper frequency hearing limit of 10 kHz, and none exhibit the ability to hear frequencies higher than 15 kHz (Dooling 2002; Dooling et al. 2000). Hearing capabilities have been studied for only a few marine birds (Beason 2004; Beuter et al. 1986; Thiessen 1958; Wever et al. 1969); these studies show that marine birds have hearing ranges and sensitivities that are consistent with what is currently known about general bird hearing capabilities.

Auditory abilities have been measured in ten diving bird species in-air using electrophysiological techniques (Crowell et al. 2015). All species tested had the best hearing sensitivity from 1 to 3 kHz. The red-throated loon (*Gavia stellata*) and northern gannet (*Morus bassanus*) (both non-duck species) had the highest thresholds of the diving species while the lesser scaup (*Aythya affinis*) and ruddy duck (*Oxyura jamaicensis*) (both duck species) had the lowest thresholds (Crowell et al. 2015). Auditory sensitivity varied amongst the species tested, spanning over 30 decibels (dB) in the frequency range of best hearing. While electrophysiological techniques provide insight into hearing abilities, auditory sensitivity is more accurately obtained using behavioral techniques. Crowell (2016) used behavioral methods to obtain an in-air audiogram of the lesser scaup. Best hearing frequency range in-air was similar to other birds, with best sensitivity of 14 dB referenced to 20 micropascals (re 20 μ Pa) at 2.68 kHz. Another study used physiological auditory evoked potential (AEP) methods to obtain and compare auditory curves of the Atlantic puffin and the common murre (Mooney et al. 2019). Hearing data for the puffin was comparable to other birds, while the responses of the murre were elevated and the frequency range was narrower (1-4 kHz with no response at 3 kHz). Crowell et al. (2015) also compared the vocalizations of the same ten diving bird species to the region of highest sensitivity of in-air hearing. Of the birds studied, vocalizations of only eight species were obtained due to the relatively silent nature of two of the species. The peak frequency of vocalizations of seven of the eight species fell within the range of highest sensitivity of in-air hearing. Crowell et al. (2015) suggested that the colonial nesters tested had relatively reduced hearing sensitivity because they relied on individually distinctive vocalizations over short ranges. Additionally, Crowell et al. (2015) observed that the species with more sensitive hearing were those associated with freshwater habitats, which are relatively quieter compared to marine habitats with wind and wave noise.

Although important to seabirds in air, it is unknown if seabirds use hearing or vocalizations underwater for foraging, communication, predator avoidance, or navigation (Crowell 2016; Dooling and Therrien 2012). Some scientists suggest that birds must rely on vision rather than hearing while underwater (Hetherington 2008), while others suggest birds must rely on an alternative sense in order to coordinate cooperative foraging and foraging in low light conditions (e.g., night, at depth) (Dooling and Therrien 2012).

There is little known about the hearing ability of birds underwater (Dooling and Therrien 2012). In air, the size of the bird is usually correlated with the sensitivity to sound (Johansen et al. 2016); for example, songbirds tend to be more sensitive to higher frequencies and larger non-songbirds tend to be more sensitive to lower frequencies (Dooling et al. 2000). Two studies have tested the ability of a single diving bird, a great cormorant (*Phalacrocorax carbo sinensis*), to respond to underwater sounds (Hansen et al. 2017; Johansen et al. 2016); cormorants would not occur within the Study Area. Until Larsen et al. (2020), all studies utilized one or two trained birds rather than a wider sample. These studies suggest that the cormorant's hearing in air is less sensitive than birds of similar size; however, the hearing capabilities in water are better than what would be expected for a purely in-air adapted ear (Hansen et al. 2017; Johansen et al. 2016). The frequency range of best hearing underwater was observed to be narrower than the frequency range of best hearing in air, with greatest sensitivity underwater observed around 2 kHz (about 71 dB referenced to 1 micropascal [re 1 μ Pa] based on behavioral responses). Another study found that the auditory brainstem response (ABR) waveshape and latency, as well as the ABR threshold in units of sound pressure, of cormorants were similar in air and water (Larsen et al. 2020). The best average sound pressure sensitivity was 1 kHz, both in air (53 decibels referenced to 1 micropascal [dB re 1 μ Pa]) and underwater (58 dB re 1 μ Pa) (Larsen et al. 2020). However, when adjusted for sound intensity, sensitivity was higher underwater than in air, suggesting that their in-water hearing is equal to if not better than their in-air hearing (Larsen et al. 2020).

Diving birds have adaptations to protect their ears from pressure changes, which may limit their ability to hear underwater (Dooling and Therrien 2012). Because reproduction and communication with conspecifics occurs in air, adaptations for diving may have evolved to protect in-air hearing ability and may contribute to reduced sensitivity underwater (Hetherington 2008). There are many anatomical adaptations in diving birds that may reduce sensitivity both in air and underwater. Common murre (*Uria aalge*) were deterred from gillnets by acoustic transmitters emitting 1.5 kHz pings at 120 dB re 1 μ Pa; however, no significant reaction was observed in rhinoceros auklet (*Cerorhinca monocerata*) bycatch in the same nets (Melvin et al. 1999). In a quiet, controlled environment, common murre showed consistent reactions to sounds ranging from 110 to 137 dB re 1 μ Pa (Hansen et al. 2020).

Gentoo penguins (another species that would not occur within the Study Area but has been the subject of underwater hearing studies) showed a gradient of reactions to noise burst played in a large aquarium tank, from no reactions at 100 dB to strong reactions at 120 dB more than half of the time (Sørensen et al. 2020). Another study found that King, Gentoo, and Macaroni penguins make short (less than 1 second) bursts of sound during feeding dives (at 998 Hz, 1,097 Hz, and 680 Hz, respectively) (Thiebault et al. 2019). While the majority of these vocalizations were directly associated with hunting behavior, acceleration, and/or an attempt to catch prey, the functions of these vocalizations is speculative.

3.2.2.3 Fish

The fish species located in the Study Area include those that are closely associated with the deep ocean habitat of the Beaufort and Chukchi Seas. Nearly 250 marine fish species have been identified in the Arctic, excluding the larger parts of the sub-Arctic Bering, Barents, and Norwegian Seas (Mecklenburg et al. 2011). However, only about 30 are known to occur in the Arctic waters of the Beaufort Sea (Christiansen and Reist 2013). Largely because of the difficulty of sampling in remote, ice-covered seas, many high-Arctic fish species are known only from rare or geographically patchy records (Mecklenburg et al. 2013). Aquatic systems of the Arctic undergo extended seasonal periods of ice cover and other harsh environmental conditions. Fish inhabiting such systems must be biologically and ecologically adapted to surviving such conditions. Important environmental factors that Arctic fish must contend with include reduced light, seasonal darkness, ice cover, low biodiversity, and low seasonal productivity. No ESA-listed fish species occur within the Study Area.

3.2.2.3.1 Major Fish Groups

Marine fish can be broadly categorized into horizontal and vertical distributions within the water column. The primary distributions of fish that occur in the marine environment of the Study Area are within the water column near the surface. While there are multiple major fish groups inhabiting the deep waters of the Beaufort and Chukchi Seas (Table 3-2), the only federally-managed species within the Study Area is the Arctic cod (*Boreogadus saida*).

Table 3-2. Major Groups of Marine Fish in the Study Area during the Proposed Action

Common Name	Scientific Name	Vertical Distribution Within the Study Area*
Cod	Order Gadiformes	Water column
Scorpionfish	Order Scorpaeniformes	Seafloor, water column
Eelpouts, Eelblennys, and Wolffishes	Order Perciformes	Seafloor

* All distribution information was obtained from Food and Agriculture Organization of the United Nations (Cohen et al. 1990), Kaschner et al. (2013), and Arctic Ocean Diversity (Mecklenburg and Mecklenburg 2009).

Cods (Order Gadiformes)

The two major species of cod within the Study Area are Arctic cod (*Boreogadus saida*) and polar cod (*Arctogadus glacialis*). Cod are an important component in the food web of the Beaufort Sea environment, preying on plankton, and being preyed upon by ringed seal (*Phoca hispida*), bearded seal (*Erignathus barbatus*), beluga whale (*Delphinapterus leucas*), and many marine birds (including gulls and guillemots) (Bluhm and Gradinger 2008; Cohen et al. 1990; Welch et al. 1993). Fish inhabiting the water column of oceanic waters seaward of the 200-m isobath comprise this assemblage; most species exhibit some preference of bathymetric stratification.

Arctic cod are the only federally managed species within the Study Area (for more information, see Section 3.2.2.4). Arctic cod is the northernmost-occurring fish species and is widespread throughout Arctic seas (Mecklenburg et al. 2013). Arctic cod are both cryopelagic (live in cold, deep water) and epontic (live on the underside of ice). They use sea ice for shelter, to capture prey, and to avoid predators. Arctic cod often occur in ice holes, cracks, hollows, and cavities in the lower surface of the ice and are most common near the ice edge or among broken ice. As the ice thaws at these margins, plankton grow and provide a food source. In the northeastern Chukchi Sea and western Beaufort Sea, they are found most abundantly in deep, cold, and highly saline water (Logerwell et al. 2018). They occur in the open-ocean waters of the Study Area from the surface to depths of 1,312 ft (400 m). Onshore-offshore movements are associated with spawning and movements of ice. Cod are generally found near the bottom in the continental shelf areas, feeding on benthic organisms (Paxton and Eshmeyer 1998). The primary offshore food source of Arctic cod is plankton (Mecklenburg et al. 2011). Specifically, they feed predominantly on epibenthic mysids, amphipods, copepods, and fish (Cohen et al. 1990). It is possible that Arctic cod also feed on the amphipod-diatom ice community inhabiting the lower ice layer. This species moves and feeds in different groupings, dispersed in small and very large schools throughout the water column (Welch et al. 1993).

Scorpionfish (Order Scorpaeniformes)

Scorpionfish, of the order Scorpaeniformes, are distinguishable by the well-developed spines on their cheeks, and the distinct ridges or spines on top of the head. Adults of most Arctic species live on the seafloor, but some are both benthic and pelagic. These fish typically consume small crustaceans, worms, clams, and fish eggs. One example of a scorpionfish that inhabits the Study Area is the gelatinous seasnail (*Liparis fabricii*), which is both benthic and pelagic, living at depths up to 8,202 ft (2,500 m) (Mecklenburg et al. 2011). Scorpionfish are prey species for other fishes and marine birds.

Eelpouts, Eelblennys, and Wolffishes (Order Perciformes)

Though most species of the order Perciformes are found in the benthic habitats of shallower shelf waters, some species are associated with deep-water marine environments. One such species is the glacial eelpout (*Lycodes frigidus*), which is endemic to the Arctic basins. This species is benthic in water depths up to 9,843 ft (3,000 m) (Mecklenburg et al. 2011). To feed themselves, these species move along the seafloor and use the cartilaginous keels on their lower jaws to stir up prey, such as crustaceans, worms, and fishes (Mecklenburg et al. 2011).

3.2.2.3.2 Hearing

All fish have two sensory systems to detect sound in the water: the inner ear, which functions very much like the inner ear in other vertebrates, and the lateral line, which consists of a series of receptors along the fish's body (Popper and Schilt 2008). The lateral line system is sensitive to external particle motion arising from sources within a few body lengths of the animal. The lateral line detects particle motion at low frequencies from 1 Hz up to at least 400 Hz (Coombs and Montgomery 1999; Hastings and Popper

2005; Higgs and Radford 2013; Webb et al. 2008). The inner ears of fish contain three dense otoliths (i.e., small calcareous bodies) that sit atop many delicate mechanoelectric hair cells, similar to the hair cells found in mammalian ears. Sound waves in water tend to pass through fish's bodies, which have a composition similar to water, and vibrate the otoliths. This causes a relative motion between the dense otoliths and the surrounding tissues causing a deflection of the hair cells, which are sensed by the nervous system.

Although a propagating sound wave contains pressure and particle motion components, particle motion is most significant at low frequencies (less than a few hundred Hz) and closer to the sound source (Popper and Fay 2010). The inner ears of fishes are directly sensitive to acoustic particle motion rather than acoustic pressure. Historically, studies that have investigated hearing in, and effects to, fishes have been carried out with sound pressure metrics. Although particle motion may be the more relevant exposure metric for many fish species, there is little data available that actually measures it due to a lack in standard measurement methodology and experience with particle motion detectors (Hawkins et al. 2015; Martin et al. 2016). In these instances, particle motion can be estimated from pressure measurements (Nedelec et al. 2016).

Some fishes possess additional morphological adaptations or specializations that can enhance their sensitivity to sound pressure, such as a gas-filled swim bladder (Astrup 1999; Popper and Hastings 2009). A fish's gas-filled swim bladder can enhance sound detection by converting acoustic pressure into localized particle motion, which may then be detected by the inner ear (Radford et al. 2012). Fish with swim bladders generally have better sensitivity and better high-frequency hearing than fish without swim bladders (Popper 2014; Popper and Hastings 2009). In addition, structures such as gas-filled bubbles near the ear or swim bladder, or even connections between the swim bladder and the inner ear, also increase sensitivity and allow for high-frequency hearing capabilities and better sound pressure detection.

Although many researchers have investigated hearing and vocalizations in fish species (Ladich and Fay 2013; Popper 2014), hearing capability data only exist for fewer than 100 of the over 35,000 fish species (Eschmeyer and Fong 2017). Data suggest that most species of marine fish either lack a swim bladder (e.g., sharks and flatfishes) or have a swim bladder not involved in hearing and can only detect sounds below 1 kHz. Some marine fishes (clupeiforms) with a swim bladder involved in hearing are able to detect sounds to about 4 kHz (Colley et al. 2016; Mann et al. 2001; Mann et al. 1997). One subfamily of clupeids (i.e., Alosinae) can detect high- and very high-frequency sounds (i.e., frequencies from 10 to 100 kHz, and frequencies above 100 kHz, respectively), although auditory thresholds at these higher frequencies are elevated and the range of best hearing is still in the low-frequency range (below 1 kHz) similar to other fishes. Mann et al. (Mann et al. 1998; Mann et al. 1997) theorize that this subfamily may have evolved the ability to hear relatively high sound levels at these higher frequencies in order to detect echolocations of nearby foraging dolphins. For fishes that have not had their hearing tested, such as deep sea fishes, the suspected hearing capabilities are based on the structure of the ear, the relationship between the ear and the swim bladder, and other potential adaptations such as the presence of highly developed areas of the brain related to inner ear and lateral line functions (Buran et al. 2005; Deng et al. 2011; Deng et al. 2013). It is believed that most fishes have their best hearing sensitivity from 100 to 400 Hz (Popper 2003).

Permanent hearing loss has not been documented in fish. A study by Halvorsen et al. (2012) found that for temporary hearing loss or similar negative impacts to occur, the noise needed to be within the fish's individual hearing frequency range; external factors, such as developmental history of the fish or environmental factors, may result in differing impacts to sound exposure in fish of the same species. The sensory hair cells of the inner ear in fish can regenerate after they are damaged, unlike in mammals

where sensory hair cells loss is permanent (Lombarte et al. 1993; Smith et al. 2006). As a consequence, any hearing loss in fish may be as temporary as the timeframe required to repair or replace the sensory cells that were damaged or destroyed (Smith et al. 2006), and no permanent loss of hearing in fish would result from exposure to sound.

Few auditory studies have been completed on Arctic cod. Using acoustic telemetry and modeled ship noise, one recent study found that the presence of vessels in the highly trafficked Resolute Bay in Nunavut, Canada resulted in home range displacement of Arctic cod near the vessel (Ivanova et al. 2020). Individuals altered their swimming behavior by moving away and food searching behavior was disrupted.

While anatomical differences may result in different hearing abilities, other Gadidae have the potential to be surrogate species for Arctic cod. Gadidae have been shown to detect sounds up to about 500 Hz (Popper 2008; Sand and Karlsen 1986). Atlantic cod are well-studied and can serve as model species for fish bioacoustic research (Pine et al. 2020). Atlantic cod (*Gadus morhua*) may detect high-frequency sounds (Astrup and Mohl 1993). Astrup and Møhl (1993) indicated that conditioned Atlantic cod have high frequency thresholds of up to 38 kHz at 185 to 200 dB re 1 μ Pa, which likely only allows for detection of predators at distances no greater than 33–98 ft (10–30 m) (Astrup 1999). A study by Schack et al. (2008) revisited the conclusions from Astrup and Mohl's study, arguing that hearing and behavioral responses in Atlantic cod would be different with unconditioned fish. They found that ultrasound exposures mimicking those of echosounders and odontocetes would not induce acute stress responses in Atlantic cod, and that frequent encounters with ultrasound sources would therefore most likely not induce a chronic state of stress (Schack et al. 2008). The discrepancies between the studies remain unresolved, but it has been suggested the cod in Astrup and Mohl's (1993) study were conditioned to artifacts rather than to the ultrasonic component of the exposure (Astrup 1999; Ladich and Popper 2004; Schack et al. 2008). Additionally, Jørgensen et al. (2005) found that juvenile Atlantic cod did not show any clear behavioral response when exposed to either 1.5 or 4 kHz simulated sonar sound. Therefore, accepted research on cod hearing indicates sensitivities limited to low-frequency sounds.

The effects of specific sources of noise pollution on Gadiformes are unresolved. While Atlantic cod exhibited an initial physiological response to particle motion from an airgun experiment, no behavioral startle response was observed, and fish seemed to habituate with repeated exposure (Davidsen et al. 2019). In a split-brood experiment of vessel noise on larval development, 2 days of both regular and random additional noise reduced growth, while regular noise resulted in faster yolk sac use. After 16 days, larval growth from control and increased noise treatments converged, but fish exposed to regular noise had lower body width-to-length ratios and were easier for predators to catch (Nedelec et al. 2015). Increased noise due to vessel traffic also reduced the spatial range (estimated 3.9–70.9 ft [1.2–21.6 m]) at which Atlantic cod were able to communicate during spawning (Stanley et al. 2017). While this could represent consequences for the affected populations' survival and reproduction, no baseline information exists on the distances cod have evolved to use acoustic communication. A size-structured model analyzed possible population effects of anthropogenic noise on Atlantic cod and found that population growth rates were sensitive to changes in energy expense (Soudijn et al. 2020). The study concluded that persistent levels of sub-lethal anthropogenic noise may lead cod to experience greater energy expenditure, thus indirectly and negatively affecting the age of maturation, fecundity, and survival. However, sources of noise pollution were not analyzed.

3.2.2.4 Essential Fish Habitat

Fish are vital components of the marine ecosystem. They have great ecological and economic aspects. To protect this resource, NMFS works with the regional fishery management councils to identify the

essential habitat for every life stage of each federally-managed species using the best available scientific information. EFH includes all types of aquatic habitat including wetlands, coral reefs, seagrasses, and rivers; all locations where fish spawn, breed, feed, or grow to maturity.

The fisheries of the U.S. are managed within a framework of overlapping international, federal, state, interstate, and tribal authorities. Individual states and territories generally have jurisdiction over fisheries in marine waters within 3 nm (5.6 km) of their coast. Federal jurisdiction includes fisheries in marine waters inside the U.S. Exclusive Economic Zone (EEZ), which encompasses the area from the outer boundary of state waters out to 200 nm (370 km) offshore of any U.S. coastline, except where it intersects closer than 200 nm (370 km) by bordering countries (61 Federal Register [FR] 19390-19429, May 1, 1996). The Study Area resides within the U.S. EEZ, but outside of state jurisdiction.

The Study Area is within the jurisdiction of the North Pacific Fishery Management Council, which is responsible for designating EFH and habitat areas of particular concern for all federally managed species occurring off the coast of Alaska. This council has prepared and implemented a Fishery Management Plan for the Arctic Management Area, which encompasses all marine waters in the U.S. EEZ. This Fishery Management Plan identifies EFH for Arctic cod, saffron cod (*Eleginus gracilis*), and snow crab (*Chionoecetes opilio*). Only EFH for Arctic cod overlaps the Study Area (Figure 3-4). No habitat areas of particular concern have been designated for any species within the Arctic Management Area Fisheries Management Plan (North Pacific Fishery Management Council 2009).

Insufficient information is available to determine EFH for eggs, larvae, and early juveniles of Arctic cod. EFH for late juvenile and adult Arctic cod within the Arctic Management Area occurs in waters from the nearshore to offshore areas along the continental shelf (0-656 ft [0-200 m]) and upper slope (656-1,640 ft [200-500 m]) throughout Arctic waters and often associated with ice floes which may occur in deeper waters (North Pacific Fishery Management Council 2009).




Figure 3-4. Essential Fish Habitat for Arctic Cod

3.2.2.5 Marine Mammals

Nine marine mammal species, which include three cetaceans, five pinnipeds, and the polar bear, are likely to occur in the Study Area during the Proposed Action. Marine mammals are found throughout the Study Area, including on the sea ice and within the water column. All marine mammals are protected under the MMPA, and some are additionally protected under the ESA. Table 3-3 lists the potential marine mammals within the Study Area, their stock, and ESA status. Details about the geographic range, habitat and distribution, and predator/prey interactions of each of these species are provided below.

Table 3-3. Mammals Found in the Study Area during the Proposed Action

<i>Common Name</i>	<i>Scientific Name</i>	<i>Stock(s) within the Study Area</i>	<i>ESA-Listing</i>
ESA-Listed Species			
Bearded seal	<i>Erignathus barbatus nauticus</i> ¹	Beringia	Threatened
Bowhead whale	<i>Balaena mysticetus</i>	Western Arctic	Endangered
Polar bear	<i>Ursus maritimus</i>	Southern Beaufort Sea, Chukchi/Bering Sea	Threatened
Ringed seal	<i>Pusa hispida</i>	Arctic ²	Threatened
Non-ESA Listed Species			
Beluga whale	<i>Delphinapterus leucas</i>	Beaufort Sea, Eastern Chukchi Sea	n/a
Gray whale	<i>Eschrichtius robustus</i>	Eastern North Pacific	n/a
Ribbon seal	<i>Histiophoca fasciata</i>	Alaska	n/a
Pacific walrus	<i>Odobenus rosmarus</i>	n/a	n/a
Spotted seal	<i>Phoca largha</i>	Bering	n/a

¹ Scientific name of subspecies within the Study Area

² Stock is designated under the MMPA.

3.2.2.5.1 ESA-listed Marine Mammals

The ESA-listed marine mammals that may occur in the Study Area are described below.

Bearded Seal

The bearded seal (*Erignathus barbatus*) is listed as threatened under the ESA, and listed as depleted under the MMPA. The bearded seal has been separated into two subspecies: *E. b. barbatus* and *E. b. nauticus*. Only the *E. b. nauticus* subspecies is located within the Study Area. Based on evidence, the *E. b. nauticus* subspecies was further divided into an Okhotsk Distinct Population Segment (DPS) and a Beringia DPS. The Beringia DPS is the only DPS of bearded seal that is located within the Study Area, along the Beaufort Sea continental shelf (Muto et al. 2021). NMFS published a final rule (on December 28, 2012) listing the Beringia and Okhotsk DPS as threatened. Critical habitat has been designated for bearded seals (87 FR 19180; April 1, 2022); however, it is located outside of the Study Area and will not be considered further herein.

Figure 3-5 shows the extent of bearded seals in the Northern Hemisphere. Bearded seals have a circumpolar distribution that does not extend farther north than 85 degrees North latitude (°N) (Muto et al. 2017; Reeves et al. 2002). Beringia bearded seals are widely distributed throughout the northern Bering, Chukchi, and Beaufort Seas and are most abundant north of the ice edge zone (MacIntyre et al. 2013). Telemetry data from Boveng and Cameron (2013) showed that large numbers of bearded seals move south in fall/winter as sea ice forms and move north as the seasonal sea ice melts in the spring. The highest densities of bearded seals are found in the central and northern Bering Sea shelf during

winter (Braham et al. 1981; Burns 1981a; Burns and Frost 1979; Fay 1974; Heptner et al. 1976; Nelson et al. 1984). In late winter and early spring bearded seals are widely (not uniformly) ranging from the Chukchi Sea south to the ice front in the Bering Sea usually on drifting pack ice (Muto et al. 2016). In a shallow water study by MacIntyre et al. (2013), bearded seal calls were recorded throughout the year (11 to 12 months) in the Beaufort Sea, with the peak of calls detected from January to July. Bearded seals inhabit the seasonally ice-covered seas of the Northern Hemisphere, where they whelp and rear their pups, and molt their coats on the ice in the spring and early summer (Muto et al. 2017; Muto et al. 2021).

Bearded seals along the Alaskan coast tend to prefer areas where sea ice covers 70 to 90 percent of the surface, and are most abundant 20 to 100 nm (37 to 185 km) offshore during the spring season (Bengtson et al. 2000; Bengtson et al. 2005; Simpkins et al. 2003). In spring, bearded seals may also concentrate in nearshore pack ice habitats, where females give birth on the most stable areas of ice (Reeves et al. 2002). Bearded seals haul out on spring pack ice (Simpkins et al. 2003) and generally prefer to be near polynyas (areas of open water surrounded by sea ice) and other natural openings in the sea ice for breathing, hauling out, and prey access (Nelson et al. 1984; Stirling 1997). While molting between April and August, bearded seals spend substantially more time hauled out than at other times of the year (Reeves et al. 2002).

In their explorations of the Canada Basin, Harwood et al. (2005) observed bearded seals in waters of less than 656 ft (200 m) during the months from August to September. These sightings were east of 140 degrees West longitude (°W). The Bureau of Ocean Energy Management (BOEM) conducted an aerial survey from June through October that covered the shallow Beaufort and Chukchi Sea shelf waters, and observed bearded seals from Point Barrow to the border of Canada (Clarke et al. 2014). The farthest from shore that bearded seals were observed was the waters of the continental slope.

Bearded seals feed on the seafloor, commonly occupying shallow waters (Fedoseev 2000; Kovacs 2002). The preferred depth range is often described as less than 656 ft (200 m) (Allen and Angliss 2014; Fedoseev 2000; Jefferson et al. 2008; Kovacs 2002), although adults have been known to dive to around 984 ft (300 m) (Cameron and Boveng 2009; Kovacs 2002). At these depths, they feed on demersal fish (e.g., Arctic and saffron cod, flatfish, and sculpins) and a variety of small invertebrates that live in the substrate or on its surface (Fedoseev 2000; Kovacs 2002). They may also opportunistically supplement their diet with crab, shrimp, mollusks, and octopus (Reeves et al. 2002).

Bearded seals may be present close to the continental shelf and therefore, may be present near the Study Area year-round. Designated critical habitat abuts with a small portion of the Study Area in its south westernmost extent.




Figure 3-5. Bearded Seal Distribution in Study Area

Bowhead Whale

The bowhead whale (*Balaena mysticetus*) is listed as endangered under the ESA, and listed as depleted under the MMPA. Bowhead whales that may be present within the Study Area are part of the Bering–Chukchi–Beaufort Seas stock (i.e., Western Arctic stock), which ranges from Siberia east to Amundsen Gulf in Canada to 74 °N, north to the Bering Sea and south to the Pribilof Islands (Figure 3-6). No critical habitat is currently designated for this species.

The bowhead whale is the northernmost of all whales, inhabiting only regions close to the ice edge. Bowhead whales are found in arctic and subarctic regions (55 °N to 75 °N) of the North Atlantic and North Pacific oceans (Rice 1998). Their range can expand and contract depending on ice cover and access to Arctic straits (Rugh et al. 2003). These whales are also found in the Bering, Beaufort, and Chukchi Seas, and the northern parts of Hudson Bay, Canada (Wiig et al. 2007). In Alaska, bowhead whales are closely associated with sea ice most of the year (Moore and Reeves 1993; Quakenbush et al. 2010).

The majority of the Western Arctic stock utilizes wintering areas in the northern Bering and southern Chukchi Seas (which are typically areas with 90 to 100 percent sea ice cover (Citta et al. 2015; Quakenbush et al. 2010); wintering areas are inhabited from December to April. The Western Arctic stock migrates through the Chukchi and Beaufort Seas in the spring (April through May) following fractures in the sea ice around Alaskan coast, generally in the shear zone between the shore-fast and mobile pack ice (Muto et al. 2021). Bowhead whales spend most of the summer (June to September) in the Beaufort Sea before returning again to the Chukchi Sea and then the Bering Sea in the fall (October through December) to overwinter in select shelf waters in all but heavy ice conditions (Braham et al. 1980; Citta et al. 2015; Moore 2000; Moore and Reeves 1993; Quakenbush et al. 2010). Mating occurs from late winter to spring and calving occurs from April to June, both in the Bering Sea (Quakenbush 2008).

Several areas within the Chukchi and Beaufort Seas along the northern coast of Alaska are important to bowhead whales. In the Alaskan Beaufort Sea and northeastern Chukchi Sea, there is a reproductive area that is in use during the month of October. Near Barrow Canyon, there is another area used from April to June for reproduction. In the eastern Chukchi and Alaskan Beaufort Sea, there is a migration area used from April to May. Finally, in the Alaskan Beaufort Sea, there is feeding ground used from September to October, a migration area used from September to October, and a reproduction area used in September (Calambokidis et al. 2015). The feeding grounds used from September to October are focused in the coastal waters of the eastern, central, and western Beaufort Sea (Lowry et al. 2004). Large groups of bowhead whales have been documented feeding in the western Alaskan Beaufort Sea as early as July and continuing into October (Clarke et al. 2014).




Figure 3-6. Bowhead Whale Distribution in the Study Area

Bowheads are one of the most commonly sighted cetaceans in the Chukchi Sea when the ice has receded during warm seasons (Aerts et al. 2013). During summer, most of the population is in relatively ice-free waters in the southeastern Beaufort Sea. Some bowhead whales are found in the western Beaufort, Chukchi, and Bering Seas in summer, and these are thought to be a part of the expanding Western Arctic stock (Citta et al. 2015; Clarke et al. 2013; Clarke et al. 2014; Clarke et al. 2015; Rugh et al. 2003). Summer aerial surveys conducted in the western Beaufort Sea during July and August of 2012-2014 have had relatively high sighting rates of bowhead whales, including cows with calves and feeding animals (Alaska Fisheries Science Center Marine Mammal Laboratory 2014; Clarke et al. 2013; Clarke et al. 2014; Muto et al. 2016). According to the annual Arctic aerial surveys conducted along the north shore of Alaska, the distribution of bowhead whales was primarily on the outer continental shelf (at depths of 167 to 656 ft [51 to 200 m]) in July, on the outer and inner continental shelf (at depths of 0 to 656 ft [0 to 200 m]) in August, and on the inner continental shelf (at depths of less than 164 ft [50 m]) in September. Sighting rate (whales per transect km) by depth zone between 140°W and 154°W in the western Beaufort Sea was highest in the 167 to 656 ft (51 to 200 m) zone in July, then less than or equal to 66 ft (20 m) zone in August, and the 69 to 164 ft (21 to 50 m) zone in September (Clarke et al. 2014). Compared to previous years that also had light sea ice cover, bowhead whale sightings (not normalized by survey effort) in the western Beaufort Sea in fall 2013 were significantly farther from shore and in deeper water in the west. Krutzikowski and Mate (2000) determined the average dive depth of bowhead whales in the Chukchi and Beaufort Sea to be less than 328 ft (100 m) with a maximum recorded dive of 1,155 ft (352 m).

Bowhead whales feed by skimming the surface or sometimes near the seafloor (Rugh and Shelden 2009). Preferred prey include various species of copepods and euphausiids (Budge et al. 2008; Rugh and Shelden 2009; Wiig et al. 2007). Likely or confirmed feeding areas include Amundsen Gulf and the eastern Canadian Beaufort Sea; the central and western U.S. Beaufort Sea; Wrangel Island; and the coast of Chukotka, between Wrangel Island and the Bering Strait (Alaska Fisheries Science Center Marine Mammal Laboratory 2014; Ashjian et al. 2010; Clarke et al. 2013; Clarke et al. 2014; Clarke et al. 2015; Lowry et al. 2004; Muto et al. 2016; Okkonen et al. 2011; Quakenbush et al. 2010).

Bowhead whales are most likely to be present in the Beaufort and Chukchi Seas from March to April and August through October.

Polar Bear

Two polar bear stocks occur within the Study Area: (1) the Southern Beaufort Sea stock and (2) the Chukchi/Bering Seas stock. Both of these stocks are listed as threatened under the ESA (73 FR 28212, May 15, 2009). The determination of polar bears as threatened under the ESA was made based on an extinction risk assessment. This assessment found that the main concern regarding the conservation of polar bears stems from the loss of habitat, particularly sea ice. Polar bears were determined to likely become endangered within the foreseeable future (defined as 45 years) throughout their range, based on expected continued decline of sea ice. Additionally, both stocks are currently listed as depleted and classified as strategic under the MMPA. Designated critical habitat for the polar bear (75 FR 76085; December 7, 2010) encompasses three areas or units: barrier islands, sea ice, and terrestrial denning habitat. The total area designated covers 187,157 mi² (484,734 km²) (Figure 3-7). About 96 percent of the critical habitat area is sea ice. Only a small portion in the southwestern corner of the Study Area overlaps with designated critical habitat.




Figure 3-7. Polar Bear At-Sea Distribution in Study Area

The USFWS identified physical and biological features essential to the conservation of the polar bear. These include:

- Sea ice habitat located over the continental shelf at depths of 984 ft (300 m) or less. In spring and summer, this habitat follows the northward progression of the ice edge as it retreats northward. In fall, this sea ice habitat follows the southward progression of the ice edge as it advances southward.
- Sea ice within 1 mile (mi; 1.6 km) of the mean high tide line of barrier island habitat. Barrier islands are used as migration corridors. Polar bears can move freely between barrier islands by swimming or walking on ice or sand bars, thereby avoiding human disturbance.

The Chukchi/Bering Seas stock is widely distributed on the pack ice in the Chukchi Sea and northern Bering Sea and adjacent coastal areas in Alaska and Russia (USFWS 2021). The stocks overlap seasonally in the eastern Chukchi and western Beaufort Seas (USFWS 2021).

The Southern Beaufort Sea population spends the summer on pack ice and moves toward the coast during fall, winter, and spring (Durner et al. 2004). Polar bears in the Southern Beaufort Sea concentrate in waters less than 984 ft (300 m) deep over the continental shelf and in areas with greater than 50 percent ice cover in all seasons except summer to access prey such as ringed and bearded seals (Durner et al. 2004; Durner et al. 2006b; Durner et al. 2009; Stirling et al. 1999). The eastern boundary of the Southern Beaufort Sea stock occurs south of Banks Island and east of the Baillie Islands, Canada (Amstrup et al. 2000). The western boundary of the Southern Beaufort Sea stock is near Point Hope, Alaska. Polar bears from this population have historically denned on both the sea ice and land. Therefore, the southern boundary of the Southern Beaufort Sea stock is defined by the limits of terrestrial denning sites inland of the coast, which follows the shoreline along the North Slope in Alaska and Canadian Arctic (Bethke et al. 1996). Polar bears could be within the Study Area at any time during the Proposed Action. General year-round distribution of polar bears within the Study Area is depicted in Figure 3-7. The size of a polar bear's range depends on a number of factors, including habitat quality and the amount of available food (Polar Bears International 2015). In the Beaufort Sea, annual polar bear activity areas for individually monitored female bears averaged 57,529 mi² (149,000 km²), ranging from 5,020 to 230,500 mi² (13,000 km² to 597,000 km²) (Amstrup et al. 2000).

Mating occurs in late March through early May. In November and December, females dig maternity dens in pressure ridges in fast ice, drifting pack ice, or on land (up to 100 mi [161 km] inland). Females give birth to their cubs from December to January and stay within their dens until spring (Reeves et al. 2002).

Each year, only 25 percent of reproductively active females produce a litter. Studies conducted between 1981 and 1994 of radio-collared bears found over half of the dens on sea ice (53 percent on pack ice and 4 percent on land fast ice) with the remainder of dens on land. Polar bears do not show fidelity to specific den sites but certain bears do show fidelity to denning on either land or sea ice. The U.S. Geological Survey mapped polar bear dens between 1910 and 2010 using satellite telemetry, very high frequency telemetry, forward-looking infrared, polar bear captures, and reports from coastal Alaskans, hunters, and industry personnel (Durner et al. 2010). The main terrestrial denning areas for the Southern Beaufort Sea population in Alaska occur on the barrier islands from Utqiagvik (Barrow) to Kaktovik and along coastal areas up to 25 mi (40 km) inland, including the Arctic National Wildlife Refuge to Peard Bay, west of Utqiagvik (Barrow) (Amstrup et al. 2000; Amstrup and Gardner 1994; Durner et al. 2001, 2006a). Denning sites in the Beaufort Sea and neighboring regions of Alaska are depicted in Figure 3-7.

Little comprehensive information exists that allows for reliable population estimates of the Chukchi/Bering Seas and Southern Beaufort Sea stocks. The Chukchi Sea population is estimated to

comprise 2,000 animals, based on extrapolation of aerial den surveys (Lunn et al. 2002). Research on the Southern Beaufort Sea population began in 1967 and is one of only four polar bear populations with long term (greater than 20 years) data. The population estimate of 1,526 bears (Regehr et al. 2006), which is based on open population capture-recapture data collected from 2001 to 2006, is considered the most current and valid population estimate (Regehr et al. 2006). The most recent stock assessment for polar bears indicates that the Southern Beaufort Sea stock is declining (Allen and Angliss 2011).

Polar bears' main prey are ringed and bearded seals (Durner et al. 2004; Durner et al. 2006b; Durner et al. 2009; Stirling et al. 1999). Occasionally, polar bears are known to prey upon walruses or beluga whales trapped by ice, and may also consume carrion when prey is scarce (U.S. Fish and Wildlife Service 2021).

Ringed Seal

The ringed seal, specifically the Arctic/Bering Sea subspecies *Pusa hispida hispida*, occurs within the U.S. EEZ of the Beaufort, Chukchi, and Bering Seas and overlaps with the Study Area (Kelly et al. 2009; Palo 2003; Palo et al. 2001). The Arctic ringed seal is listed as threatened under the ESA (77 FR 76706; December 28, 2012). The Arctic/Bering Sea subspecies is listed as depleted and strategic under the MMPA.

Critical habitat for the ringed seal was designated on April 1, 2022 (87 FR 19232) and includes areas of the Bering, Chukchi, and Beaufort Seas, including a small area of overlap with the Study Area (Figure 3-8). A large portion of the Study Area overlaps with a military readiness activity exclusion area for critical habitat, as designated by NMFS in the Final Rule for ringed seal critical habitat. Physical and biological features essential to the conservation of the Arctic ringed seal include:

- Snow-covered sea ice habitat suitable for the formation and maintenance of subnivean birth lairs used for sheltering pups during whelping and nursing, which is defined as waters 10 ft (3 m) or more in depth (relative to mean lower low water [MLLW]) containing areas of seasonal landfast (shorefast) ice or dense, stable pack ice, that have undergone deformation and contain snowdrifts of sufficient depth to form and maintain birth lairs (typically at least 21 in [54 cm] deep).
- Sea ice habitat suitable as a platform for basking and molting, which is defined as areas containing sea ice of 15 percent or more concentration in waters 10 ft (3 m) or more in depth (relative to MLLW).
- Primary prey resources to support Arctic ringed seals, which are defined to be small, often schooling, fishes, in particular Arctic cod, saffron cod, and rainbow smelt; and small crustaceans, in particular, shrimps and amphipods.




Figure 3-8. Ringed Seal Distribution in Study Area

The area designated as critical habitat was deemed to have one or more of these essential features that may require special management or protection. Critical habitat includes areas with physical or biological features that support functions of whelping and nursing, where birth lairs are constructed and maintained, and molting. The specific geographic locations of essential sea ice habitat vary annually, or even daily, depending on many factors, including time of year, local weather, and oceanographic conditions. In addition, the duration that any given location has sea ice habitat essential for birth lairs or for molting can vary annually depending on the rate of ice melt and other factors. The southern boundary suggests that sea ice essential for Arctic ringed seal birth lairs extends to some point south of St. Matthew Island and Nunivak Island. Given the inherent variability in the spatial distribution of sea ice and the widespread distribution of Arctic ringed seals, including in offshore pack ice, the northern and eastern boundaries of the area were identified as the outer extent of the U.S. exclusive economic zone (Figure 3-8). NMFS determined that the essential features of the habitat of the Arctic ringed seal may require special management considerations or protection in the future to minimize the risks posed to these features by potential shipping and transportation activities, because: (1) both the physical disturbance and noise associated with these activities could displace seals from favored habitat that contains the essential features, thus altering the quantity and/or quality of these features; and (2) in the event of an oil spill, sea ice essential for birth lairs and for molting could become oiled, and the quantity and/or quality of the primary prey resources could be adversely affected.

Ringed seals are the most common pinniped in the Study Area and have wide distribution in seasonally and permanently ice-covered waters of the Northern Hemisphere (North Atlantic Marine Mammal Commission 2004). Throughout their range, ringed seals have an affinity for ice-covered waters and are well adapted to occupying both shore-fast and pack ice (Kelly 1988c). Ringed seals can be found further offshore than other pinnipeds since they can maintain breathing holes in ice thickness greater than 6.6 ft (2 m) (Smith and Stirling 1975). Breathing holes are maintained by ringed seals' sharp teeth and claws on their fore flippers. They remain in contact with ice most of the year and use it as a platform for molting in late spring to early summer, for pupping and nursing in late winter to early spring, and for resting at other times of the year (Muto et al. 2017).

Ringed seals have at least two distinct types of subnivean lairs: haulout lairs and birthing lairs (Smith and Stirling 1975). Haulout lairs are typically single-chambered and offer protection from predators and cold weather. Birthing lairs are larger, multi-chambered areas that are used for pupping in addition to protection from predators. Ringed seals pup on both land-fast ice as well as stable pack ice. Lentfer (1972) found that ringed seals north of Barrow, Alaska (Figure 3-8), build their subnivean lairs on the pack ice near pressure ridges. Since subnivean lairs were found north of Barrow, Alaska, in pack ice, they are also assumed to be found within the sea ice in the Study Area. Ringed seals excavate subnivean lairs in drifts over their breathing holes in the ice, in which they rest, give birth, and nurse their pups for 5–9 weeks during late winter and spring (Chapskii 1940; McLaren 1958; Smith and Stirling 1975). Snow depths of at least 20–26 in (50–65 cm) are required for functional birth lairs (Kelly 1988a; Lydersen 1998; Lydersen and Gjertz 1986; Smith and Stirling 1975), and such depths typically are found only where 8–12 in (20–30 cm) or more of snow has accumulated on flat ice and then drifted along pressure ridges or ice hummocks (Hammill 2008; Lydersen et al. 1990; Lydersen and Ryg 1991; Smith and Lydersen 1991). Ringed seals are born beginning in March, but the majority of births occur in early April. About a month after parturition, mating begins in late April and early May.

In Alaskan waters, during winter and early spring when sea ice is at its maximal extent, ringed seals are abundant in the northern Bering Sea, Norton and Kotzebue Sounds, and throughout the Chukchi and Beaufort Seas (Frost 1985; Kelly 1988c) and, therefore, are in the Study Area (Figure 3-8). Passive acoustic monitoring of ringed seals from a high frequency recording package deployed at a depth of

787 ft (240 m) in the Chukchi Sea, 65 nm (120 km) north-northwest of Barrow, Alaska detected ringed seals in the area between mid-December and late May over the four year study (Jones et al. 2014). With the onset of the fall freeze, ringed seal movements become increasingly restricted and seals will either move west and south with the advancing ice pack with many seals dispersing throughout the Chukchi and Bering Seas, or remain in the Beaufort Sea (Crawford et al. 2012; Frost and Lowry 1984; Harwood et al. 2012). Kelly et al. (2010a) tracked home ranges for ringed seals in the subnivean period (using shorefast ice); the size of the home ranges varied from less than 0.39 up to 10.8 mi² (1 up to 27.9 km²); (median is 0.24 mi² [0.62 km²] for adult males and 0.25 mi² [0.65 km²] for adult females). Most (94 percent) of the home ranges were less than 1.15 mi² (3 km²) during the subnivean period (Kelly et al. 2010a). Near large polynyas, ringed seals maintain ranges, up to 2,703 mi² (7,000 km²) during winter and 811 mi² (2,100 km²) during spring (Born et al. 2004). Some adult ringed seals return to the same small home ranges they occupied during the previous winter (Kelly et al. 2010a). The size of winter home ranges can vary by up to a factor of 10 depending on the amount of fast ice; seal movements were more restricted during winters with extensive fast ice, and were much less restricted where fast ice did not form at high levels (Harwood et al. 2015). Ringed seals may occur within the Study Area throughout the year and during the Proposed Action.

Ringed seal population surveys in Alaska have used various methods and assumptions, had incomplete coverage of their habitats and range, and were conducted more than a decade ago; therefore, current, comprehensive, and reliable abundance estimates or trends for the Alaska stock are not available (Muto et al. 2021). Although a reliable population estimate is not available for the entire stock, survey methods have been developed and applied to substantial portions of the Arctic stock's range in U.S. waters. Frost et al. (2004) conducted surveys within 21.6 nm (40 km) of shore in the Alaska Beaufort Sea during May-June 1996-1999, and observed ringed seal densities ranging from 0.81 seal per km² in 1996 to 1.17 seals per km² in 1999. Moulton et al. (2002) conducted similar, concurrent surveys in the Alaska Beaufort Sea during 1997-1999 but reported substantially lower ringed seal densities (0.43, 0.39, and 0.63 seals per km² in 1997-1999, respectively) than Frost et al. (2004). Using the most recent estimates from surveys by Bengtson et al. (2005) and Frost et al. (2004) in the late 1990s and 2000, Kelly et al. (2010b) estimated the total population in the Alaska Chukchi and Beaufort seas to be at least 300,000 ringed seals, which Kelly et al. (2010b) states is likely an underestimate since the Beaufort surveys were limited to within 21.6 nm (40 km) of shore.

In general, ringed seals prey upon fish and crustaceans. Ringed seals are known to consume up to 72 different species in their diet; their preferred prey species is the polar cod (Jefferson et al. 2008). Ringed seals also prey upon a variety of other members of the cod family, including Arctic cod (Holst et al. 2001), and saffron cod, with the latter being particularly important during the summer months in Alaskan waters (Lowry et al. 1980). Invertebrate prey seems to become prevalent in the ringed seals diet during the open-water season and often dominates the diet of young animals (Holst et al. 2001; Lowry et al. 1980). Large amphipods (e.g., *Themisto libellula*), krill (e.g., *Thysanoessa inermis*), mysids (e.g., *Mysis oculata*), shrimps (e.g., *Pandalus* spp., *Eualus* spp., *Lebbeus polaris*, and *Crangon septemspinosa*), and cephalopods (e.g., *Gonatus* spp.) are also consumed by ringed seals.

3.2.2.5.2 Non ESA-listed Marine Mammals

Marine mammals that may occur in the Study Area, and are not listed under the ESA, are described below.

Beluga Whale

In the United States and Canada, individual populations have been assessed for status under the applicable conservation statutes. Five stocks of beluga whales are recognized within U.S. waters:

(1) Cook Inlet, (2) Bristol Bay, (3) Eastern Bering Sea, (4) Eastern Chukchi Sea, and (5) Beaufort Sea. Only the Cook Inlet population of the beluga whale is listed as endangered under the ESA. The Beaufort Sea, Eastern Chukchi Sea, Eastern Bering Sea, and Bristol Bay stocks of beluga whales are not listed as threatened or endangered under the ESA. Additionally, those stocks not listed as threatened or endangered under the ESA are not listed as depleted or classified as strategic under the MMPA. Only the Beaufort Sea and Eastern Chukchi Sea stocks of beluga whales are expected to occur in the Study Area.

The majority of belugas are distributed discontinuously around the Arctic Ocean and adjacent seas, primarily on the continental shelf and near coasts around North America, Russia, and Greenland (Rice 1998). Beluga whales are found primarily in shallow coastal waters (in depths as shallow as 3 to 10 ft [1 to 3 m]), but can be found in waters deeper than 2,624 ft (800 m) (Jefferson et al. 2012; Richard et al. 2001). Beluga whales are distributed throughout the seasonally ice-covered arctic and subarctic waters of the Northern Hemisphere (Gurevich 1980). Their range includes Greenland, the Arctic coast of Eurasia and central Asia, the Arctic coast of Siberia to the Bering Sea, the Arctic coast of Alaska and northwestern Canada, and from the Chukchi Sea east to the Beaufort Sea. Disjoined populations are located in the St. Lawrence estuary, Sea of Okhotsk, Cook Inlet, and northern Gulf of Alaska.

Belugas are both migratory and residential (non-migratory), depending on the population. Migratory populations move between seasonal ranges. During winter, migratory belugas can be found foraging around the pack ice. When the sea ice melts in summer, they move to warmer river estuaries and coastal areas for molting and calving (Muto et al. 2017; Muto et al. 2021). These annual migrations can span over thousands of kilometers (Frost et al. 1985). Irregular sea ice conditions during the spring and summer months can influence beluga whales to adjust their migratory tracks to summering areas (O'Corry-Crowe et al. 2016). Habitat selection can differ between sexes, with males selecting areas with higher ice concentrations whereas females prefer to remain proximal (less than 124 mi [200 km]) to the shore during summers and September migrations (Hauser et al. 2017). There are two migration areas used by belugas that overlap the Study Area. One, located in the Eastern Chukchi and Alaskan Beaufort Sea, is a migration area in use from April to May. The second, located in the Alaskan Beaufort Sea, is used by migrating belugas from September to October (Calambokidis et al. 2015). The residential populations participate in short distance movements within their range throughout the year. Seasonal distribution is affected by ice cover, tidal conditions, prey availability, temperature, and human interaction (Frost et al. 1985). Belugas are closely associated with open leads and polynyas in ice-covered regions (Hazard 1988).

Near the Study Area, beluga whales may spend summer in both offshore and coastal waters, with concentrations in Kasegaluk Lagoon (on the north slope of Alaska) and the Mackenzie Delta (in the Beaufort Sea) (Hazard 1988). Most beluga whales from these summering areas overwinter in the Bering Sea, excluding those found in the northern Gulf of Alaska (Shelden 1994). The Eastern Chukchi Sea belugas move into coastal areas along Kasegaluk Lagoon in late June and remain in the area until mid-July (Frost and Lowry 1990; Frost et al. 1993). Telemetry tags attached to belugas within Kasegaluk Lagoon in June and July of 1998 showed that whales traveled 594 nm (1,100 km) north of the Alaska coastline and to the Canadian Beaufort Sea within three months (Suydam et al. 2001), which indicated an overlap in distribution with the Beaufort Sea stock of beluga whales. Adult males appear to use deep waters rather than shallow shelf areas and remain in these deep waters for the duration of the summer. All belugas that moved into the Arctic Ocean (north of 75 °N) were males that can travel through 90 percent pack ice cover to reach deeper waters of the Beaufort Sea and Arctic Ocean (approximately 79 to 80 °N) by late July/early August, while the adult and immature females remain at or near the shelf break of the Chukchi Sea. After October, the remaining females in the Chukchi Sea migrate south, through the Bering Strait into the Bering Sea north of Saint Lawrence Island, which suggests that some

belugas that summer in the eastern Chukchi Sea overwinter in the waters north of Saint Lawrence Island (Suydam 2009).

The Beaufort Sea beluga whale stock range includes the Alaska north coast and the Canadian Arctic Archipelago northward to the pack-ice (Department of Fisheries and Oceans 2000). Beaufort Sea belugas congregate in the Mackenzie Estuary in early summer. Later in summer, belugas move eastward toward Canada. By mid-August and early September, belugas begin their migration westward along the Alaska coast and far offshore to the pack-ice of the Chukchi Sea. The winter range is thought to include the offshore areas of the Chukchi and Bering Seas.

Belugas are opportunistic feeders that vary their diets according to their location and the season. Fish (eulachon, salmon, capelin, cod, herring, smelt, flounder, sole, lamprey and lingcod), crustaceans (crab, clams, mussels and shrimp) and other deep-sea invertebrates (octopus and squid) are their main prey (Reeves et al. 2002). Belugas are shallow divers with typical dives of about 66 ft (20 m) or less (Goetz et al. 2012). Goetz et al. (2012) recorded belugas in the Cook Inlet of Alaska diving to mean depths ranging from (5.2 to 22 ft (1.6 to 6.7 m) with mean durations ranging from 1.1 to 6.9 minutes.

According to the annual BOEM aerial surveys along the north coast of Alaska, beluga distribution in the western Beaufort Sea was centered over the continental slope and Barrow Canyon, with few sightings nearshore. There were more beluga whales in the Chukchi Sea than the Beaufort Sea (Clarke et al. 2014).

Beluga whales may be present within the Beaufort Sea during the summer.

Gray Whale

Two genetically distinct populations of Pacific gray whales (*Eschrichtius robustus*) are currently recognized (Reilly et al. 2008): (1) the Eastern North Pacific stock and (2) the Western North Pacific stock (Bonner 1986; LeDuc et al. 2002; Weller et al. 2013). Although the Western North Pacific stock is listed as endangered under the ESA and depleted under the MMPA, only the Eastern North Pacific stock is expected to be in the Study Area. The Eastern North Pacific stock is not listed under the ESA.

The Eastern North Pacific stock lives along the West Coast of North America (Rice 1981; Rice et al. 1984; Swartz et al. 2006). Gray whale occurrence is primarily in shallow waters over the continental shelf. Breeding and calving are seasonal and closely synchronized with migratory timing. An important area for reproduction stretches from Point Lay to Point Barrow, west of the Study Area, and is in use from June to September. Gray whale migration typically follows the coastline (within 1.1 nm [2 km] of the coast), except when crossing major bays, straits and inlets from southeastern Alaska to the eastern Bering Sea. The northbound migration from low latitude winter calving grounds in Mexico begins about mid-February (Rice and Wolman 1971). In summer and fall, gray whales feed in the Chukchi, Beaufort, and northwestern Bering Seas and are among the most commonly observed cetaceans in the Chukchi Sea (Aerts et al. 2013; Carretta et al. 2021). Then, by late November, the southbound migration is underway as whales begin to travel from summer feeding areas to winter calving areas off the West Coast of Baja California, Mexico, and the southeastern Gulf of California (Rugh et al. 2001; Swartz et al. 2006). Migrating whales move southward through the Unimak Pass and follow a shoreline route to the winter grounds (Rice 1998). Gray whales typically migrate to nearly landlocked lagoons and bays in Baja California, Mexico and give birth to calves between January and mid-February (Rice et al. 1981). Gray whale feeding grounds are generally in waters less than 223 ft (68 m) in depth. An important feeding ground for gray whales stretches from Point Lay to Point Barrow, west of the Study Area, and is in use from June to October. During summer and fall, most whales in the Eastern North Pacific stock feed in the Chukchi and Beaufort Seas, between 174 degrees East longitude (°E) and 130 °W, and the northwestern Bering Sea south to Russia (Rice 1998).

Prey of gray whales consists primarily of swarming mysids, polychaete tube worms, and amphipods in the northern parts of their range (Jefferson et al. 2008). They will also take crabs, baitfish, and other food opportunistically. Killer whales (*Orcinus orca*) are the only known non-human predator to the gray whale. Gray whales feed by swimming slowly over the seafloor at depths up to 197 ft (60 m) (Golda 2015).

During the annual BOEM Arctic survey, gray whales were observed east of Point Barrow in August and October. However, primarily they were seen nearshore and west of the Study Area. Gray whales may be present in the Beaufort Sea in the late summer and early fall, but are unlikely to occur within the deeper waters of the Study Area.

Ribbon Seal

The ribbon seal (*Histriophoca fasciata*) does not have any subspecies, and is therefore considered a single species throughout its range. Ribbon seals are not listed under the ESA, although the species is a Species of Concern. Ribbon seals are protected under the MMPA.

The ribbon seal's range includes the Sea of Okhotsk, Bering Sea, and southern Chukchi Sea (Reeves et al. 2002). Their range stretches throughout the Bering Sea, including the Aleutian Islands, the western Pacific around the Kamchatka Peninsula and Kuril Islands (Russia), as well as the Sea of Okhotsk. The southern distribution within their effective range is strongly associated with the extent of ice formation in the Bering Sea and Sea of Okhotsk, which can drive large numbers of these seals further south in years with heavy ice. The inverse is also true; years of light ice formation causes greater numbers of seals to remain further north. The northernmost record of a ribbon seal was in the western Beaufort Sea, which is considered outside of the typical range of the ribbon seal, in August of 1983.

Ribbon seals are found in the open sea and on the free-floating pack ice rather than shore-fast ice (Kelly 1988b). From late March to early May, ribbon seals inhabit the Bering Sea ice front (Braham et al. 1984; Burns 1970, 1981b) and are most abundant in the central and western parts of the Bering Sea along the southern edge of the ice front (Burns 1970, 1981b). As the ice front recedes, most seals move further north in the Bering Sea between May and mid-July, using the ice edge or ice remnants to haul out (Burns 1970, 1981b; Burns et al. 1981). The Bering Sea and Sea of Okhotsk are the principal breeding grounds for this species (Reeves et al. 2002). During summer, from July through October, these seals do not occur near shore, nor do they migrate northward to the fringe of polar ice as do bearded and ringed seals. Although their distribution is not completely understood, the most likely explanation is that they spend the summer at sea. During open water seasons, the Beaufort Sea shelf, Northern Bering Strait, and Southern Chukchi Sea are ecologically important areas for ribbon seals (Frouin-Mouy et al. 2019). Using passive acoustic data, the northeastern Chukchi Sea was identified as a possible part of a migration corridor between the Chukchi Plateau and the Beaufort Sea (Frouin-Mouy et al. 2019). A 2005 study using satellite telemetry has shown that animals tagged near the eastern coast of the Kamchatka Peninsula (Russia) spent the summer and fall in the Bering Sea and Aleutian Islands, while others moved from the central Bering Sea to the Bering Strait, Chukchi Sea, or Arctic Basin in a 2010 study as the seasonal ice receded (Boveng et al. 2008; Muto et al. 2021). In Alaskan waters, ribbon seals range northward from Bristol Bay in the Bering Sea into the Chukchi and western Beaufort Seas (Muto et al. 2017; Muto et al. 2021). Little is known about the range of ribbon seals during the rest of the year. In their explorations of the Canada Basin, Harwood et al. (2005) observed ribbon seals east of 140 °W from the coast to waters as deep as 9,843 ft (3,000 m).

Ribbon seals in the Bering Sea and Sea of Okhotsk consume 35 different species of fish and invertebrates (Jefferson et al. 2008). Pollock and Arctic cod are among the known prey species for the ribbon seal (Reeves et al. 2002). Juvenile ribbon seals feed on euphausiids after weaning until they reach one year

of age when they feed predominantly on shrimp for one year (Jefferson et al. 2008). In the Bering Sea, 65 percent of the ribbon seal's diet consists of squid and octopus. Potential predators include polar bears, killer whales, sharks, and walrus. Ribbon seals often dive to depths of 656 ft (200 m) while foraging, and have been recorded diving to depths greater than 1,969 ft (600 m) (London et al. 2015).

Ribbon seals are typically close to shore, but may be rarely encountered in the Beaufort Sea in the summer and fall.

Pacific Walrus

The Pacific walrus (*Odobenus rosmarus*) is the only subspecies occurring in U.S. waters, and is considered a single stock. On October 4, 2017, the Pacific walrus was removed as a candidate species by the USFWS, and determined that the population did not warrant listing at this time. On June 3, 2021, the U.S. 9th Circuit Court of Appeals issued a decision in a case brought by the Center for Biological Diversity finding that the USFWS had not sufficiently explained why it changed its prior position on the need to list the walrus under the ESA. The matter was remanded to the USFWS with instructions for the Service to provide a sufficient explanation for its conclusion that listing was no longer warranted (Center for Biological Diversity v. Haaland, No. 19-35981 (9th Cir. 2021)). The Pacific walrus may be under consideration for federal protection again, but a final decision has not been issued by the USFWS. Additionally, the Pacific walrus within the U.S. EEZ is not designated as depleted under the MMPA, but is classified as strategic because the level of human-caused mortality exceeds the rate of reproduction and survival required for a stable population. The walrus is managed by the USFWS.

Walrus have a circumpolar distribution in the Arctic Ocean and are associated with pack ice everywhere they are found, at least during winter. Pacific walrus range throughout the continental shelf waters of the Bering and Chukchi Seas, occasionally moving into the East Siberian Sea and the Beaufort Sea (Muto et al. 2017). A significant proportion of the Pacific walrus population migrates into the Chukchi Sea region each summer. Walrus are known to stay fairly close to land for most of their lives and make shallow dives inshore (depths of roughly 262 ft [80 m]) (Kastelein 2002) from the continental shelf and slope, so they do not regularly occur in deep oceanic waters. Walrus haul out on ice floes and sandy beaches or rocky shores, along remote stretches of mainland coastlines or islands (Jefferson et al. 2008; Kastelein 2009). Walrus haul out on land to a greater extent during years with reduced pack ice. The movements of walrus generally follow the movements of pack ice. However, some individuals do travel farther from the ice during summer months.

The shallow, productive, ice covered waters of the eastern Chukchi Sea are considered particularly important habitat for female walrus and their dependent young. Several thousand animals (primarily adult males) aggregate near coastal haulouts in the Bering Strait region and Bristol Bay, as well as several areas near Russia and Japan. During the late winter breeding season, most walrus are found in two major Bering Sea concentration areas where open leads, polynyas, or thin ice allows open water access (Fay et al. 1984; Garlich-Miller et al. 2011). While the specific location of these groupings can vary annually and seasonally depending upon the extent of the sea ice, one group will generally range from the Gulf of Anadyr in Russia into a region southwest of St. Lawrence Island (northern Bering Sea), and the second group will aggregate somewhere in the southeastern Bering Sea from south of Nunivak Island to northwestern portions of Bristol Bay.




Figure 3-9. Pacific Walrus Distribution Near the Study Area

In the annual BOEM survey, only a few walrus were observed east of Barrow, and then, only in shallow waters in August (Clarke et al. 2014; Kuletz et al. 2015). In their explorations of the Canada Basin, Harwood et al. (2005) only saw walrus in the Chukchi Sea at the Chukchi shelf break and at the Northwind Ridge, located just east of 160 °W.

Walrus feed on bottom-dwelling invertebrates and slow-moving fish to depths of roughly 262 ft (80 m) (Kastelein 2002). Preferred prey include clams, snails, shrimp and slow-moving fish (Jefferson et al. 1993). Walrus have been observed preying on seabirds, seals, and Northern sea lions (Reeves et al. 2002). Walrus are known to consume between 88 and 176 pounds (lbs; 40 and 80 kilograms [kg]) of food per day (Jefferson et al. 2008; Kastelein and Wiepkema 1989). Common predators to the walrus are killer whales and polar bears.

Pacific walrus may rarely be encountered in the Study Area, though it is unlikely.

Spotted Seal

Within the Study Area, spotted seals (*Phoca largha*) are not listed as threatened or endangered under the ESA. The Bering Sea DPS is located off the coast of Alaska within the Study Area. Spotted seals are protected under the MMPA.

Spotted seals are widespread in the Sea of Okhotsk, Yellow, Japan, and Bering Seas. Spotted seals are closely related to and are often mistaken for Pacific harbor seals. The two species are often seen together and are partially sympatric with range overlap in the southern part of the Bering Sea (Quakenbush 1988). The key difference between the two species is spotted seals breed earlier than harbor seals and they are noticeably less social during the breeding season. Additionally, spotted seals are strongly associated with pack ice whereas harbor seals are not (Quakenbush 1988; Shaughnessy and Fay 1977).

Spotted seals inhabit the southern edges of the pack ice in the Chukchi Sea from winter to early summer. Spotted seals also overwinter in the Bering Sea, tending to remain associated with the ice edge and moving in an east to west direction (Lowry et al. 1998). To the south, and along the West Coast of Alaska, spotted seals can be found at the Pribilof Islands (in the Bering Sea), in Bristol Bay, and along the eastern Aleutian Islands. As mentioned above, a large percent of haulouts are associated with pack ice and their movements tend to remain associated with ice.

Breeding takes place on pack ice from January to mid-April, with the peak of pups born in mid to late March. Eight offshore breeding areas have been described, three of which are in the Bering Sea (Shaughnessy and Fay 1977). The seals remain at the breeding sites until the end of the breeding season which coincides with the break-up of ice in spring.

As ice begins to break up in the Bering Sea, seals follow the retreating ice edge and disperse northward along the shores of Alaska and Siberia (Bigg 1981). During spring, spotted seals tend to prefer the small, broken up floes (i.e., less than 66 ft [20 m] in diameter) and remain at the southern margin of the ice in areas where the water depth does not exceed 656 ft (200 m). Once the sea ice retreats in early summer, seals move to coastal habitats, including the mouths of rivers, where they remain until the fall (Fay 1974; Lowry et al. 2000; Shaughnessy and Fay 1977; Simpkins et al. 2003). In the summer and fall, spotted seals occupy coastal haulouts regularly using sand bars and beaches as resting places between longer foraging periods at sea (Frost et al. 1993; Lowry et al. 1998), and can be found as far north as 69 to 72 °N in the Chukchi and Beaufort Seas (Porsild 1945; Shaughnessy and Fay 1977). When the cold season begins, some seals in the northeastern Chukchi Sea move south in October and pass through the Bering Strait during November (Porsild 1945; Shaughnessy and Fay 1977).

Spotted seals feed opportunistically on a variety of fish, cephalopods, and crustaceans (Bigg 1981). While juveniles and adults eat a variety of schooling fish (pollock, capelin, Arctic cod and herring), epibenthic fish (especially flounder, halibut and sculpin), crabs, and octopus at depths up to 984 ft (300 m) (Reeves et al. 2002), pups feed on small amphipods found around ice floes. Predators of spotted seals include Pacific sleeper sharks, killer whales, polar and brown bears, walruses and Steller sea lions; predators to pups include golden eagles (*Aquila chrysaetos*), Steller's sea eagles (*Haliaeetus pelagicus*), ravens, gulls, and Arctic foxes (*Vulpes lagopus*) (Quakenbush 1988).

Spotted seals typically remain close to shorelines, but may be encountered in the Beaufort Sea during the summer and fall.

3.2.2.5.3 Marine Mammal Hearing

All marine mammals studied can use sound to forage, orient, socially interact with others, and detect and respond to predators. Measurements of marine mammal sound production and hearing capabilities provide some basis for assessment of whether exposure to a particular sound source may affect a marine mammal behaviorally or physiologically.

The hearing mechanism for marine mammals is similar to that of terrestrial mammals. It is comprised of an outer ear, a fluid-filled inner ear with a frequency-tuned membrane interacting with sensory cells, and an air-filled middle ear, which provides a connection between the outer ear and inner ear (Nedwell et al. 2004). The discussion on hearing below is broken down into the hearing groups of each species within the Study Area. Hearing groups for each species is shown in Table 3-4 below.

The typical terrestrial mammalian ear (which is ancestral to that of marine mammals) consists of an outer ear that collects and transfers sound to the tympanic membrane and then to the middle ear (Popper and Fay 1994; Rosowski 1994). The middle ear contains ossicles that amplify and transfer acoustic energy to the sensory cells (called hair cells) in the cochlea, which transforms acoustic energy into electrical neural impulses that are transferred by the auditory nerve to high levels in the brain (Møller 2012). All marine mammals display some degree of modification to the terrestrial ear; however, there are differences in the hearing mechanisms of marine mammals with an amphibious ear versus those with a fully aquatic ear (Wartzok and Ketten 1999). Marine mammals with an amphibious ear include the marine carnivores: pinnipeds, sea otters, and polar bears (Ghoul and Reichmuth 2014; Owen and Bowles 2011; Reichmuth et al. 2013). Outer ear adaptations in this group include external pinnae (ears) that are reduced or absent, and in the pinnipeds, cavernous tissue, muscle, and cartilaginous valves seal off water from entering the auditory canal when submerged (Wartzok and Ketten 1999). Marine mammals with the fully aquatic ear (cetaceans and sirenians) use bone and fat channels in the head to conduct sound to the ear; while the auditory canal still exists in pinnipeds, it is narrow and sealed with wax and debris, and external pinnae are absent (Ketten 1998).

For this analysis, marine mammals are arranged into the following functional hearing groups based on their generalized hearing sensitivities: mid-frequency cetaceans (odontocetes), low-frequency cetaceans (mysticetes), otariids and other non-phocid marine carnivores in water and air (walruses, polar bears), and phocids in water and air (true seals). Note that the designations of mid- and low-frequency cetaceans are based on relative differences of hearing sensitivity between groups, as opposed to conventions used to describe active sonar systems. For discussion of all marine mammal functional hearing groups and their derivation, see technical report Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects (U.S. Department of the Navy 2017a).

Table 3-4. Species in Marine Mammal Hearing Groups Potentially Within the Study Area

<i>Functional Hearing Group</i>	<i>Species in the Study Area</i>
Low-frequency cetaceans (mysticetes)	Bowhead whale
	Gray whale
Mid-frequency cetaceans (odontocetes)	Beluga whale
Phocids	Bearded seal
	Ribbon seal
	Spotted seal
	Ringed seal
Polar bear	Polar bear
Odobenids	Pacific walrus

Mysticete/Low-Frequency Cetacean Hearing

Anatomical and paleontological evidence suggests that the inner ears of mysticetes (baleen whales) are well adapted for hearing at lower frequencies (Ketten 1998; Richardson et al. 1995b). Functional hearing in low-frequency cetaceans is conservatively estimated to be between about 7 Hz and 22 kHz (Southall et al. 2007). In updating their 2007 hearing range determinations, Southall et al. (2019) estimated functional hearing range for this group at 30 Hz to 30 kHz with peak sensitivity between 1 and 8 kHz, although the authors noted that there are no direct measurements of species in this group. Their determinations were based on anatomical and physiological parameters of the group’s species as well as extrapolation from other hearing groups.

Non-biosonar communication signals span a wide frequency range, primarily having energy up into the tens of kilohertz range. Of particular note are the very low-frequency calls of mysticete whales that range from tens of hertz to several kilohertz, and have source levels of 150 to 200 dB re 1 μ Pa (Cummings and Thompson 1971; Edds-Walton 1997; Širović et al. 2007; Stimpert et al. 2007; Wartzok and Ketten 1999). These calls most likely serve social functions such as mate attraction, but may serve an orientation function as well (Green 1994; Green et al. 1994; Richardson et al. 1995b).

Odontocete/Mid-Frequency Cetacean Hearing

In general, odontocete hearing is very broad, including low-frequency, mid-frequency, and high-frequency cetaceans. Beluga whales are members of the mid-frequency cetacean functional hearing group, which also includes 32 species of dolphins and sperm whales. Functional hearing in mid-frequency cetaceans is conservatively estimated to be between 150 Hz and 160 kHz (Southall et al. 2007) with best hearing sensitivity at frequencies of several tens of kilohertz or higher (Southall et al. 2019). Mid-frequency cetaceans also generate short-duration (50-200 microseconds) specialized clicks used in echolocation with peak frequencies between 10 and 200 kHz (Au 1993; Wartzok and Ketten 1999). Echolocation is used to detect, localize, and characterize underwater objects, including prey items (Au 1993). These clicks are often more intense than other communicative signals, with reported source levels as high as 229 dB re 1 μ Pa at 1 m peak-to-peak (Au et al. 1974). Castellote et al. (2014) found that wild beluga whales can hear in the range of 4 to 150 kHz. Klishin et al. (2000) tested a single beluga whale and found its hearing to be most sensitive from 32 kHz to 108 kHz.

Phocid Hearing

Phocids can make calls between 90 Hz and 16 kHz (Richardson et al. 1995b). The generalized hearing for phocids (underwater) ranges from 50 Hz to 86 kHz (National Marine Fisheries Service 2016), which includes the suggested auditory bandwidth for pinnipeds in water proposed by Southall et al. (2007),

ranging between 75 Hz to 75 kHz. Phocid functional hearing in air is estimated to be 75 Hz to 30 kHz (Carretta et al. 2008; Kastak et al. 1999; Kastelein et al. 2009a; Kastelein et al. 2009b; Møhl 1968a, 1968b; Reichmuth 2008; Terhune and Ronald 1971, 1972).

Polar Bear Hearing

Airborne hearing threshold measurements of polar bears have shown best hearing sensitivity between 8 and 14 kHz, with a rapid decline in sensitivity below 125 Hz and above 20 kHz (Bowles et al. 2008; Nachtigall et al. 2007; Owen and Bowles 2011). Like the pinnipeds, polar bears are amphibious mammals in the order Carnivora. However, unlike pinnipeds, polar bears spend only a few minutes submerged and spend the majority of their time above water, thus limiting any potential for acoustic exposure. Additionally, the polar bear ear is very similar to the otariid ear and therefore the polar bear is placed within the same hearing group as otariids (Nummela 2008a; Nummela 2008b). Hearing limits are 50 Hz to 35 kHz in air and 50 Hz to 50 kHz in water (Southall et al. 2007).

Odobenid Hearing

The walrus is the only extant Odobenid pinniped and may be found within the Study Area. Walruses react to airborne sounds at 250 Hz to 8 kHz, but absolute thresholds were not determined (Kastelein et al. 1993). The walrus is adapted to low-frequency sound with a range of best hearing in water from 1 to 12 kHz; its hearing ability falls off sharply at frequencies above 14 kHz (Kastelein 2002; Kastelein et al. 1996). Walrus hearing sensitivity is most similar to otariids, and therefore the walrus is assigned the same functional hearing range as otariids for this analysis. Functional hearing limits are conservatively estimated to be 50 Hz to 35 kHz in air and 50 Hz to 50 kHz in water (Southall et al. 2007). Walrus produce low frequency (100-1,200 Hz) sounds including barks (females) and bell sounds and whistles (males), as well as some grunts, guttural sounds, and roars (Charrier et al. 2010; Richardson et al. 1995b). Hearing in odobenids is very similar to that of Otariids (sea lions and fur seals).

4 Environmental Consequences

This chapter discusses the potential environmental consequences of the Proposed Action to the natural and physical environments described in Chapter 3. Stressors resulting from the Proposed Action that may potentially harm the biological environment include:

- Acoustic: non-impulsive acoustic sources, icebreaking noise, and vessel noise
- Physical: icebreaking (physical impacts), vessel and in-water device strike, and bottom disturbance
- Expended Material: entanglement and ingestion

Appendix A provides a description of each stressor, as well as matrices showing which activities generate each stressor and what resources are impacted by each stressor.

4.1 Stressors Associated with the Proposed Action

4.1.1 Acoustic Stressors

The acoustic stressors from the Proposed Action include non-impulsive acoustic sources, icebreaking noise, and vessel noise.

4.1.1.1 Non-Impulsive Acoustic Sources

The Proposed Action includes non-impulsive acoustic sources that require quantitative analysis. Some of the acoustic sources are either above the known hearing range of marine species or have narrow beam widths and short pulse lengths that would not result in effects to marine species, and are not quantitatively analyzed. Potential effects from these “*de minimis*” sources are analyzed qualitatively in accordance with current Navy policy.

Active acoustic sources would be lowered from a cruise vessel while stationary, deployed on gliders and UUVs, or deployed on fixed moorings. The total amount of active source testing for ship-deployed sources used over the duration of a cruise would be 120 hours. The testing would take place in the vicinity of the source locations in Figure 1-1, with UUVs running tracks within the designated box. During this testing, 35 Hz, 900 Hz and 10 kHz sources would be employed. Up to seven fixed acoustic navigation sources, transmitting at 900 Hz, would remain in place for a year at the locations given in Figure 1-1. Up to two VLF sources, transmitting at 35 Hz, would be deployed in a similar manner (Alternative 2 only). These sources would be deployed in two of the three VLF source positions in Figure 1-1. Two drifting IGBs would also be configured with active acoustic sources at 900 Hz and 10 kHz.

In assessing the potential for environmental harm to biological resources from non-impulsive acoustic sources, a variety of factors must be considered, including source characteristics, animal presence and associated density, duration of exposure, and thresholds for harm and harassment for the species that may occur in the Study Area. The severity of the potential consequences, such as physiological stress and behavioral response, depends on the received sound level at the animal, the details of the sound-producing activity, the animal’s life history stage (e.g., juvenile or adult, breeding or feeding season), and past experience with the stimuli. An animal’s life history stage is an important factor to consider when predicting whether a stress response is likely. An animal’s life history stage includes its level of physical maturity (i.e., larva, infant, juvenile, sexually mature adult) and the primary activity in which it is engaged such as mating, feeding, or rearing/caring for young. Prior experience with a stressor may be of particular importance because repeated experience with a stressor may dull the stress response via

acclimation (St Aubin and Dierauf 2001) or increase the response via sensitization. The types of potential consequences to marine species from acoustic sources can be described by the following categories:

Non-auditory injury: Non-auditory injury can occur to lungs and other organs and can cause tissue damage. Resonance occurs when the frequency of the sound waves matches the frequency of vibration of the air-filled organ or cavity, causing it to resonate. This can, in certain circumstances, lead to damage to the tissue making up the organ or air-filled cavity. Tissue damage also can be inflicted directly by sound waves in cases of sound waves with high amplitude and rapid rise time.

Hearing Loss: Also called a noise-induced threshold shift. Hearing loss manifests itself as loss in hearing sensitivity across part of an animal's hearing range, which is dependent upon the specifics of the noise exposure. Hearing loss may be either a permanent threshold shift (PTS) or a temporary threshold shift (TTS). If the threshold shift does not return to zero but leaves some finite amount of threshold shift, then that remaining threshold shift is a PTS. The intensity and duration of a sound that will cause PTS varies across species and even between individual animals. PTS is a consequence of the death of sensory hair cells of the auditory epithelia of the ear and a resultant loss of hearing ability in the general vicinity of the frequencies of stimulation (Myrberg 1990; Richardson et al. 1995b).

Physiological stress: Marine animals naturally experience physiological stress as part of their normal life histories. The physiological response to a stressor, often termed the stress response, is an adaptive process that helps an animal cope with changing external and internal environmental conditions. Sound-producing activities have the potential to cause additional stress. However, too much of a stress response can be harmful to an animal, resulting in physiological dysfunction.

If a sound is detected (i.e., heard or sensed) by an animal, a stress response can occur. Additionally, if an animal suffers injury or hearing loss, a physiological stress response will occur. The generalized stress response is characterized by a release of hormones (Reeder and Kramer 2005) and other chemicals (e.g., stress markers) such as reactive oxidative compounds associated with noise-induced hearing loss (Henderson et al. 2006). An acute stress response is traditionally considered part of the startle response and is hormonally characterized by the release of the catecholamines. Annoyance type reactions may be characterized by the release of either or both catecholamines and glucocorticoid hormones. Regardless of the physiological changes that make up the stress response, the stress response may contribute to an animal's decision to alter its behavior.

Behavioral response: Marine animals may exhibit short-term behavioral reactions, such as cessation of feeding, resting, or social interaction, and they also may exhibit alertness or avoidance behavior (Richardson et al. 1995b).

Masking: The presence of intense sounds or sounds within a mammal's hearing range in the environment potentially can interfere with an animal's ability to hear relevant sounds. This effect, known as "auditory masking," could interfere with the animal's ability to detect biologically relevant sounds, such as those produced by predators or prey, thus increasing the likelihood of the animal not finding food or being preyed upon (Myrberg 1981; Popper et al. 2004). Masking only occurs in the frequency band of the sound that causes the masking condition. Other relevant sounds with frequencies outside of this band would not be masked.

Non-impulsive acoustic sources are analyzed for their potential effects on invertebrates (Section 4.3.2.1.1), marine birds (Section 4.3.2.2.1), fish (Section 4.3.2.3.1), EFH (Section 4.3.2.4.1), and marine mammals (Section 4.3.2.5.1).

4.1.1.1.1 Alternative 1

Under Alternative 1, all acoustic sources listed in the Proposed Action (Section 2.2) would be used except for VLF sources.

4.1.1.1.2 Alternative 2 (Preferred Alternative)

Under Alternative 2 (Preferred Alternative), all acoustic sources listed in the Proposed Action (Section 2.2), including VLF sources, would be used.

4.1.1.2 Icebreaking Noise

CGC HEALY, the only potential icebreaking vessel associated with the Proposed Action, travels at a maximum speed of 3 knots when traveling through 3.5 ft (1.07 m) of sea ice (Murphy 2010). CGC HEALY may be required to perform icebreaking to deploy the moored acoustic sources. CGC HEALY has proven capable of breaking ice up to 8 ft (2.4 m) thick while backing and ramming (Roth et al. 2013). A study in the western Arctic Ocean was conducted while CGC HEALY was mapping the seafloor north of the Chukchi Cap in August 2008. During this study, CGC HEALY icebreaker events generated centered frequencies near 10, 50, and 100 Hz with maximum source levels of 190 to 200 dB re 1 μ Pa at 1 m (Roth et al. 2013).

The type of ice in the Study Area during the icebreaking would influence the type of organisms present and their reaction to icebreaking. Icebreaking would occur in the warm season, usually between July and October, each year through 2025, when ice thickness is expected to be at or near its lowest levels, which would minimize the time required for icebreaking. Icebreaking was modeled for seven days per year during this timeframe. In loose pack ice, the speed and noise of CGC HEALY is expected to be similar to those produced in the open ocean. In heavier pack ice or thick landfast ice, CGC HEALY would operate at a maximum speed of 3 knots, but power levels would be higher, which would increase the sound produced by CGC HEALY.

Marine species within the Study Area may be exposed to icebreaking noise associated with CGC HEALY during the Proposed Action. The potential harm from icebreaking noise is from masking other biologically important sounds or behavioral reactions such as alerting, avoidance, or other behavioral reactions.

The potential effects of icebreaking noise are analyzed for invertebrates (Section 4.3.2.1.2), marine birds (Section 4.3.2.2.2), fish (Section 4.3.2.3.2), EFH (Section 4.3.2.4.2), and marine mammals (Section 4.3.2.5.2).

4.1.1.2.1 Alternative 1

Icebreaking from both Alternatives 1 and 2 would result in the same potential for effects to biological resources, in that the same vessels would be utilized for either alternative.

4.1.1.2.2 Alternative 2 (Preferred Alternative)

Icebreaking from both Alternatives 1 and 2 would result in the same potential for effects to biological resources, in that the same vessels would be utilized for either alternative.

4.1.1.3 Vessel Noise

During the Proposed Action, vessel noise would be generated from either the R/V Sikuliaq or CGC HEALY. The R/V Sikuliaq has a one-third octave signature band range of 10 Hz to 200 kHz and a source level of 130 to 172 dB re 1 μ Pa at 1 m when traveling at 11 knots, and an one-third octave signature band range of 10 Hz to 200 kHz with a source level of 127 to 154 dB re 1 μ Pa at 1 m when traveling at 4

knots (Naval Sea Systems Command 2015). CGC HEALY travels at a maximum speed of 17 knots with a cruising speed of 12 knots (United States Coast Guard 2013), and a maximum speed of 3 knots when traveling through 1.07 m of sea ice (Murphy 2010). Icebreaking noise associated with CGC HEALY is discussed in Section 4.1.1.2.

Marine species within the Study Area may be exposed to vessel noise associated with the R/V Sikuliaq or the CGC HEALY during the Proposed Action. Vessel noise would result from open-ocean movement. The potential harm from vessel noise is from masking of other biologically relevant sounds as well as behavioral reactions, such as alerting, avoidance, or other behavioral reaction. Although unlikely due to the low volume of shipping traffic of the Arctic, some marine species may have habituated to vessel noise and may be more likely to respond to the sight of a vessel rather than the sound of a vessel, although both could play a role in prompting reactions (Hazel et al. 2007).

Auditory masking can occur due to vessel noise, potentially masking vocalizations and other biologically important sounds (e.g., sounds of prey or predators) on which marine organisms may rely. Potential masking can vary depending on the ambient noise level within the environment, the received level and frequency of the vessel noise, and the received level and frequency of the sound of biological interest. In the open ocean, ambient noise levels are between about 60 and 80 dB re 1 μ Pa, especially at lower frequencies (below 100 Hz) (National Research Council 2003). When the noise level is above the sound of interest, and in a similar frequency band, auditory masking could occur. Any sound that is above ambient noise levels and within an animal's hearing range needs to be considered in the analysis; however, noise that is just detectable over ambient levels is unlikely to actually cause any substantial masking.

Analysis of vessel noise associated with the Proposed Action has been completed for invertebrates (Section 4.3.2.1.3), marine birds (Section 4.3.2.2.3), fish (Section 4.3.2.3.3), and marine mammals (Section 4.3.2.5.3).

4.1.1.3.1 Alternative 1

Vessel noise would be the same in both Alternatives 1 and 2.

4.1.1.3.2 Alternative 2 (Preferred Alternative)

Vessel noise would be the same in both Alternatives 1 and 2.

4.1.2 Physical Stressors

Physical stressors resulting from the Proposed Action include icebreaking (physical impacts), vessel and in-water device strike, and bottom disturbance.

4.1.2.1 Icebreaking (Physical Impacts)

Icebreaking could occur in the Study Area when transiting out to deploy sources, at speeds of 3 to 6 knots. CGC HEALY would be icebreaking, as needed, during the warm season while the ice is at its lowest extent of the year.

Icebreaking has the potential to harm sea ice (Section 4.2.2.1), invertebrates (Section 4.3.2.1.4), fish (Section 4.3.2.3.4), EFH (Section 4.3.2.4.3), and marine mammals (Section 4.3.2.5.4) by causing behavioral reactions, mortality upon impact, and/or altering habitats.

4.1.2.1.1 Alternative 1

Icebreaking would occur in equal measure under both Alternatives 1 and 2.

4.1.2.1.2 Alternative 2 (Preferred Alternative)

Icebreaking would occur in equal measure under both Alternatives 1 and 2.

4.1.2.2 Vessel and In-Water Device Strike

The vessels that would be utilized during the Proposed Action are the R/V Sikuliaq (maximum speed of 12 knots), and CGC HEALY (maximum speed of 17 knots). These vessels would not be operating at their maximum speed due to travel through the marginal ice zone. Gliders also have the potential to result in strike to marine resources, though this would be very unlikely. Gliders are slow moving, travelling at a speed under 1 knot. Physical disturbance from the use of in-water devices would not be expected to result in more than a momentary behavioral response. Any change to an individual animal's behavior from in-water devices would not be expected to result in long-term or population-level effects. Research on marine animal's responses to gliders has not been conducted; the discussion below is based on potential reactions to vessels, which is used as a surrogate for this analysis.

Vessels have the potential to affect invertebrates, fish, or marine mammals by eliciting a behavioral response or causing mortality or serious injury from collisions. It is difficult to differentiate between behavioral responses to vessel sound and visual cues associated with the presence of a vessel; thus, it is assumed that both play a role in prompting reactions from animals. Reactions to vessels often include changes in general activity (e.g., from resting or feeding to active avoidance), changes in surfacing-respiration-dive cycles, and changes in speed and direction of movement. Past experiences of the animals with vessels are important in determining the degree and type of response elicited from an animal-vessel encounter. Some species have been noted to tolerate slow-moving vessels within several hundred meters, especially when the vessel is not directed toward the animal and when there are no sudden changes in direction or engine speed (Heide-Jørgensen et al. 2003; Richardson et al. 1995b).

Vessel and in-water device strike would not affect bottom substrates, as none of the vehicles would be at bottom depth, nor would they affect EFH or marine birds. The potential effects on invertebrates (Section 4.3.2.1.5), fish (Section 4.3.2.3.5), and marine mammals (Section 4.3.2.5.5) have been analyzed.

4.1.2.2.1 Alternative 1

Vessel and in-water device use would be the same under both Alternatives 1 and 2.

4.1.2.2.2 Alternative 2 (Preferred Alternative)

Vessel and in-water device use would be the same under both Alternatives 1 and 2.

4.1.2.3 Bottom Disturbance

Various components of the Proposed Action would have the potential to alter the bottom substrate. These would include expenditure of anchors and other materials that would sink to the bottom.

During activities in the Study Area, various items would be introduced and expended into the marine environment, including onto the soft bottom seafloor (Section 3.1.1.2). These expended materials have the potential to strike a resource once they sink to the seafloor and settle in the bottom substrate. Expended materials that would be expected to sink to the seafloor include expended buoys and other anchors or tethers. The Proposed Action would utilize various anchored and tethered equipment. These anchors would be bottom placed and could weigh up to 800 lbs (363 kg).

Bottom disturbance is not expected to affect marine birds, EFH, or marine mammals as they do not inhabit the seafloor within the Study Area. Therefore, these resources are not further analyzed. The

potential effects on the physical environment (Section 4.2.2.2), invertebrates (Section 4.3.2.1.6), and fish (Section 4.3.2.3.6) have been analyzed.

4.1.2.3.1 Alternative 1

Bottom disturbance would be the same under both Alternatives 1 and 2.

4.1.2.3.2 Alternative 2 (Preferred Alternative)

Bottom disturbance would be the same under both Alternatives 1 and 2.

4.1.3 Expended Materials

4.1.3.1 Entanglement

Devices that pose an entanglement risk are those with lines or tethers; devices with a potential for entanglement include moored or ice-tethered sensors. All lines hanging from buoys or ice tethered equipment would be weighted, and therefore would not have any loops or slack.

The final line that could be a threat for entanglement is the use of a device tethered to an unmanned underwater vehicle (depth of 295 ft [91 m]). The tether for this research initiative has a diameter of 0.35 in (0.9 cm), and is made of Kevlar. This tether has a very high breaking strength (1,543 lb force [700 kg force]), but environmental resources should not be at risk due to the small likelihood of any loops or slack developing in this line, since it would be under positive pressure. No mooring lines would be expended during the proposed action, so this further limits the chance for entanglement.

It is not anticipated that entanglement would affect marine habitats, marine birds, or EFH, as they are not within an area to be adversely affected or cannot become entangled in expended material. Therefore, they will not be further analyzed. The potential effects on invertebrates (Section 4.3.2.1.7), fish (Section 4.3.2.3.7), and marine mammals (Section 4.3.2.5.6) have been analyzed.

4.1.3.1.1 Alternative 1

Under Alternative 1, the potential for entanglement would be from mooring lines. Alternative 1 has fewer mooring lines associated with the Proposed Action due to the exclusion of the VLF sources. In the upper portion of the water column object deployment would be controlled.

4.1.3.1.2 Alternative 2 (Preferred Alternative)

Under Alternative 2, the potential for entanglement would be from mooring lines. Lines extending from the moorings would be retrieved at the completion of the Proposed Action. In the upper portion of the water column object deployment would be controlled.

4.1.3.2 Ingestion

During the Proposed Action, the only expended materials available for ingestion include the on-ice measurement systems. On-ice measurement systems include the autonomous weather station and the ice mass balance buoy. The autonomous weather station would be deployed on a tripod with insulated foot platforms frozen into the ice. While the ice mass balance buoy would be lowered into the water column through a two-inch hole in the ice, there would be a tripod located on the ice. All other expended objects would be expended into the water column and would sink to the seafloor. Ingestion of these materials does not require the entire object to be ingested; pieces of objects that either break off or are bitten off are included in this analysis.

Ingestion stressors are not anticipated to affect any resources other than marine mammals, specifically polar bears, due to the large size of the material that is expended in the water column or stationed on the sea ice. These objects (e.g., anchors, buoys) would be too large for any other marine resource to eat. Additionally, within the Study Area marine mammals would not be feeding near the seafloor further eliminating any overlap of expended materials and marine mammals. Therefore, harm to marine resources (other than polar bears) is not discussed further. The objects deployed and expended within the water column would be too deep to overlap with a swimming polar bear. The potential effects from ingestion of expended materials have been analyzed for polar bears (Section 4.3.2.5.7).

4.1.3.2.1 Alternative 1

Potential risk for ingestion would be the same under both Alternatives 1 and 2.

4.1.3.2.2 Alternative 2 (Preferred Alternative)

Potential risk for ingestion would be the same under both Alternatives 1 and 2.

4.2 Physical Resources

4.2.1 No Action Alternative

Under the No Action Alternative, the Proposed Action would be limited to equipment retrieval pursuant to the existing supplemented 2018 OEA and regulatory authorizations (U.S. Department of the Navy 2018a, 2021b); any equipment that cannot be retrieved will go silent and be abandoned. There would be no change to the baseline physical environment. Therefore, no significant harm to the physical environment would occur with implementation of the No Action Alternative. However, the No Action Alternative would not meet the purpose and need of the Proposed Action.

4.2.2 Action Alternatives

4.2.2.1 Icebreaking (Physical Impacts)

Potential Harm

Sea ice is considered important habitat for many polar species including diatoms, Arctic cod, ringed seals, and polar bears. Many species feed along the ice edge, while others use it for resting, pupping, or traveling.

Ice, however thin, does not fracture by itself, but rather, wind, pressure systems, and ocean gyres transport ice and often cause fractures to form. Cracks are a regular feature of ice. During winter when fractures appear, leads form but quickly freeze over again. From May onwards, with the sun shining down on the Arctic, the thin ice will disappear, leaving behind stretches of open water, sometimes well within the ice pack. The total sea ice extent was 4.14 million mi² (10.71 million km²) in the Arctic in June of 2021, the month when the sun's energy is strongest (NOAA National Centers for Environmental Information 2021). An icebreaker cruising through the ice for 620 mi (1,000 km) and leaving an ice-free wake of 33 ft (10 m) would open an area of water 3.9 mi² (10 km²) over the entire cruise. In contrast, the Arctic sea ice cover decreases by an average of over 3.5 million mi² (9 million km²) each year during the melt season (National Snow and Ice Data Center 2012). This is an area larger than the contiguous United States. In total, researchers estimate that the number of icebreakers traversing the Arctic at any given time is usually less than three. Thus, the actual contribution of icebreaking to sea ice reduction is miniscule—only one part in a million of the total ice cover (National Snow and Ice Data Center 2012). As

the ice pack has started to break up in ever smaller parts earlier in the year, it has also become easier for vessels to move the ice around.

Alternative 1

CGC HEALY does not diminish or destroy ice habitat, and the amount of ice that is broken up relative to the overall total amount of ice is small. In accordance with E.O. 12114 physical impacts from icebreaking associated with Alternative 1 would not significantly harm the physical environment (sea ice habitat).

Alternative 2 (Preferred Alternative)

CGC HEALY does not diminish or destroy ice habitat, and the amount of ice that is broken up relative to the overall total amount of ice is small. In accordance with E.O. 12114 physical impacts from icebreaking associated with Alternative 2 would not significantly harm the physical environment (sea ice habitat).

4.2.2.2 Bottom Disturbance

Potential Harm

In general, three things happen to expended materials that come to rest on the ocean floor: (1) they lodge in sediment where there is little or no oxygen, usually below 4 in (10 cm), (2) they remain on the ocean floor and begin to react with seawater, or (3) they remain on the ocean floor and become encrusted by marine organisms. As a result, rates of deterioration depend on the material and the conditions in the immediate marine and benthic environment. If buried deep in ocean sediments, materials tend to decompose at much lower rates than when exposed to seawater (Ankley et al. 1996). In those situations where metals are exposed to seawater, they begin to slowly corrode, a process that creates a layer of corroded material between the seawater and uncorroded metal. This layer of corrosion removes the metal from direct exposure to the corrosiveness of seawater, a process that further slows movement of the metals into the adjacent sediment and water column. Any elevated levels of metals in sediment would be restricted to a small zone around the metal, and any release to the overlying water column would be diluted. In a similar fashion, as materials become covered by marine life, the direct exposure of the material to seawater decreases and the rate of corrosion decreases (Little and Ray 2002). Dispersal of these materials in the water column is controlled by physical mixing and diffusion, both of which tend to vary with time and location. The disturbance of bottom sediments by objects settling onto the seafloor could result in temporary and localized increases in turbidity that would quickly dissipate.

Large-scale processes control sediment composition in the deep sea, so the seafloor tends to be uniform over hundreds of square miles. At the spatial scale at which most individual organisms experience their environment (millimeters to meters), the seafloor is typically heterogeneous (Thistle 2003). The instances of bottom disturbance during the Proposed Action would be minimal, due to the few items expended over the large region of the Study Area.

Alternative 1

Based on the geographically expansive size of the Study Area in comparison to the small area of the anchors, the physical marine environment would not be altered in any meaningful way. In accordance with E.O. 12114, bottom disturbance associated with Alternative 1 would not result in significant harm to the physical environment (bottom substrate).

Alternative 2 (Preferred Alternative)

Based on the geographically expansive size of the Study Area in comparison to the small area of the anchors, the physical marine environment would not be altered in any meaningful way. In accordance

with E.O. 12114, bottom disturbance associated with Alternative 2 would not result in significant harm to the physical environment (bottom substrate).

4.3 Biological Resources

4.3.1 No Action Alternative

Under the No Action Alternative, the Proposed Action would be limited to equipment retrieval pursuant to the existing supplemented 2018 OEA and regulatory authorizations (U.S. Department of the Navy 2018a, 2021b); any equipment that cannot be retrieved will go silent and be abandoned. Any abandoned equipment would eventually sink to the seafloor; further details can be found in the 2021 Supplemental OEA (U.S. Department of the Navy 2021b). There would be no change to biological resources. Therefore, no significant harm to biological resources would occur with implementation of the No Action Alternative. However, the No Action Alternative would not meet the purpose and need of the Proposed Action.

4.3.2 Action Alternatives

4.3.2.1 Invertebrates

Excluding microbes, approximately 5,000 known marine invertebrates have been documented in the Arctic; the number of species is likely higher, though, since this area is not well studied (Josefson et al. 2013). Although most species are found within the benthic zone, marine invertebrates can be found in all zones (sympagic [within the sea ice], pelagic [open ocean], or benthic [bottom dwelling]) of the Beaufort Sea (Josefson et al. 2013). Sea ice provides a habitat for algae and a nursery ground for invertebrates during times when the water column does not support phytoplankton growth (Winfrey 2005). Sympagic zone invertebrates live within the pores and brine channels of the ice (small spaces within the sea ice which are filled with a salty solution called brine) or at the ice-water interface. Biodiversity of species is low within the sympagic zone due to the extreme conditions of the sea ice (Leet et al. 2001). Within the Study Area, many sympagic species also exist in and along the edges of ice coverage, feeding on blooms of phytoplankton and other algae which grow in, on, or adjacent to the ice (Wyllie-Echeverria and Ackerman 2003). Marine invertebrate distribution in the Beaufort Sea is influenced by habitat and oceanographic conditions (e.g., depth, temperature, salinity, nutrient concentrations, and ocean currents) (Levinton 2009). No ESA-listed invertebrate species are present in the Study Area.

Acoustic stressors that may have potential impacts on invertebrates include non-impulsive acoustic sources, icebreaking noise, and vessel noise. Physical stressors that may have potential impacts on invertebrates include icebreaking (physical impacts), vessel and in-water device strike, and bottom disturbance. The only stressor associated with expended materials that may have potential impacts on invertebrates is entanglement.

4.3.2.1.1 Non-Impulsive Acoustic Sources

Potential Impacts

Hearing capabilities of invertebrates are largely unknown, although they are not expected to hear sources above 3 kHz (see Section 3.2.2.1.5 for invertebrate hearing information). Invertebrates are only expected to potentially perceive the signals of a few sources used during the Proposed Action. In addition, most marine invertebrates in water are known to detect only particle motion associated with

sound waves (Graduate School of Oceanography 2021), which drop off rapidly with distance, limiting exposure to the period when an invertebrate is close to a sound source.

Outside of studies conducted to test the sensitivity of invertebrates to vibrations, very little is known on the effects of anthropogenic underwater noise on invertebrates (Edmonds et al. 2016). While data are limited, research suggests that some of the major cephalopods and decapods may have limited hearing capabilities (Hanlon 1987; Offutt 1970), and may hear only low-frequency (less than 1 kHz) sources (Offutt 1970), which is most likely within the frequency band of biological signals (Hill 2009). In a review of crustacean sensitivity of high amplitude underwater noise by Edmonds et al. (2016), crustaceans may be able to hear the frequencies at which they produce sound, but it remains unclear which noises are incidentally produced and if there are any negative effects from masking them. Acoustic signals produced by crustaceans range from low frequency rumbles (20-60 Hz) to high frequency signals (20-55 kHz) (Henninger and Watson 2005; Patek and Caldwell 2006; Staaterman et al. 2016). Aquatic invertebrates that can sense local water movements with ciliated cells include cnidarians, flatworms, segmented worms, urochordates (tunicates), mollusks, and arthropods (Budelmann 1992a, 1992b; Popper et al. 2001). Some aquatic invertebrates have specialized organs called statocysts for determination of equilibrium and, in some cases, linear or angular acceleration. Statocysts allow an animal to sense movement and may enable some species, such as cephalopods and crustaceans, to be sensitive to water particle movements associated with sound (Goodall et al. 1990; Hu et al. 2009; Kaifu et al. 2008; Montgomery et al. 2006; Popper et al. 2001; Roberts and Breithaupt 2016; Salmon 1971). Because any acoustic sensory capabilities, if present at all, are limited to detecting water motion, and water particle motion near a sound source falls off rapidly with distance, aquatic invertebrates are probably limited to detecting nearby sound sources rather than sound caused by pressure waves from distant sources.

Studies of sound energy effects on invertebrates are few, and identify only behavioral and physiological responses. Non-auditory injury, PTS, TTS, and masking studies have not been conducted for invertebrates. Both behavioral and auditory brainstem response studies suggest that crustaceans may sense frequencies up to 3 kHz, but best sensitivity is likely below 200 Hz (Goodall et al. 1990; Lovell et al. 2005; Lovell et al. 2006). Most cephalopods likely sense low-frequency sound below 1 kHz, with best sensitivities at lower frequencies (Budelmann 2010; Mooney et al. 2010; Offutt 1970). A few cephalopods may sense higher frequencies up to 1,500 Hz (Hu et al. 2009). Hudson et al. (2022) recently examined the effects of low- (<1 kHz) and mid-frequency (1.6 to 4.0 kHz) sounds on two crustacean species (blue crabs [*Callinectes sapidus*] and American lobsters [*Homarus americanus*]). They observed that physiological indicators of stress returned to baseline levels within 48 hours of exposure with the exception of an amplified hemolymph glucose signal for seven days after exposure to mid-frequency sonar.

Within the Study Area, marine invertebrate abundance is low within the sea ice and in the water column. The highest densities are on the seafloor, further reducing the likelihood of invertebrates hearing the frequencies of the active acoustic sources due to the dissipation of the non-impulsive acoustic sources in the water column. In studies by Christian et al. (2003) and Payne et al. (2007), neither found damage to lobster or crab statocysts from high intensity air gun firings (which is of greater intensity than the non-impulsive acoustic sources in the Proposed Action). Furthermore, in the study by Christian et al. (2003), no changes were found in biochemical stress markers in snow crabs.

Alternative 1

There is a low likelihood that invertebrates would be able to perceive the non-impulsive acoustic sources, and if perceived, that an individual animal would react. Therefore, in accordance with E.O.

12114, non-impulsive acoustic sources associated with Alternative 1 would not result in significant harm to invertebrates.

Alternative 2 (Preferred Alternative)

There is a low likelihood that invertebrates would be able to perceive the non-impulsive acoustic sources, and if perceived, that an individual animal would react. The inclusion of VLF sources would not be likely to increase the potential for impacts from non-impulsive acoustic sources. Therefore, in accordance with E.O. 12114, non-impulsive acoustic sources associated with Alternative 2 would not result in significant harm to invertebrates.

4.3.2.1.2 Icebreaking Noise

Potential Impacts

Icebreaking noise is generally low frequency impulsive sound similar in frequency to vessel noise, with the impulsive nature being the primary difference. As such, the species expected to respond and the levels of response to icebreaking noise would be expected to be similar for icebreaking and vessel noise. As addressed in Section 3.2.2.1.5, hearing capabilities of invertebrates are largely unknown, although they are not expected to hear sources above 3 kHz (Lovell et al. 2005; Popper 2008). Impacts to invertebrates from icebreaking noise is relatively unknown, but it is likely that some species including crustaceans and cephalopods would be able to perceive the low frequency sources generated from icebreaking that occurs during the Proposed Action, which could result in masking acoustic communication in invertebrates such as crustaceans (Staaterman et al. 2011). Avoidance behavior, short term temporary responses (such as feeding cessation, increased stress, or other minor physiological harm) may occur if invertebrates were close enough to the icebreaking (Edmonds et al. 2016; Roberts and Breithaupt 2016). Masking of important acoustic cues used by invertebrates during larval orientation and settlement may lead to maladaptive behavior that could reduce successful recruitment (Simpson et al. 2011).

Icebreaking associated with the Proposed Action would be short-term and temporary as the vessel moves through an area, and it is not anticipated that this short-term noise would result in significant harm via masking; nor is it expected to result in more than a temporary behavioral reaction of marine invertebrates in the vicinity of the icebreaking event. It is expected that invertebrates would return to their normal behavior shortly after exposure.

Alternative 1

Icebreaking noise, if perceived by an invertebrate, would likely result in temporary behavioral reactions and would not result in any population level impacts. In accordance with E.O. 12114, icebreaking noise associated with Alternative 1 would not result in significant harm to invertebrates.

Alternative 2 (Preferred Alternative)

Icebreaking noise, if perceived by an invertebrate, would likely result in temporary behavioral reactions and would not result in any population level impacts. In accordance with E.O. 12114, icebreaking noise associated with Alternative 2 would not result in significant harm to invertebrates.

4.3.2.1.3 Vessel Noise

Potential Impacts

As addressed in Section 3.2.2.1.5, hearing capabilities of invertebrates are largely unknown, although they are not expected to hear sources above 3 kHz (Lovell et al. 2005; Popper 2008). Impacts to invertebrates from vessel noise is relatively unknown, but it is likely that some species would be able to

perceive the low frequency sounds generated from the vessels (see Section 2.2.2) used during the Proposed Action, which could result in masking acoustic communication in invertebrates such as crustaceans (Staaterman et al. 2011). Masking of important acoustic cues used by invertebrates during larval orientation and settlement may lead to maladaptive behavior that could reduce successful recruitment (Simpson et al. 2011). Recent research suggests that some invertebrates may experience sub-lethal physiological impacts from prolonged exposure to high amplitude, low frequency sound (Celi et al. 2014; Wale et al. 2013); however, the Study Area is over deep water, which would limit the exposure of benthic invertebrates. Since vessels are generally transiting through, prolonged exposure to high amplitudes such as those used in these studies is unlikely. The low-frequency component of vessel noise would likely be detected by some invertebrates, although the number of individuals affected would be limited to those near enough to a source to experience particle motion.

Several studies have found physiological and behavioral responses in some invertebrate species in response to playback of vessel noise, although one study found no reaction by krill to an approaching vessel. Physiological effects included biochemical changes indicative of stress in crustacean species, decreased growth and reproduction in shrimp, and changes in sea hare embryo development. Nedelec et al. (2014a) exposed sea hares to vessel noise playback for 45 seconds every five minutes over a 12-hour period and found reduced embryo development and increased larvae mortality. Hudson et al. (2022) observed no significant biochemical changes following exposure to low-frequency simulated vessel noise. It is also possible that vessel noise may contribute to masking of relevant environmental sounds, such as predator detection.

Behavioral effects resulting from vessel noise playback have been observed in various crustacean, cephalopod, and bivalve species and include shell closing and changes in feeding, coloration, swimming, and other movements. Hudson et al. (Hudson et al. 2022) did not observe activity level changes in American lobsters or blue crabs exposed to simulated vessel noise; however, they did observe behavioral changes to blue crabs' competitive behaviors for 24 hours following exposure to simulated vessel noise when also exposed to a competitor, the green crab (*Callinectes maenas*). In addition to disruption of important processes like feeding or seeking shelter, behavioral reactions can result in increased energy expenditure (Hudson et al. 2022). However, because a vessel would only be within a given area for a brief period of time, behavioral reactions would be short-term.

Vessel noise associated with the Proposed Action would be short-term and temporary as the vessel moves through an area, and it is not anticipated that this short-term noise would result in significant harm via masking; nor is it expected to result in more than a temporary behavioral reaction of marine invertebrates in the vicinity of the vessel noise. It is expected that invertebrates would return to their normal behavior shortly after exposure.

Alternative 1

Vessel noise from both Alternatives 1 and 2 would result in the same potential for effects to invertebrates, in that the same vessels would be utilized for both Alternatives. Vessel noise, if perceived by an invertebrate, would likely result in temporary behavioral reactions and would not result in any population level impacts. In accordance with E.O. 12114, vessel noise associated with Alternative 1 would not result in significant harm to invertebrates.

Alternative 2 (Preferred Alternative)

Vessel noise from both Alternatives 1 and 2 would result in the same potential for effects to invertebrates, in that the same vessels would be utilized for both Alternatives. Vessel noise, if perceived by an invertebrate, would likely result in temporary behavioral reactions and would not result in any

population level impacts. In accordance with E.O. 12114, vessel noise associated with Alternative 2 would not result in significant harm to invertebrates.

4.3.2.1.4 Icebreaking (Physical Impacts)

Potential Impacts

The population of invertebrates with the most potential for harm from icebreaking associated with the Proposed Action are the sympagic invertebrates that live on or in the sea ice (Guglielmo et al. 2000; Kohlbach et al. 2016; Kramer et al. 2011). Individuals of these species could be killed or displaced by the icebreaking. Because the impact would be localized to the immediate path of the vessel, icebreaking disturbance would not be expected to harm the vast majority of the biomass of sympagic invertebrates and therefore, no population level impacts would be expected. Though many other communities are also dependent on sympagic production (Kohlbach et al. 2016), the impact on those food web dynamics would be similarly small, since the ratio of affected area to unaffected area is extremely small.

Alternative 1

Icebreaking from both Alternatives 1 and 2 would result in the same potential for effects to invertebrates, in that the same vessels would be utilized for both alternatives. Although some invertebrates could be disturbed or killed by icebreaking, population level effects are not anticipated. In accordance with E.O. 12114, physical impacts from icebreaking associated with Alternative 1 would not result in significant harm to invertebrates.

Alternative 2 (Preferred Alternative)

Icebreaking from both Alternatives 1 and 2 would result in the same potential for effects to invertebrates, in that the same vessels would be utilized for both alternatives. Although some invertebrates could be disturbed or killed by icebreaking, population level effects are not anticipated. In accordance with E.O. 12114, physical impacts from icebreaking associated with Alternative 2 would not result in significant harm to invertebrates.

4.3.2.1.5 Vessel and In-Water Device Strike

Potential Impacts

Vessels and in-water devices have the potential to harm marine invertebrates by disturbing the water column or directly striking organisms (Bishop 2008). Vessel movement may result in short-term and localized disturbances to invertebrates, such as zooplankton and cephalopods, utilizing the upper water column. Propeller wash (water displaced by propellers used for propulsion) from vessel, and vehicle movement can potentially disturb marine invertebrates in the water column and are a likely cause of zooplankton mortality (Bickel et al. 2011). Since most of the macroinvertebrates within the Study Area are benthic and the Proposed Action takes place within the water column, potential for macroinvertebrate vessel or vehicle strike is extremely low. No measurable effects on invertebrate populations in the water column would occur because the number of organisms exposed to vessel movement would be low relative to total invertebrate biomass.

Alternative 1

Vessel and in-water device strike from both Alternatives 1 and 2 would result in the same potential for effects to invertebrates, in that the same vessels and in-water devices would be utilized for both Alternatives. Although some invertebrates could be disturbed or killed by vessel and in-water device strike, population level effects are not anticipated. In accordance with E.O. 12114, vessel and in-water device strike associated with Alternative 1 would not result in significant harm to invertebrates.

Alternative 2 (Preferred Alternative)

Vessel and in-water device strike from both Alternatives 1 and 2 would result in the same potential for effects to invertebrates, in that the same vessels and in-water devices would be utilized for both Alternatives. Although some invertebrates could be disturbed or killed by vessel and in-water device strike, population level effects are not anticipated. In accordance with E.O. 12114, vessel and in-water device strike associated with Alternative 2 would not result in significant harm to invertebrates.

4.3.2.1.6 Bottom Disturbance

Potential Impacts

Effects to invertebrates from bottom disturbance would be either from the temporary and localized disturbance of the sediment or the bottom habitat changing from a soft bottom habitat to hard bottom due to the expended material. Expended material that would eventually sink may cause disturbance, injury, or mortality within the footprint of the device, may disturb marine invertebrates outside the footprint of the device, and would cause temporary local increases in turbidity near the seafloor. The overall footprint of the expended materials is minor compared to the size of the Study Area. The sediment disturbance would be temporary causing increased turbidity in the water locally. Objects that sink to the seafloor or are moored to the seafloor may attract invertebrates, or provide temporary attachment points for invertebrates. Some invertebrates attached to the devices would be removed from the habitat when the objects are recovered. In the immediate area where the expended material settled the bottom type would change from soft to hard substrate and may displace any invertebrates requiring soft bottom habitat. This may also attract invertebrates that attach to hard bottom substrate. The impact of expended materials on invertebrates may cause injury or mortality to individuals, but impacts to populations would be inconsequential due to the short-term disturbance during installation and removal of these devices.

Alternative 1

Invertebrates may be displaced, temporarily disturbed, or killed due to bottom disturbance as items are deployed or sink to the seafloor, but no population level effects are expected to occur. Under Alternative 1, the disturbance associated with the Proposed Action would be localized and temporary. In accordance with E.O. 12114, bottom disturbance associated with Alternative 1 would not result in significant harm to invertebrates.

Alternative 2 (Preferred Alternative)

Invertebrates may be displaced, temporarily disturbed, or killed due to bottom disturbance as items are deployed or sink to the seafloor, but no population level effects are expected to occur. Under Alternative 2, the disturbance associated with the Proposed Action would be localized and temporary. In accordance with E.O. 12114, bottom disturbance associated with Alternative 2 would not result in significant harm to invertebrates.

4.3.2.1.7 Entanglement

Potential Impacts

A marine invertebrate that might become entangled may only be temporarily confused and escape unharmed, it could be held tightly enough that it could be injured during its struggle to escape, it could be preyed upon while entangled, or it could starve while entangled. The likelihood of these outcomes cannot be predicted with any certainty because interactions between invertebrate species and entanglement hazards are not well known. Potential entanglement scenarios are based on observations of how marine invertebrates are entangled in marine debris that typically floats at the sea surface for

long periods of time (e.g., plastic bags and food wrappers), which is far more prone to tangling than weighted sensors dangling from buoys or floats, because these devices would not have materials prone to developing loops (Environmental Sciences Group 2005; Ocean Conservancy 2010). Deployments of the moorings and floats could cause short-term and localized disturbances to invertebrates utilizing the upper water column. Since most of the invertebrates within the Study Area are benthic, the risk of entanglement from deployment of moorings is extremely low.

Invertebrates also have an entanglement risk from the expended materials as they sink and land on the seafloor. Since all devices are lowered from a winch system in a controlled manner, the risk of entanglement from deployment of moorings is extremely low. Unlike marine mammals and fish, some invertebrates are sessile and would not be able to move out of the path of an expended material as it sinks and settles on the seafloor. Although there is a risk of an expended material entangling around and potentially injuring or killing an individual invertebrate, there would be no long term population level effects due to the small amount of expended materials over the large Study Area and the limited number of organisms potentially exposed to the material.

Alternative 1

Under Alternative 1, the potential for entanglement would be from mooring lines. Alternative 1 has fewer mooring lines associated with the Proposed Action due to the exclusion of the VLF sources. Lines extending from the moorings would be retrieved at the completion of the Proposed Action. In the upper portion of the water column object deployment would be controlled, which would greatly limit entanglement with invertebrates found in the sympagic or pelagic zones. Invertebrates within the benthic zone may be displaced, temporarily disturbed, or killed, but no population level effects are expected to occur. Therefore, in accordance with E.O. 12114, entanglement associated with Alternative 1 would not result in significant harm to invertebrates.

Alternative 2 (Preferred Alternative)

Under Alternative 2, the potential for entanglement would be from mooring lines. Alternative 2 has more mooring lines associated with the Proposed Action due to the inclusion of the VLF sources. Lines extending from the moorings would be retrieved at the completion of the Proposed Action. In the upper portion of the water column object deployment would be controlled, which would greatly limit entanglement with invertebrates found in the sympagic or pelagic zones. Invertebrates within the benthic zone may be displaced, temporarily disturbed, or killed, but no population level effects are expected to occur. Therefore, in accordance with E.O. 12114, entanglement associated with Alternative 2 would not result in significant harm to invertebrates.

4.3.2.2 Marine Birds

A combination of short-distance migrants, long-distance migrants, and year-round resident marine bird species occur within the Study Area (Section 3.2.2.2). Of all the marine birds that occur in the vicinity of the Study Area, only the thick-billed murre exhibits foraging diving behaviors at distances greater than 90 nm (167 km) from the shoreline during the timeframe of the Proposed Action. However, the thick-billed murre is not expected to forage in the deep waters of the Study Area. Therefore, no birds are expected to be foraging or migrating through the deep water Study Area. All other marine bird species in the area would either not travel over 90 nm (167 km) offshore or are not expected to forage underwater within the Study Area. No ESA-listed birds would be present in the Study Area during the Proposed Action.

Acoustic stressors that may have potential impacts on birds include non-impulsive acoustic sources, icebreaking noise, and vessel noise. Physical stressors would not impact birds and are not discussed further.

4.3.2.2.1 Non-Impulsive Acoustic Sources

Potential Impacts

Information regarding the impacts of sonar on birds is unavailable. Little is known about the ability for birds to hear underwater, although researchers have recently begun to examine this topic (Section 3.2.2.2.1). The limited information indicates that diving birds have a more narrow hearing range in water than in air (Dooling and Therrien 2012; Johansen et al. 2016). Birds have been reported to hear best at mid-frequencies (1 to 5 kHz), and are likely to be able to hear the low- and mid-frequency signals associated with the Proposed Action. No birds are expected to forage in the deep waters of the Study Area.

Alternative 1

Under Alternative 1, marine birds are not expected to encounter non-impulsive acoustic sources within the Study Area. The potential for a marine bird to be underwater and within receiving distance of an acoustic source is unlikely due to short duration of their dives, the ice cover in the Study Area, and the spread nature of the acoustic sources. Therefore, pursuant to E.O. 12114, non-impulsive acoustic sources associated with the Proposed Action would not result in significant harm to marine birds. Pursuant to the MBTA, non-impulsive acoustic sources associated with the Proposed Action would have no effect on migratory bird populations.

Alternative 2 (Preferred Alternative)

Under Alternative 2, marine birds are not expected to encounter non-impulsive acoustic sources within the Study Area. The potential for a marine bird to be underwater and within receiving distance of an acoustic source is unlikely due to short duration of their dives, the ice cover in the Study Area, and the spread nature of the acoustic sources. Therefore, pursuant to E.O. 12114, non-impulsive acoustic sources associated with the Proposed Action would not result in significant harm to marine birds. Pursuant to the MBTA, non-impulsive acoustic sources associated with the Proposed Action would have no effect on migratory bird populations.

4.3.2.2.2 Icebreaking Noise

Potential Impacts

Auditory masking related to marine bird hearing is unlikely, as marine birds spend a limited amount of time underwater and it is not thought that they use underwater sound related to their biologically relevant sounds. Marine birds would not be diving in ice-covered areas; therefore, they would not be underwater directly near the icebreaking vessel. However, noise propagating from the location of icebreaking could elicit short-term behavioral (startle response, swimming away, looking up) or physiological responses (increased heart rate), but are not likely to disrupt major behavior patterns, such as migrating, breeding, feeding, and sheltering, or to result in serious injury to any seabirds. Icebreaking noise is generally described as a low frequency, 10 to 100 Hz (Roth et al. 2013), non-impulsive sound. While Godin (2006) states that the air-water interface is nearly transparent when it comes to the transmission of low-frequency sound, this is not within the range of best hearing for birds in air. In addition, any noise associated with icebreaking, both in-air and underwater, would likely fall within the spectrum of natural ice-related sounds expected in the Arctic environment. Thus, icebreaking noise is unlikely to be detected by seabirds, either in air or if the sound transmission carries underwater.

Icebreaking noise from the Proposed Action is not expected to be received at levels that would elicit a response.

Alternative 1

Icebreaking noise from both Alternatives 1 and 2 would result in the same potential for effects to marine birds because the same vessels would be utilized for both alternatives. Due to the insignificant and short-term reactions to icebreaking noise, and in accordance with E.O. 12114, icebreaking noise associated with Alternative 1 would not result in significant harm to marine birds. Icebreaking noise associated with Alternative 1 would not result in a significant adverse effect on populations of marine birds protected under the MBTA.

Alternative 2 (Preferred Alternative)

Icebreaking noise from both Alternatives 1 and 2 would result in the same potential for effects to marine birds because the same vessels would be utilized for both alternatives. Due to the insignificant and short-term reactions to icebreaking noise, and in accordance with E.O. 12114, icebreaking noise associated with Alternative 2 would not result in significant harm to marine birds. Icebreaking noise associated with Alternative 2 would not result in a significant adverse effect on populations of marine birds protected under the MBTA.

4.3.2.2.3 Vessel Noise

Potential Impacts

Auditory masking related to marine bird hearing would not have an impact on marine birds, as they spend a limited amount of time underwater and it is not thought that they use underwater sound related to their biologically relevant sounds. However, vessel noise could elicit short-term behavioral (startle response, flying away, looking up) or physiological responses (increased heart rate), but is not likely to disrupt major behavior patterns, such as migrating, breeding, feeding, and sheltering, or to result in serious injury to any seabirds. In air, Beason (2004) notes that birds exposed to 146 dB re 20 μ Pa sound pressure level in air within 325 ft (99 m) of the noise source flushed, but then returned within minutes of the disturbance. Vessel noise from the Proposed Action is not expected to be as high as the noise level in the Beason study.

Alternative 1

Vessel noise from both Alternatives 1 and 2 would result in the same potential for effects to marine birds because the same vessels would be utilized for both alternatives. Due to the insignificant and short-term reactions to vessel noise, and in accordance with E.O. 12114, vessel noise associated with Alternative 1 would not result in significant harm to marine birds. Vessel noise associated with Alternative 1 would not result in a significant adverse effect on populations of marine birds protected under the MBTA.

Alternative 2 (Preferred Alternative)

Vessel noise from both Alternatives 1 and 2 would result in the same potential for effects to marine birds because the same vessels would be utilized for both alternatives. Due to the insignificant and short-term reactions to vessel noise, and in accordance with E.O. 12114, vessel noise associated with Alternative 2 would not result in significant harm to marine birds. Vessel noise associated with Alternative 2 would not result in a significant adverse effect on populations of marine birds protected under the MBTA.

4.3.2.3 Fish

The fish species located in the Study Area include those that are closely associated with the deep ocean habitat of the Beaufort Sea (Section 3.2.2.3). Only about 30 species are known to occur in the Arctic waters of the Beaufort Sea (Christiansen and Reist 2013). Aquatic systems of the Arctic undergo extended seasonal periods of ice cover and other harsh environmental conditions. Fish inhabiting such systems must be biologically and ecologically adapted to surviving such conditions. Important environmental factors that Arctic fish must contend with include reduced light, seasonal darkness, ice cover, low biodiversity, and low seasonal productivity. No ESA-listed fish species occur within the Study Area.

Acoustic stressors that may have potential impacts on fish include non-impulsive acoustic sources, icebreaking noise, and vessel noise. Physical stressors that may have potential impacts on fish include icebreaking (physical impacts), vessel and in-water device strike, and bottom disturbance. The only stressor associated with expended materials that may have potential impacts on fish is entanglement.

4.3.2.3.1 Non-Impulsive Acoustic Sources

Potential Impacts

As discussed in Section 3.2.2.3.2, data on hearing sensitivities of fish species occurring in the Study Area are not known. Research on fish hearing is limited; however, there is the potential for a fish with hearing sensitivities yet to be determined to perceive the sound of the Proposed Action. The region of hearing sensitivity in fish is generally within the lower frequencies, ranging from 100 to 400 Hz (Popper 2003). PTS has not been documented in fish. A study regarding mid-frequency sonar exposure by Halvorsen et al. (2012) found that for temporary hearing loss or similar negative impacts to occur, the noise needed to be within the fish's individual hearing frequency range; external factors, such as developmental history of the fish or environmental factors, may result in differing impacts to sound exposure in fish of the same species. The sensory hair cells of the inner ear in fish can regenerate after they are damaged, unlike in mammals where sensory hair cell loss is permanent (Lombarte et al. 1993; Smith et al. 2006). As a consequence, any hearing loss in fish may be as temporary as the timeframe required to repair or replace the sensory cells that were damaged or destroyed (Smith et al. 2006), and no permanent loss of hearing in fish would result from exposure to sound.

Studies of the effects of long-duration sounds with sound pressure levels (SPLs) below 170–180 dB re 1 μ Pa indicate that there is little to no effect of long-term exposure on species that lack notable anatomical hearing specialization (Amoser and Ladich 2003; Scholik and Yan 2001; Smith et al. 2004a, 2004b; Wysocki et al. 2006). The longest of these studies exposed young rainbow trout (*Onorhynchus mykiss*) to a level of noise equivalent to one that fish would experience in an aquaculture facility (e.g., on the order of 150 dB re 1 μ Pa) for about nine months. The investigators found no effect on hearing (i.e., TTS) as compared to fish raised at 110 dB re 1 μ Pa. Though these studies have not directly determined impacts to the fish expected to be present within the Study Area, it can be assumed that they would react in a similar manner to sound exposure.

Behavioral responses to noise in wild fish could alter the behavior of a fish in a manner that would affect its way of living, such as where it tries to locate food or how well it can locate a potential mate.

Behavioral responses to loud noise could include a startle response, such as the fish swimming away from the source, the fish “freezing” and staying in place, or scattering (Popper 2003).

Fish use sounds to detect both predators and prey, and for schooling, mating, and navigating (Myrberg 1981; Popper 2003). Masking of sounds associated with these behaviors could have impacts to fish by reducing their ability to perform these biological functions. Any noise (i.e., unwanted or irrelevant

sound, often of an anthropogenic nature) detectable by a fish can prevent the fish from hearing biologically important sounds including those produced by prey or predators (Myrberg 1981; Popper 2003). The immediate elevated stress response to anthropogenic noise can inhibit the survival of certain prey fish, reducing their ability to react to predator attacks during noise exposure (Simpson et al. 2016). The frequency of the sound is an important consideration for fish because many marine fish are limited to detection of the particle motion component of low frequency sounds at relatively high sound intensities (Amoser and Ladich 2005). Some of the frequencies of the non-impulsive acoustic sources associated with the Proposed Action are higher than those expected to be perceived by those species within the Study Area; therefore, masking is not likely as the mid- and high-frequency sources are not within the hearing range a fish would use to detect predators or prey. Behavioral responses are possible for those fish close to the active sonar sources since most would be within or near the frequencies of highest hearing sensitivity (100 to 400 Hz), but there is little evidence of these responses at most of the frequency levels of the Proposed Action. Individual fish may avoid an area in which a low-frequency moored source is present, but population level effects would not be anticipated from placement of these sources.

Alternative 1

There is a chance that fish within the Study Area would be able to perceive the non-impulsive acoustic sources, and if perceived, that an individual fish would react; however, this reaction would be temporary or minimal, and the fish would be expected to resume normal behavior after exposure. In accordance with E.O. 12114, non-impulsive acoustic sources associated with Alternative 1 would not result in significant harm to fish.

Alternative 2 (Preferred Alternative)

There is a chance that fish within the Study Area would be able to perceive the non-impulsive acoustic sources, and if perceived, that an individual fish would react; however, this reaction would be temporary or minimal, and the fish would be expected to resume normal behavior after exposure. The inclusion of two VLF sources would not increase risk of acoustic exposure to fish in a meaningful way. In accordance with E.O. 12114, non-impulsive acoustic sources associated with Alternative 2 would not result in significant harm to fish.

4.3.2.3.2 Icebreaking Noise

Potential Impacts

Icebreaking noise has the potential to expose fish to both sound and general disturbance, which could result in short-term behavioral or physiological responses (e.g., avoidance, stress, increased heart rate). Misund (1997) found that fish ahead of a ship showed avoidance reactions at ranges of 160 to 489 ft (49 to 149 m). When the vessel passed over them, some species of fish exhibited sudden escape responses that included lateral avoidance or downward compression of the school. Avoidance behavior of vessels, vertically or horizontally in the water column, has been reported for cod and herring, and was attributed to vessel noise; similar behavioral response could be expected due to icebreaking noise. Vessel activity can also alter schooling behavior and swimming speed of fish (United Nations Environment Programme 2012).

It is not anticipated that temporary behavioral reactions (e.g., temporary cessation of feeding) would harm the individual fitness of a fish as individuals are expected to resume feeding upon cessation of the sound exposure and unconsumed prey would still be available in the environment. Furthermore, while icebreaking noise may influence the behavior of some fish species (e.g., startle response, masking), other fish species can be equally unresponsive (Becker et al. 2013).

Alternative 1

Icebreaking noise from both Alternatives 1 and 2 would result in the same potential for effects to fish, in that the same vessel would be utilized for both Alternatives. Due to the insignificant and short-term reactions to icebreaking noise, and in accordance with E.O. 12114, icebreaking noise associated with Alternative 1 would not result in significant harm to fish.

Alternative 2 (Preferred Alternative)

Icebreaking noise from both Alternatives 1 and 2 would result in the same potential for effects to fish, in that the same vessel would be utilized for both Alternatives. Due to the insignificant and short-term reactions to icebreaking noise, and in accordance with E.O. 12114, icebreaking noise associated with Alternative 2 would not result in significant harm to fish.

4.3.2.3.3 Vessel Noise

Potential Impacts

Vessel noise has the potential to expose fish to both sound and general disturbance, which could result in short-term behavioral or physiological responses (e.g., avoidance, stress, increased heart rate). Noise from the vessels associated with the Proposed Action is not expected to impact fish, as available evidence does not suggest that ship noise can injure or kill a fish (Popper 2014). Misund (1997) found that fish ahead of a ship showed avoidance reactions at ranges of 161 to 489 ft (49 to 149 m). When the vessel passed over them, some species of fish exhibited sudden escape responses that included lateral avoidance or downward compression of the school. Avoidance behavior of vessels, vertically or horizontally in the water column, has been reported for cod and herring, and was attributed to vessel noise. Vessel activity can also alter schooling behavior and swimming speed of fish (United Nations Environment Programme 2012). Using acoustic telemetry and modeled ship noise, one recent study found that the presence of vessels in the highly trafficked Resolute Bay in Nunavut, Canada resulted in home range displacement of Arctic cod near the vessel (Ivanova et al. 2020). Individuals altered their swimming behavior by moving away and food searching behavior was disrupted; it is unknown for how long the behavioral disruption lasted before fish resumed their normal behavioral pattern (Ivanova et al. 2020).

It is not anticipated that temporary behavioral reactions (e.g., temporary cessation of feeding) would harm the individual fitness of a fish as individuals are expected to resume feeding upon cessation of the sound exposure and unconsumed prey would still be available in the environment. Furthermore, while vessel sounds may influence the behavior of some fish species (e.g., startle response, masking), other fish species can be equally unresponsive (Becker et al. 2013).

Alternative 1

Vessel noise from both Alternatives 1 and 2 would result in the same potential for effects to fish, in that the same vessels would be utilized for both Alternatives. Due to the insignificant and short-term reactions to vessel noise and in accordance with E.O. 12114, vessel noise associated with Alternative 1 would not result in significant harm to fish.

Alternative 2 (Preferred Alternative)

Vessel noise from both Alternatives 1 and 2 would result in the same potential for effects to fish, in that the same vessels would be utilized for both Alternatives. Due to the insignificant and short-term reactions to vessel noise and in accordance with E.O. 12114, vessel noise associated with Alternative 2 would not result in significant harm to fish.

4.3.2.3.4 Icebreaking (Physical Impacts)

Potential Impacts

Fish species within the Study Area are distributed throughout the surface, water column, and seafloor. Based on the existing scientific information on Arctic cod in the Beaufort Sea, Arctic cod would be nearshore, feeding in late summer and early autumn. As the autumn ice thickens and eventually freezes to the bottom in shallow nearshore areas, Arctic cod would move farther offshore where they spawn under the ice between November and February (Office of Environment Alaska OCS Region 2012). Icebreaking associated with the Proposed Action is scheduled during the warm season, between July and October in the deep water area of the Study Area. Arctic cod are expected to be nearshore during this timeframe and would not likely be exposed to icebreaking activities. However, Arctic cod have been observed among broken ice floes in the wake of icebreakers or splashed on top of ice floes (Crawford 2003; Gradinger and Bluhm 2004) indicating that individual Arctic cod and other ice-associated fish could be injured or killed along the icebreaker track lines. However, mortality is unlikely, because fish are highly mobile and are likely to avoid icebreaking activities since CGC HEALY would travel at a maximum speed of 3 knots during icebreaking activities.

Alternative 1

Icebreaking from both Alternatives 1 and 2 would result in the same potential for effects to fish, in that the same vessel would be utilized for both Alternatives. The icebreaking may result in short-term and local displacement of fishes in the water column. However, these behavioral reactions are not expected to result in substantial changes to an individual's fitness, or population recruitment, and are not expected to result in any harm at the population-level. Isolated cases of icebreaking striking a fish would potentially injure or result in the mortality of individuals, but would not result in population-level effects. In accordance with E.O. 12114, physical impacts from icebreaking associated with Alternative 1 would not result in significant harm to fish.

Alternative 2 (Preferred Alternative)

Icebreaking from both Alternatives 1 and 2 would result in the same potential for effects to fish, in that the same vessel would be utilized for both Alternatives. The icebreaking may result in short-term and local displacement of fishes in the water column. However, these behavioral reactions are not expected to result in substantial changes to an individual's fitness, or population recruitment, and are not expected to result in any harm at the population-level. Isolated cases of icebreaking striking a fish would potentially injure or result in the mortality of individuals, but would not result in population-level effects. In accordance with E.O. 12114, physical impacts from icebreaking associated with Alternative 2 would not result in significant harm to fish.

4.3.2.3.5 Vessel and In-Water Device Strike

Potential Impacts

Fish species within the Study Area are distributed throughout the surface, water column, and seafloor. Seafloor species would be unlikely to come into contact with vessels and in-water devices. Arctic cod and other pelagic species would be exposed to vessels and in-water devices, as their distribution within the water column is from the surface to 1,312 ft (400 m), as discussed in Section 3.2.2.3.

The potential for fish to be struck by vessels or in-water devices from the Proposed Action would be extremely low because most fish can detect and avoid vessel and in-water device movements. Fish would not be impacted by any wave produced by a vessel in motion. The fish lateral line system can detect changing water flow, which would allow fish to evade approaching objects (Stewart et al. 2014).

As a vehicle approaches a fish, the fish could have a behavioral or physiological response (e.g., swimming away and increased heart rate) as the passing vehicle displaces them. Potential harm from exposure to vessels, vehicles, and devices is not expected to result in substantial changes to an individual's overall behavior patterns, or species fitness and recruitment, and is not expected to result in any harm at the population-level. Any isolated cases of vessels or vehicles striking an individual could injure that individual, impacting its fitness, but not to the extent that there would be harm to the viability of populations based on the small number of vessels involved and the normal response of fish avoiding vessels and in-water devices.

Alternative 1

The potential for vessel and in-water device strike would be the same under Alternatives 1 and 2. Vessel and in-water device use may result in short-term and local displacement of fishes in the water column. However, these behavioral reactions are not expected to result in substantial changes to an individual's fitness, or species recruitment, and are not expected to result in any harm at the population-level, for the reasons described above. Isolated cases of vessel strike would potentially injure individuals, but would not result in population-level effects. In accordance with E.O. 12114, vessel and in-water device strike associated with Alternative 1 would not result in significant harm to fish.

Alternative 2 (Preferred Alternative)

The potential for vessel and in-water device strike would be the same under Alternatives 1 and 2. Vessel and in-water device use may result in short-term and local displacement of fishes in the water column. However, these behavioral reactions are not expected to result in substantial changes to an individual's fitness, or species recruitment, and are not expected to result in any harm at the population-level, for the reasons described above. Isolated cases of vessel strike would potentially injure individuals, but would not result in population-level effects. In accordance with E.O. 12114, vessel and in-water device strike associated with Alternative 2 would not result in significant harm to fish.

4.3.2.3.6 Bottom Disturbance

Potential Impacts

Items on the seafloor may attract benthic fish, including fish of the Orders Scorpaeniformes and Perciformes, but their sensory abilities allow them to avoid colliding with expended materials (Bleckmann and Zelick 2009). Those materials expended by the Proposed Action would fall to the seafloor in a manner dictated by ocean currents, but would be unlikely to do so nearby each other. Moorings would be anchored on the seabed. All sources would be deployed by shipboard winches, which would lower sources and receivers in a controlled manner. Anchors would be steel "wagon wheels" typically used for this type of deployment. Since fish are able to sense and avoid materials within their path, and expended materials would be drifting with the currents, rather than being self-propelled, it is highly unlikely that a fish would collide with an anchor or other tethering mechanism, either while it is sinking to the ocean floor or once it is on the ocean floor. Any turbidity associated with expended material hitting the seafloor would be expected to dissipate quickly, and not have any impacts on water quality.

Alternative 1

The impact to bottom habitats from bottom disturbance under Alternative 1 would be slightly lower than that under Alternative 2, based on the lower number of expended anchors associated with no VLF moorings being deployed. The disturbance would be localized and temporary as the equipment settles on the seafloor, which may cause scatter behavior in fish. However, the overall effects would be minimal due to the large size of the area and the low number of items expended over the expanse of the Study

Area. Therefore, in accordance with E.O. 12114, expended material associated with Alternative 1 would not result in significant harm to fish.

Alternative 2 (Preferred Alternative)

The impact to bottom habitats from bottom disturbance under Alternative 2 would be slightly higher than that under Alternative 1, based on the addition of two expended anchors from the VLF moorings. The disturbance would be localized and temporary as the equipment settles on the seafloor, which may cause scatter behavior in fish. However, the overall effects would be minimal due to the large size of the area and the low number of items expended over the expanse of the Study Area. Therefore, in accordance with E.O. 12114, expended material associated with Alternative 2 would not result in significant harm to fish.

4.3.2.3.7 Entanglement

Potential Impacts

The likelihood of fish being affected by an entanglement stressor is a function of the physical properties, location, and buoyancy of the object, as well as the behavior of the fish. Most entanglement observations involve abandoned or discarded nets, lines, and other materials that form loops or incorporate rings (Derraik 2002; Keller et al. 2010; Laist 1987; Macfadyen et al. 2009). A 25-year dataset assembled by the Ocean Conservancy (2010) reported that fishing line, rope, and fishing nets accounted for approximately 68 percent of fish entanglements, with the remainder due to encounters with various items such as bottles, cans, and plastic bags.

Fish entanglement occurs most frequently at or just below the surface or in the water column where objects are suspended; however, the physical properties (taut lines with no slack) of the materials associated with ARA are not expected to cause any entanglement. More fish species are entangled in coastal waters and the continental shelf than elsewhere in the marine environment because of higher concentrations of human activity (e.g., fishing, sources of entangling debris), higher fish abundances, and greater species diversity (Helfman et al. 2009; Macfadyen et al. 2009). The consequences of entanglement range from temporary and inconsequential to major physiological stress or mortality.

Some fish are more susceptible to entanglement in derelict fishing gear and other marine debris, compared to other fish groups. Physical features, such as rigid or protruding snouts of some elasmobranchs (e.g., the wide heads of hammerhead sharks), increase the risk of entanglement compared to fish with smoother, more streamlined bodies (e.g., lamprey and eels). Most other fish, except for jawless fish and eels that are too smooth and slippery to become entangled, are susceptible to entanglement gear specifically designed for that purpose (e.g., gillnets); however, no items would be expended that are designed to function as entanglement objects, nor are they designed to have slack or form loops. Expended materials have the potential to strike fish as they sink to the seafloor. Although individual fish may be at some marginal risk of injury, population-level impacts from these materials would not occur due to the dispersed nature and small amount of the expended material.

Alternative 1

Under Alternative 1, the potential for entanglement would be from mooring lines. Alternative 1 would have fewer mooring lines associated with the Proposed Action due to the exclusion of the VLF sources. Lines extending from the moorings would be retrieved at the completion of the Proposed Action. Entanglement of fish in the lines associated with the Proposed Action is not anticipated, given the mobility of the fish and the weighted (e.g., no slack or loops) lines that would be used. In accordance with E.O. 12114, entanglement associated with Alternative 1 would not result in significant harm to fish.

Alternative 2 (Preferred Alternative)

Under Alternative 2, the potential for entanglement would be from mooring lines, with an increased number of mooring lines as compared to Alternative 1 due to the inclusion of moored VLF sources. Entanglement of fish in the lines associated with the Proposed Action is not anticipated, given the mobility of the fish and the weighted (e.g., no slack or loops) lines that would be used. In accordance with E.O. 12114, entanglement associated with Alternative 2 would not result in significant harm to fish.

4.3.2.4 Essential Fish Habitat

The only species for which EFH has been designated within the Study Area is Arctic cod. Insufficient information is available to determine EFH for eggs, larvae, and early juveniles of Arctic cod. Essential Fish Habitat for late juvenile and adult Arctic cod within the Arctic Management Area occurs in waters from the nearshore to offshore areas along the continental shelf (0-656 ft [0-200 m]) and upper slope (656-1,640 ft [200-500 m]) throughout Arctic waters and often associated with ice floes which may occur in deeper waters (North Pacific Fishery Management Council 2009). EFH designation only occurs within the U.S. EEZ.

Acoustic stressors that may have potential impacts on EFH include non-impulsive acoustic sources and icebreaking noise. The only physical stressor that may have potential impacts on EFH is the physical impact of icebreaking.

4.3.2.4.1 Non-Impulsive Acoustic Sources

Potential Impacts

Non-impulsive acoustic sources could have an effect on the water column within the epipelagic zone, which is designated as EFH in a large portion of the Study Area, due to the increase in ambient sound level during the transmissions. However, this potential reduction in the quality of the acoustic habitat would be localized to the area of the sound sources. The quality of the water column would only be disturbed while the sound source is broadcasting and only in the area immediately ensonified around the non-impulsive acoustic source.

Moored and drifting non-impulsive acoustic sources would be deployed to transmit signals every day for the duration of the Proposed Action. Of those sources that would be frequently active, only the drifting IGB sources might be within designated EFH; no LF or VLF sources would be moored within Arctic cod EFH. The AMOS sources would operate intermittently, but outside of EFH. Sources on the drifting IGBs would be intermittent, with 30 second pings every 4 hours. Non-impulsive acoustic sources may be used in EFH during research cruises, but these would be on moving UUVs or on the vessel.

Exposure to individual fish would be limited, as would any noise added into the environment. The shallow moored sources would also be likely to increase ambient noise in the vicinity of the devices; only two would be deployed, which would limit overall impacts. Secondary effects to federally managed fish species (i.e., Arctic cod) are considered in Section 4.3.2.3.1 above.

Alternative 1

The quality of the water column as EFH would only be affected locally and temporarily overall. In accordance with E.O. 12114, non-impulsive acoustic sources associated with Alternative 1 would not result in significant harm to EFH.

Alternative 2 (Preferred Alternative)

The quality of the water column as EFH would only be affected locally and temporarily overall. In accordance with E.O. 12114, non-impulsive acoustic sources associated with Alternative 2 would not result in significant harm to EFH.

Pursuant to the MSA, an action may adversely affect EFH when it may reduce the quantity or quality of EFH, because it could be meaningfully measured or observed individually or cumulatively (regardless of duration or scale), or is likely to occur. Due to the potential for non-impulsive acoustic sources to alter Arctic cod EFH by temporarily and locally ensonifying the water column, non-impulsive acoustic sources associated with the Proposed Action may result in a reduction of the quantity or quality of EFH and therefore consultation under the MSA was initiated on July 6, 2022, with concurrence received from NMFS on July 29, 2022.

4.3.2.4.2 Icebreaking Noise

Potential Impacts

Icebreaking activities could have an effect on the features of the EFH, due to the increase in ambient sound level during icebreaking. However, this potential reduction in the quality of the acoustic habitat would be localized to the area of the icebreaking, which would be transient. The icebreaker is actively moving during icebreaking; therefore, any noise generated by the icebreaking activity would only affect the water column in close proximity to the ship and would be temporary in nature and would not ensonify the entire water column, but only the upper few meters. Icebreaking would be limited to up to eight days annually during a few weeks in the warm season, further reducing the amount of icebreaking noise entering the water column. Secondary effects to federally managed fish species (i.e., Arctic cod) are considered in Section 4.3.2.3.2 above.

Alternative 1

Icebreaking noise from both Alternatives 1 and 2 would result in the same potential for effects to EFH. In accordance with E.O. 12114, icebreaking noise associated with Alternative 1 would not result in significant harm to EFH.

Alternative 2 (Preferred Alternative)

Icebreaking noise from both Alternatives 1 and 2 would result in the same potential for effects to EFH. In accordance with E.O. 12114, icebreaking noise associated with Alternative 2 would not result in significant harm to EFH.

Pursuant to the MSA, an action may adversely affect EFH when it may reduce the quantity or quality of EFH, because it could be meaningfully measured or observed individually or cumulatively (regardless of duration or scale), or is likely to occur. While the quality of the water column as Essential Fish Habitat would only be affected locally and temporarily and cannot be meaningfully measured, in accordance with the MSA there would be a reduction in the overall quality of EFH and therefore consultation under the MSA was initiated on July 6, 2022, with concurrence received from NMFS on July 29, 2022.

4.3.2.4.3 Icebreaking (Physical Impacts)

Potential Impacts

EFH for Arctic cod in the Study Area includes areas of ice floe. Arctic cod are commonly found among ice floes and vessel movement through these areas could alter EFH via icebreaking activities. Icebreaking activities would be limited to up to eight days of active icebreaking annually during the warm season from July to October. Only areas of thick, wide concentrations of sea ice would require icebreaking by

the CGC HEALY. During August and September, these areas are expected to be at a minimum, which would reduce the impact to ice floes and EFH. The use of an icebreaking vessel may result in localized changes to Essential Fish Habitat as larger sheets of floating ice are broken down into smaller sizes, potentially reducing the coverage of ice in certain areas. However, icebreaking is not expected to significantly alter Arctic cod ice floe habitat as it will occur mainly in deep, ice-covered water and will be limited in duration. Secondary effects to federally managed fish species (i.e., Arctic cod) are considered in Section 4.3.2.3.4 above.

Alternative 1

Icebreaking from both Alternatives 1 and 2 would result in the same potential for effects to EFH, in that the same icebreaking vessel (CGC HEALY) would be utilized for both Alternatives. The use of an icebreaking vessel may result in localized changes to EFH as larger sheets of floating ice are broken down into smaller sizes. However, icebreaking is not expected to significantly alter Arctic cod ice floe habitat. Therefore, in accordance with E.O. 12114 physical impacts from icebreaking associated with Alternative 1 would not result in significant harm to EFH.

Alternative 2 (Preferred Alternative)

Icebreaking from both Alternatives 1 and 2 would result in the same potential for effects to EFH, in that the same icebreaking vessel (CGC HEALY) would be utilized for both Alternatives. The use of an icebreaking vessel may result in localized changes to EFH as larger sheets of floating ice are broken down into smaller sizes. However, icebreaking is not expected to significantly alter Arctic cod ice floe habitat. Therefore, in accordance with E.O. 12114 physical impacts from icebreaking associated with Alternative 2 would not result in significant harm to EFH.

Pursuant to the MSA, an action may adversely affect EFH when it may reduce the quantity or quality of EFH, because it could be meaningfully measured or observed individually or cumulatively (regardless of duration or scale), or is likely to occur. Due to the potential for icebreaking to alter Arctic cod EFH, physical impacts from icebreaking associated with the Proposed Action may result in a reduction of the quantity or quality of EFH, and therefore, consultation under the MSA was initiated on July 6, 2022, with concurrence received from NMFS on July 29, 2022.

4.3.2.5 Marine Mammals

Nine marine mammal species, which include three cetaceans, five pinnipeds, and the polar bear, are likely to occur in the Study Area during the Proposed Action. Marine mammals are found throughout the Study Area, including on the sea ice and within the water column. ESA-listed marine mammals, including the bearded seal, bowhead whale, polar bear, and ringed seal, would be present in the Study Area.

Acoustic stressors that may have potential impacts on marine mammals include non-impulsive acoustic sources, icebreaking noise, and vessel noise. Physical stressors that may have potential impacts on marine mammals include icebreaking (physical impacts) and vessel and in-water device strike. The stressors associated with expended materials that may have potential impacts on marine mammals are entanglement and ingestion.

4.3.2.5.1 Non-Impulsive Acoustic Sources

Potential Impacts

In assessing the potential effects on marine mammals from the Proposed Action, a variety of factors must be considered, including source characteristics, animal presence, animal hearing range, duration of exposure, and impact thresholds for species that may be present. Potential acoustic impacts could

include PTS, TTS, or behavioral effects. To make these assessments, a model was used to quantitatively estimate the potential number of exposures that could occur, followed by a qualitative analysis to account for other factors not reflected by the model.

The Navy Acoustic Effects Model (NAEMO) was used to produce a quantitative estimate of PTS, TTS, and behavioral exposures for marine mammals. The Navy then further analyzed the data and conducted an in-depth qualitative analysis of the species distribution and likely responses to the non-impulsive acoustic sources based on available scientific literature. The determination of the effects to marine mammals was based on this combination of quantitative and qualitative analyses. Additional details on the acoustic modeling can be found in Appendix B.

Quantitative Analysis

A quantitative analysis of the potential effects to marine mammals from the proposed non-impulsive acoustic sources was conducted using a method that calculates the total sound exposure level (SEL) and maximum sound pressure level that a marine mammal may receive from the non-impulsive acoustic sources. NAEMO was used for all modeling analysis (U.S. Department of the Navy 2017d). Environmental characteristics (e.g., bathymetry, wind speed, and sound speed profiles) and source characteristics (i.e., source level, source frequency, transmit pulse length and interval, horizontal and vertical beam width and source depth) were used to determine the propagation loss of the acoustic energy, which was calculated using the Comprehensive Acoustic System Simulation/Gaussian Ray Bundle (CASS/GRAB) propagation model. Additionally, an under-ice model (Oceanographic and Atmospheric Master Library [OAML] ICE) for surface interaction was implemented in NAEMO. The propagation loss then was used in NAEMO to create acoustic footprints. The NAEMO model then simulated source movement through the Study Area and calculated sound energy levels around the source. Animats, or representative animals, were distributed based on density data estimates obtained from Kaschner et al. (2006) and Cañadas et al. (2020). The Navy used a Seasonal Relative Environmental Suitability model (Kaschner et al. 2006), based on seasonal habitat preferences and requirements of known occurrences, such as temperature, bathymetry, and distance to land data and literature review, because occurrence information for marine mammals in the Study Area is not well known. Empirical data is coupled with Relative Environmental Suitability modeling data to generate predictions of density data for locations where no survey data exist. Densities derived from survey data were used when available, primarily in the southernmost portions of the Study Area during the warm season (Cañadas et al. 2020). The energy received by each animat distributed within the model was summed into a total sound exposure level. Additionally, the maximum sound pressure level received by each animat was also recorded.

NAEMO provides the predicted number of exposures that could result in effects as determined by the application of acoustic threshold criteria. Criteria and thresholds for measuring these effects induced from underwater acoustic energy have been established for marine mammals. Marine mammal criteria were established based on the following hearing groups: low-, mid-, and high-frequency cetaceans, otariid and non-phocid marine carnivores, and phocid pinnipeds. A summary of physiological and behavioral criteria is provided in Table 4-1 for groups of marine mammals that are found within the Study Area. The thresholds established for physiological effects (sound exposure levels for PTS and TTS) for groups of marine mammals that are found in the Study Area are described in detail in National Marine Fisheries Service (2016), and behavioral effects are described in detail in Department of the Navy (2017a).

Table 4-1. Acoustic In-Water Criteria and Thresholds for Predicting Physiological and Behavioral Effects on Marine Mammals Potentially Occurring in the Study Area

<i>Group</i>	<i>Species</i>	<i>Behavioral Criteria</i>	<i>Physiological Criteria</i>	
			<i>Onset TTS</i>	<i>Onset PTS</i>
Low Frequency Cetaceans	Gray whale, bowhead whale	Low-Frequency BRF dose response function*	179 dB SEL cumulative	199 dB SEL cumulative
Mid Frequency Cetaceans	Beluga whale	Mid-Frequency BRF dose response function*	178 dB SEL cumulative	198 dB SEL cumulative
Phocidae (in water)	Bearded seal, ribbon seal, spotted seal, ringed seal	Pinniped Dose Response Function*	181 dB SEL cumulative	201 dB SEL cumulative
Otariidae (in water) and other non-phocid marine carnivores	Polar bear, Pacific walrus	Pinniped Dose Response Function*	199 dB SEL cumulative	219 dB SEL cumulative

BRF = Behavioral Response Function

*See Figure 4-1




Figure 4-1. The Bayesian biphasic dose-response BRF for A) Odontocetes, B) Pinnipeds, and C) Mysticetes.

Note: The blue solid line represents the Bayesian Posterior median values, the green dashed line represents the biphasic fit, and the grey represents the variance. [X-Axis: Received Level (dB re 1 μ Pa), Y-Axis: Probability of Response]

The results from the NAEMO acoustic analysis are provided in Table 4-2. Non-impulsive acoustic sources would be active throughout the duration of the Proposed Action. Although the Proposed Action would occur over a multi-year period, estimated acoustic exposures were calculated on an annual basis. Exposures were calculated based on deployment of all sources. No marine mammals are likely to experience received SELs that may result in PTS or TTS. Beluga whales and ringed seals were calculated to potentially be exposed to sound pressure levels that may elicit a behavioral response.

Table 4-2. NAEMO-Calculated Marine Mammal Estimated Yearly Exposures

<i>Species</i>	<i>Alternatives 1 and 2</i>		
	<i>Behavioral</i>	<i>TTS</i>	<i>PTS</i>
Beluga whale	268	0	0
Bowhead whale ¹	0	0	0
Gray whale	0	0	0
Polar bear ¹	0	0	0
Bearded seal ¹	0	0	0
Ribbon seal	0	0	0
Ringed seal ¹	2,839	0	0
Spotted seal	0	0	0
Pacific Walrus	0	0	0

¹ESA listed species

These quantitative calculations were then analyzed qualitatively, taking into account the best available data on the species itself, and how the species has been observed to respond to similar types of influences.

Qualitative Analysis

No research has been conducted on the potential behavioral responses of ice-associated seals and other marine mammals occurring in the Study Area to the type of non-impulsive acoustic sources used during the Proposed Action. However, data are available on effects of non-impulsive acoustic sources (e.g., sonar transmissions) on marine mammals, which were assessed and incorporated into the findings of this analysis. Polar bears are anticipated to remain on the ice surface the majority of the time, and are not expected to be exposed to acoustic transmissions when in the water column since the acoustic sources would be active infrequently and widely distributed; effects from non-impulsive acoustic sources would be discountable.

Effects of Non-Impulsive Acoustic Sources on Phocids in Water

For non-impulsive sounds (i.e., similar to the sources used during the Proposed Action), data suggest that exposures of pinnipeds to sources between 90 and 140 dB re 1 μ Pa do not elicit strong behavioral responses; no data were available for exposures at higher received levels for Southall et al. (2007) to include in the severity scale analysis. Reactions of harbor seals (*Phoca vitulina*) were the only available data for which the responses could be ranked on the severity scale. For reactions that were recorded, the majority (17 of 18 individuals/groups) were ranked on the severity scale as a 4 (moderate change in movement, brief shift in group distribution, or moderate change in vocal behavior) or lower; the remaining response was ranked as a 6 (minor or moderate avoidance of the sound source). Additional data on hooded seals (*Cystophora cristata*) indicate avoidance responses to signals above 160–170 dB re 1 μ Pa (Kvadsheim et al. 2010), and data on gray (*Halichoerus grypus*) and harbor seals indicate avoidance response at received levels of 135–144 dB re 1 μ Pa (Götz et al. 2010). In each instance where food was available, which provided the seals motivation to remain near the source, habituation to the signals occurred rapidly. In the same study, it was noted that habituation was not apparent in wild seals where no food source was available (Götz et al. 2010). This implies that the motivation of the animal is necessary to consider in determining the potential for a reaction. In one study that investigated the under-ice movements and sensory cues associated with under-ice navigation of ice seals, acoustic transmitters (60–69 kHz at 159 dB re 1 μ Pa at 1 m) were attached to ringed seals (Wartzok et al. 1992a; Wartzok et al. 1992b). An acoustic tracking system then was installed in the ice to receive the acoustic signals and provide real-time tracking of ice seal movements. Although the frequencies used in this

study are at the upper limit of ringed seal hearing, the ringed seals appeared unaffected by the non-impulsive acoustic sources, as they were able to maintain normal behaviors (e.g., finding breathing holes).

Seals exposed to non-impulsive acoustic sources with a received sound pressure level within the range of calculated exposures (142–193 dB re 1 μ Pa), have been shown to change their behavior by modifying diving activity and avoiding the sound source (Götz et al. 2010; Kvadsheim et al. 2010). Although a minor change to a behavior may occur as a result of exposure to the sources in the Proposed Action, these changes would be within the normal range of behaviors for the animal (e.g., the use of a breathing hole further from the source, rather than one closer to the source, would be within the normal range of behavior) (Kelly et al. 1988).

Adult ringed seals spend up to 20 percent of their time in subnivean lairs during the winter season (Kelly et al. 2010a). Ringed seal pups spend about 50 percent of their time in the lair during the nursing period (Lydersen and Hammill 1993). Ringed seal lairs are typically used by individual seals (haul-out lairs) or by a mother with a pup (birthing lairs); large lairs used by many seals for hauling out are rare (Smith and Stirling 1975). The acoustic modeling does not account for seals within subnivean lairs, and all animals are assumed to be in the water and susceptible to hearing non-impulsive acoustic sources 100 percent of the time. Therefore, the acoustic modeling output likely represents an overestimate, given the percentage of time that ringed seals are expected to be in subnivean lairs, rather than in the water. Although the exact amount of transmission loss of sound traveling through ice and snow is unknown, it is clear that some sound attenuation would occur due to the environment itself. In air (i.e., in the subnivean lair), the best hearing sensitivity for ringed seals has been documented between 3 and 5 kHz; at higher frequencies, the hearing threshold rapidly increases (Sills et al. 2015).

If the non-impulsive acoustic sources are heard and are perceived as a threat, ringed seals within subnivean lairs could react to the sound in a similar fashion to their reaction to other threats, such as polar bears and arctic foxes (their primary predators), although the type of sound would be novel to them. Responses of ringed seals to a variety of human-induced noises (e.g., helicopter noise, snowmobiles, dogs, people, and seismic activity) have been variable; some seals entered the water and some seals remained in the lair (Kelly et al. 1988). However, in all instances in which observed seals departed lairs in response to noise disturbance, they subsequently reoccupied the lair (Kelly et al. 1988).

Ringed seal mothers have a strong bond with their pups and may physically move their pups from the birth lair to an alternate lair to avoid predation, sometimes risking their lives to defend their pups from potential predators (Smith 1987). Additionally, it is not unusual to find up to three birth lairs within 328 ft (100 m) of each other, probably made by the same female seal, as well as one or more haul-out lairs in the immediate area (Smith et al. 1991). If a ringed seal mother perceives the non-impulsive acoustic sources as a threat, the network of multiple birth and haul-out lairs allows the mother and pup to move to a new lair (Smith and Hammill 1981; Smith and Stirling 1975). However, the non-impulsive acoustic sources are unlike the low frequency sounds and vibrations felt from approaching predators. Additionally, the non-impulsive acoustic sources are not likely to impede a ringed seal from finding a breathing hole or lair, as captive seals have been found to primarily use vision to locate breathing holes and no effect to ringed seal vision would occur from the non-impulsive acoustic sources (Elsner et al. 1989; Wartzok et al. 1992a). It is anticipated that a ringed seal would be able to relocate to a different breathing hole relatively easily without impacting their normal behavior patterns.

Effects of Non-Impulsive Acoustic Sources on Mysticetes within the Study Area

While not many studies have been done on mysticete (i.e., low-frequency cetaceans) responses to sonar, behavioral response studies have been conducted. Although some strong responses have been

observed in mysticetes to sonar and other active acoustic sources, for the most part mysticete responses appear to be fairly moderate across all received levels. While some responses such as cessation of foraging or changes in dive behavior could carry short-term impacts, in all cases behavior returned to normal after the signal stopped. Mysticete responses also seem to be highly mediated by behavioral state, with no responses occurring in some behavioral states, and contextual factors and signal characteristics having more impact than received level alone. Many of the contextual factors resulting from the behavioral response studies (e.g., close approaches by multiple vessels or tagging) would not occur during the Proposed Action. Mysticete behavioral responses to acoustic transmission from the Proposed Action would likely be a result of the animal's behavioral state and prior experience rather than external variables such as ship proximity; thus, if significant behavioral responses occur they would likely be short-term. In fact, no significant behavioral responses such as panic, stranding or other severe reactions have been observed during monitoring of actual training exercises (Smultea and Mobley 2009; U.S. Department of the Navy 2011, 2014; Watwood et al. 2012).

Effects of Non-Impulsive Acoustic Sources on Odontocetes within the Study Area

Research shows that if odontocetes do respond to a sound, they may react in a number of ways depending on the characteristics of the sound source, their experience with the sound source, and whether they are migrating or on seasonal grounds (i.e., breeding or feeding grounds). Behavioral reactions may include an alert response; terminating a feeding dive and surfacing; a shift in normal dive depth (e.g., several consecutive shallow dives), or swimming away. Animals disturbed while engaged in activities, such as feeding or reproductive behaviors, may be more likely to ignore or tolerate the disturbance and continue with their behavior. Therefore, most behavioral reactions from odontocetes are likely to be short-term, with low to moderate severity.

Behavioral research indicates that most odontocetes would likely avoid sound sources at levels that would cause any hearing loss (i.e., TTS) (Finneran 2015; Finneran et al. 2010; Finneran et al. 2005; Finneran and Schlundt 2003; Finneran and Schlundt 2010; Finneran and Schlundt 2013; Mooney et al. 2009; Popov et al. 2011; Schlundt et al. 2000).

In studies that examined sperm whales and false killer whales (both in the mid-frequency cetacean hearing group), the marine mammals showed temporary cessation of calling and avoidance of sonar sources (Akamatsu et al. 1993; Watkins and Schevill 1975). Sperm whales resumed calling and communication approximately two minutes after the pings stopped (Watkins and Schevill 1975). False killer whales did move away from the sound source, but returned to the area between 0 and 10 minutes after the end of the transmissions (Akamatsu et al. 1993). Many of the contextual factors resulting from the behavioral response studies (e.g., close approaches by multiple vessels or tagging) would not occur during the Proposed Action. Odontocete behavioral responses to acoustic transmissions from non-impulsive acoustic sources used during the Proposed Action would likely be a result of the animal's behavioral state and prior experience rather than external variables such as ship proximity; thus, if significant behavioral responses occur they would likely be short-term. In fact, no significant behavioral responses such as panic, stranding or other severe reactions have been observed during monitoring of actual training exercises (Smultea and Mobley 2009; U.S. Department of the Navy 2011, 2014; Watwood et al. 2012).

Effects of Non-Impulsive Acoustic Sources on Walrus within the Study Area

Typical behavioral responses by Pacific walrus to disturbances include: altered headings; increased swimming rates; increased vigilance; changes in dive, surfacing, respiration, feeding, and vocalization patterns; and hormonal stress production (Ellison et al. 2012; Richardson et al. 1995b; Southall et al. 2007). Low-level reactions are common and can be caused by both natural and anthropogenic sources.

Significant behavioral responses include displacement from preferred foraging areas, increased stress levels or energy expenditures, or cessation of feeding. Noise may evoke behavioral responses in addition to the possible impacts to hearing (i.e., TTS or PTS). Passive acoustic monitoring conducted during 2016 cable laying on the Beaufort and Chukchi shelf documented Pacific walrus vocalizing in the local area before and after, but not during, cable-laying work. There is a possibility that the Pacific walrus either moved or ceased vocalizing due to the project's noise (Owl Ridge Natural Resource Consultants Inc 2017). This may be an indication of auditory masking (a change in the ability to detect relevant sounds in the presence of other sounds (Wartzok et al. 2003)). The biological implications of anthropogenic masking among walrus are unknown, but if the Pacific walrus' response to masking is to leave the area, then the physiological costs are similar to those of other disturbances that trigger the same response. The response of walrus to disturbance stimuli is highly variable. Observations by walrus hunters and researchers suggest that males tend to be more tolerant of disturbances than females, and individuals tend to be more tolerant than groups; females with dependent calves are considered least tolerant of disturbances.

The most likely behaviorally significant responses that the Proposed Action could evoke among Pacific walrus include temporary cessation of feeding, resting, or communicating. Effects of these types of mid-level responses include increased energy expenditures and stress levels. Energetic costs are incurred from loss of forage and energy expended while travelling to another region.

Similarly, a controlled exposure study to simulated mid-frequency sonar was conducted with U.S. Navy California sea lions (*Zalophus californianus*; an appropriate surrogate for Pacific walrus based on similarities in hearing and ear morphology) at the Navy Marine Mammal Program facility specifically to study behavioral reactions (Houser et al. 2013a). Animals were trained to swim across a pen, touch a panel, and return to the starting location. During transit, a simulated mid-frequency sonar signal was played. Behavioral reactions included increased respiration rates, prolonged submergence, and refusal to participate, among others. Younger animals were more likely to respond than older animals, while some sea lions did not respond consistently at any sound source level.

Alternative 1

Non-impulsive acoustic sources from both Alternatives 1 and 2 would have the same potential for effects to marine mammals and are expected to result in, at most, minor to moderate avoidance responses of animals, over short and intermittent periods of time.

Non-impulsive acoustic sources associated with the Proposed Action may affect, and are likely to adversely affect, ringed seals. The Proposed Action would not result in the destruction or adverse modification of critical habitat for ringed seals or polar bears. Since quantitative modeling in NAEMO showed no acoustic exposures at or above the behavioral thresholds, effects from non-impulsive acoustic sources associated with the Proposed Action would be insignificant or discountable; therefore, the Proposed Action would have no effect on the ESA-listed bowhead whale, bearded seal, and polar bear. In accordance with E.O. 12114, non-impulsive acoustic sources from Alternative 1 would not result in significant harm to marine mammals.

Alternative 2 (Preferred Alternative)

Non-impulsive acoustic sources from both Alternatives 1 and 2 would have the same potential for effects to marine mammals and are expected to result in, at most, minor to moderate avoidance responses of animals, over short and intermittent periods of time. The addition of VLF sources under Alternative 2 does not result in additional takes of marine mammals as compared to Alternative 1.

Due to the number of behavioral exposures, the Navy submitted an application for an IHA with NMFS for Level B take of ringed seals and beluga whales for the period from September 2022 – September 2023 on March 21, 2022; a Notice of availability of the draft IHA was published in the Federal Register on July 26, 2022. Since acoustic sources used during the Proposed Action can change based on previous results and changing scientific objectives, annual requests for IHAs would be completed throughout the duration of the Proposed Action to capture any changes in estimated take numbers.

In accordance with the ESA, the Navy consulted with NMFS based on the determination the Proposed Action may affect, and is likely to adversely affect, ringed seals. The Proposed Action would not result the destruction of adverse modification of critical habitat for ringed seals or polar bears. Since quantitative modeling in NAEMO showed no acoustic exposures at or above the behavioral thresholds, effects from non-impulsive acoustic sources associated with the Proposed Action would be insignificant or discountable; therefore, the Proposed Action would have no effect on the ESA-listed bowhead whale, bearded seal, and polar bear. Formal consultation was initiated with NMFS regarding ringed seals on April 29, 2022.

In accordance with E.O. 12114, non-impulsive acoustic sources from Alternative 2 would not result in significant harm to marine mammals.

4.3.2.5.2 Icebreaking Noise

Potential Impacts

Icebreaking noise was modeled using similar methods to those described in Section 4.1.1.2. Below is a quantitative analysis of the modeling results for CGC HEALY icebreaking, as well as a qualitative analysis for icebreaking noise.

Quantitative Analysis

The underwater radiated noise signature for icebreaking in the central Arctic Ocean by CGC HEALY during different types of ice cover was characterized in Roth et al. (2013). The radiated noise signatures were characterized for various fractions of ice cover. For modeling, the 8/10 and 3/10 ice cover were used. Each modeled day of icebreaking consisted of 16 hours of 8/10 ice cover and 8 hours of 3/10 ice cover. Icebreaking was modeled for eight days each year. Figure 5a and 5b in Roth et al. (2013) depicts the source spectrum level versus frequency for 8/10 and 3/10 ice cover, respectively. The sound signature of each of the ice coverage levels was broken into 1-octave bins (Table 4-3 and Table 4-4). In the model, each bin was included as a separate source on the modeled vessel. When these independent sources go active concurrently, they simulate the sound signature of CGC HEALY. The modeled source level summed across these bins was 196.2 dB for the 8/10 signature and 189.3 dB for the 3/10 ice signature. These source levels are a good approximation of the icebreaker's observed source level (provided in Figure 4b of (Roth et al. 2013)). Each frequency and source level was modeled as an independent source, and applied simultaneously to all of the animats within NAEMO. Each second was summed across frequency to estimate sound pressure level (root mean square [SPL_{RMS}]). This value was incorporated into the behavioral risk function to estimate behavioral exposures. For PTS and TTS determinations, sound exposure levels were summed over the duration of the test and the transit through the Study Area.

Table 4-3. Modeled Bins for 8/10 Ice Coverage (Full Power) Ice Breaking on CGC HEALY

<i>Frequency (Hz)</i>	<i>Source Level (dB)</i>
25	189
50	188
100	189
200	190
400	188
800	183
1600	177
3200	176
6400	172
12800	167

Table 4-4. Modeled Bins for 3/10 Ice Coverage (Quarter Power) Ice Breaking on CGC HEALY

<i>Frequency (Hz)</i>	<i>Source Level (dB)</i>
25	187
50	182
100	179
200	177
400	175
800	170
1600	166
3200	171
6400	168
12800	164

The output from the acoustic model is the calculated number of marine mammals exposed at or above acoustic effects thresholds listed in Table 4-5. Icebreaking could occur on any research cruise using CGC HEALY. Although the Proposed Action would occur over a multi-year period, estimated acoustic exposures were calculated on an annual basis. Due to the changing environmental conditions in the Study Area it is unknown how long icebreaking would occur each year. However, it is anticipated from previous cruises that no more than eight days of icebreaking would be required to reach the areas for deployment during the summer months. Exposures provided in Table 4-5 are for the maximum amount of icebreaking during a CGC HEALY cruise. A maximum of one icebreaking cruise would occur annually.

Table 4-5. Model-Calculated Yearly Acoustic Exposures for CGC HEALY Icebreaking

<i>Species</i>	<i>Alternatives 1 and 2</i>		
	<i>Behavioral</i>	<i>TTS</i>	<i>PTS</i>
Beluga whale	21	0	0
Bowhead whale ¹	0	0	0
Gray whale	0	0	0
Polar bear ¹	0	0	0
Bearded seal ¹	0	0	0
Ringed seal ¹	538	1	0
Ribbon seal	0	0	0
Spotted seal	0	0	0
Pacific walrus	0	0	0

¹ESA Listed Species

The quantitative analysis calculated that most marine mammals in the Study Area would not experience behavioral response, TTS, or PTS from the Proposed Action under either Alternative for sound generated from icebreaking. However, modeling results indicated that icebreaking would result in 538 behavioral exposures and 1 TTS exposure to ringed seals and 21 behavioral exposures to beluga whales under both Alternatives, suggesting the possibility of eliciting a behavioral response.

The likelihood of a behavioral response is dependent upon the received sound pressure level. NAEMO provides two outputs. The first is the number of animals recorded with received levels within 1 dB bins at and greater than 100 dB re 1 µPa, prior to effect thresholds being applied (referred to as unprocessed animal exposures). These results are used to determine if a marine mammal may be exposed to the acoustic energy resulting from the Proposed Action, but they do not infer that any such exposure results in an effect from the action. The second output, referred to as calculated exposures (as seen in Table 4-5), is the predicted number of exposures that could result in effects from the Proposed Action after the application of the behavioral risk function (Figure 4-1) and acoustic threshold criteria. Additional details on the acoustic modeling can be found in Appendix B.

As discussed above, the quantitative output calculated that 538 ringed seals and 21 beluga whales could be exposed to sound pressure levels that may elicit a behavioral response. These quantitative calculations are then analyzed qualitatively by marine biologists and acoustic experts, taking into account the best available data on the species itself, and how the species has been observed to respond to similar types of influences.

Qualitative Analysis – All Species

Variables such as the marine mammal’s gender, age, the activity it is engaged in during a sound exposure, its distance from a sound source, the number of sound sources, and whether the sound sources are approaching or moving away from the animal can be critically important in determining whether and how a marine mammal would respond to a sound source. Furthermore, the BRF does not differentiate between different types of behavioral reactions (e.g., area avoidance, diving avoidance, or alteration of natural behavior) or provide information regarding the predicted consequences to the animal of the reaction. At present, available data do not allow for incorporation of these other variables in the current BRF; they must be assessed qualitatively.

Southall et al. (2007) summarized data on behavioral reactions of pinnipeds in water to non-impulsive and impulsive sources (termed nonpulse and pulse sources, respectively), and ranked these reactions on a severity scale. For impulsive sources (e.g., airguns), data indicate that exposures between 150 and 180 dB re 1 µPa generally have limited potential to induce avoidance responses in pinnipeds, whereas higher received levels have exhibited some responses. Data used to identify the severity of behavioral reactions

are based primarily on ringed seals, but also include bearded and spotted seals (Blackwell et al. 2004; Harris et al. 2001; Miller et al. 2005). For received sound pressure levels between 140 dB re 1 μ Pa and 200 dB re 1 μ Pa, responses to impulsive sources were either 0 on the severity scale (no observable response; 49 percent of responses) or 6 on the severity scale (minor or moderate avoidance of the sound source; 51 percent of responses). The majority of the severity 6 responses (92 percent) occurred at sound pressure levels between 190 dB re 1 μ Pa and 200 dB re 1 μ Pa. Southall et al. (2007) found that within the range of sound pressure levels of approximately 150–190 dB re 1 μ Pa, 91 percent of individuals/groups were observed to have no response (severity scale ranking of 0) to the impulsive source. The remaining 9 percent were ranked on the severity scale as a 6, as minor or moderate avoidance reactions were observed. All of the reactions noted as a 6 on the severity scale (avoidance) are attributed to open-water use of a full-array of up to eleven 120 in³ (1,966 cm³) airguns. The avoidance of the area was relatively minor; some (but not all) seals avoided the zone within 492 ft (150 m) of the source, but did not move much beyond 820 ft (250 m) from the source. Additionally, the seismic operations with the full-array did not cause seals to desert the general area of the activity (Harris et al. 2001).

Although the icebreaking associated with the Proposed Action is not impulsive in a strict sense, the data on ringed seal reactions during seismic surveys nonetheless indicate that ringed seals have shown little reaction to noise disturbance in general within the sound pressure levels potentially received from the Proposed Action. Any behavioral reaction is expected to be short term, as icebreaking would occur in small areas and would be transient in nature, which reduces the probability of encountering a marine mammal during icebreaking activities. Behavioral reactions would be limited to swimming away, hauling out, diving underwater and, in some cases, avoidance behavior. These short-term reactions are not expected to significantly disrupt behavioral patterns such as migration, breathing, nursing, breeding, feeding and sheltering to a point where the behavior pattern is abandoned or significantly altered.

Marine mammals have been recorded in several instances altering and modifying their vocalizations to compensate for the masking noise from vessels, or other similar sounds (Holt et al. 2011; Parks et al. 2011). Vocal changes in response to anthropogenic noise can occur across the repertoire of sound production modes used by marine mammals, such as whistling, echolocation click production, calling, and singing. Changes to vocal behavior and call structure may result from a need to compensate for an increase in background noise. In cetaceans, vocalization changes have been reported from exposure to anthropogenic sources such as sonar, vessel noise, and seismic surveying.

Icebreaking noise has the potential to disturb marine mammals and elicit an alerting, avoidance, or other behavioral reaction (Huntington et al. 2015; Pirota et al. 2015; Williams et al. 2014). Icebreaking in fast ice during the spring can cause behavioral reactions in beluga whales. Erbe and Farmer (2000) calculated the zone of impacts to beluga whales from icebreakers in the Beaufort Sea using data from Canadian icebreakers. Beluga whales had a zone of behavioral disturbance out to 25 nm (46 km) in a shipping corridor near Beluga Bay, and 16 nm (30 km) when the icebreaker was over the abyssal plain in response to ramming noise from an icebreaker. Bowheads have been observed avoiding areas within 13 nm (25 km) of an icebreaking site (Richardson et al. 1995b). Icebreaking associated with the Proposed Action would occur in the August through October timeframe, which lessens the probability of a whale encountering the vessel.

Fay et al. (1984) compared the behavioral reactions of walrus to both icebreaking vessels and vessels in open water. Walrus tended to exhibit behavioral reactions to icebreaking at longer distances than from vessels in open water. Aerial surveys also indicated that walrus appeared to avoid areas within 5 to 8 nm (10 to 15 km) of an icebreaking vessel (Brueggeman et al. 1991). However, walrus are not located in the areas where icebreaking would occur and would not be affected by icebreaking.

Phocids are known to flush in the presence of vessels, including icebreakers, which can lead to displacement and potential separation of mothers and pups (Lomac-MacNair et al. 2019a). Ringed seals and bearded seals on pack ice showed various behaviors when approached by an icebreaking vessel; a majority of seals dove underwater when the ship was within 0.5 nm (0.93 km) while others remained on the ice. However, as icebreaking vessels came closer to the seals, most dove underwater. Ringed seals have also been observed foraging in the wake of an icebreaking vessel (Richardson et al. 1995b). Lomac-MacNair et al. (2019a) observed that four species of phocids (including bearded and ringed seals) flushed most frequently when the icebreaking vessel was within 1,969 ft (600 m), and no flushing was observed when the icebreaking vessel was greater than 2,625 ft (800 m) from the seal.

In studies by Alliston (1980; 1981), there was no observed change in the density of ringed seals in areas that had been subject to icebreaking. Alternatively, ringed seals may have preferentially established breathing holes in the ship tracks after the icebreaker moved through the area. Due to the time of year of the activity (August through October), ringed seals are not expected to be within the subnivean lairs nor pupping (Chapskii 1940; McLaren 1958; Smith and Stirling 1975). Therefore, icebreaking would not impact seals which could not visually detect an oncoming vessel.

Polar bears do not appear to be significantly affected by icebreaking noise and show very little reaction to icebreaking vessels (Richardson et al. 1995b). Polar bears that did react to icebreaker presence had the following reactions: walking away, running away, approaching, vigilance, and no reaction (Lomac-MacNair et al. 2019a). Vigilance was the most common observed reaction in a study by Smultea et al. (2016). Polar bear reactions involving walking or running away were brief in duration (less than five minutes) when the icebreaker was within 1,640 ft (500 m) or less.

Alternative 1

Icebreaking noise from both Alternatives 1 and 2 would have the same potential for effects to marine mammals and not be expected to have more than a short-term and temporary impact on any individual marine mammals exposed.

Non-impulsive acoustic sources associated with the Proposed Action may affect, and are likely to adversely affect, ringed seals. The Proposed Action would not result in the destruction of adverse modification of critical habitat for ringed seals or polar bears. Since quantitative modeling in NAEMO showed no acoustic exposures at or above the behavioral thresholds, effects from icebreaking noise associated with the Proposed Action would be insignificant or discountable; therefore, the Proposed Action would have no effect on the ESA-listed bowhead whale, bearded seal, and polar bear.

In accordance with E.O. 12114, icebreaking noise associated with Alternative 1 would not result in significant harm to marine mammals.

Alternative 2 (Preferred Alternative)

Icebreaking noise from both Alternatives 1 and 2 would have the same potential for effects to marine mammals and would not be expected to have more than a short-term and temporary impact on any individual marine mammals exposed.

Icebreaking noise associated with the Proposed Action may cause a behavioral reaction or TTS to the ringed seal and beluga whale. Due to the number of icebreaking noise exposures, the Navy submitted an application for an IHA with NMFS for Level B take of ringed seals and beluga whales for the period from September 2022 – September 2023 on March 21, 2022; a Notice of availability of the draft IHA was published in the Federal Register on July 26, 2022. Since the yearly amount of icebreaking is unknown, annual requests for IHAs would be completed throughout the duration of the Proposed Action.

In accordance with the ESA, the Navy consulted with NMFS based on the determination that the Proposed Action may affect, and is likely to adversely affect, ringed seals. The Proposed Action would not result in the destruction of adverse modification of critical habitat for ringed seals or polar bears. Since quantitative modeling in NAEMO showed no acoustic exposures at or above the behavioral thresholds, effects from icebreaking noise associated with the Proposed Action would be insignificant or discountable; therefore, the Proposed Action would have no effect on the ESA-listed bowhead whale, bearded seal, and polar bear. Formal consultation was initiated with NMFS regarding ringed seals on April 29, 2022.

In accordance with E.O. 12114, icebreaking noise associated with Alternative 2 would not result in significant harm to marine mammals.

4.3.2.5.3 Vessel Noise

Potential Impacts

Vessel noise associated with the Proposed Action could result from sound generated by the R/V Sikuliaq and CGC HEALY. Marine mammals have been recorded in several instances altering and modifying their vocalizations to compensate for the masking noise from vessels, or other similar sounds (Holt et al. 2011; Parks et al. 2011). Vocal changes in response to anthropogenic noise can occur across the repertoire of sound production modes used by marine mammals, such as whistling, echolocation click production, calling, and singing. Changes to vocal behavior and call structure may result from a need to compensate for an increase in background noise. In cetaceans, vocalization changes have been reported from exposure to anthropogenic sources such as sonar, vessel noise, and seismic surveying.

Since many marine mammals rely on sound to find prey, moderate social interactions, and facilitate mating (Pine et al. 2021), noise from anthropogenic sound sources like ships can interfere with these functions, but only if the noise spectrum overlaps with the hearing sensitivity of the marine mammal (Clark et al. 2009; Hatch et al. 2012; Southall et al. 2007; Southall et al. 2019). It is difficult to differentiate between behavioral responses to vessel sound and visual cues associated with the presence of a vessel; thus, it is assumed that both play a role in prompting reactions from animals. Vessel noise has the potential to disturb marine mammals and elicit an alerting, avoidance, or other behavioral reaction (Huntington et al. 2015; Pirotta et al. 2015; Williams et al. 2014). Most studies have reported that marine mammals react to vessel sounds and traffic when received levels were over 20 dB greater than ambient noise levels with short-term interruption of feeding, resting, or social interactions (Huntington et al. 2015; Magalhães et al. 2002; Merchant et al. 2014; Pirotta et al. 2015; Richardson et al. 1995b; Williams et al. 2014).

Some species respond negatively by retreating or responding to the vessel antagonistically, while other animals seem to ignore vessel noises altogether (Watkins 1986). Beluga whales can exhibit a variety of reactions from fleeing the area to no response at all to the vessel (Wartzok et al. 2003). Polar bears do not appear to be significantly affected by vessel noise. Some polar bears have been observed walking, running, and swimming away from approaching vessels, but these reactions were brief and localized. Other bears have been observed approaching vessels or having no reaction to vessels (Richardson et al. 1995b). The presence of boats and tourists did not significantly disturb walrus' haul-out behavior in common tourism sites (Øren et al. 2018).

Overall baleen whale responses to vessel noise and traffic are varied but are generally minor, and habituation or disinterest seems to be the predominant long-term response. When baleen whales avoid ships they do so by altering their swim and dive patterns to move away from the vessel, but no strong reactions have been observed. In fact, in many cases the whales do not appear to change their behavior at all. This may result from habituation by the whales, but may also result from reduced received levels

near the surface due to propagation, or due to acoustic shadowing of the propeller cavitation noise by the ship's hull. Based on studies on a number of species, mysticetes (such as bowhead and gray whales) are not expected to be disturbed by vessels that maintain a reasonable distance from them, though this varies with vessel size, geographic location, and tolerance levels of individuals. Bernasconi et al. (2012) observed the reactions of six individual baleen whales of unknown species at distances of 164 to 1,312 ft (50 to 400 m) from a fishing vessel conducting an acoustic survey of pelagic fisheries, with only a slight change in swim direction when the vessel began moving around the whales. Bowhead whales avoided the area around icebreaker ship noise and increased their time at the surface and number of blows (Richardson et al. 1995a). The noise generated from the R/V Sikuliaq is at a low source level (less than 160 dB) for the vessel speeds of the Proposed Action (Naval Sea Systems Command 2015), and at very small distances from the vessel the sound would be below the level capable of producing a behavioral response. The noise generated from CGC HEALY is at a similarly low source level at frequencies associated with vessel noise (100-1000 Hz). The noise from CGC HEALY when icebreaking is significantly higher (~ 10 dB) and will have enhanced propagation due to the introduction of additional low-frequency components (Roth et al. 2013).

In general, studies of pinniped reactions to vessels are limited. Pinnipeds have shown substantial tolerance to anthropogenic noise stressors. Pinniped reactions to vessels are variable and reports include a wide spectrum of possibilities from avoidance and alert, to cases where animals in the water are attracted, and cases on land where there is lack of significant reaction suggesting habituation to or tolerance of vessels (Richardson et al. 1995b). Another study of reactions of harbor seals hauled out on ice to cruise ship approaches in Disenchantment Bay, Alaska, revealed that animals are more likely to flush and enter the water when cruise ships approach within 1,640 ft. (500 m) and four times more likely when a cruise ship approaches within 328 ft. (100 m) (Jansen et al. 2010). Brueggeman et al (1992), observed ringed seals hauled out on ice sheets showing a short term behavioral reaction by diving into the water when a vessel came within 0.13-0.27 nm (0.25-0.5 km).

Vessels associated with the Proposed Action would not purposefully approach marine mammals and noise generated by these vessels is not expected to elicit significant behavioral responses. Such reactions are not expected to significantly disrupt behavioral patterns such as migration, breathing, nursing, breeding, feeding and sheltering to a point where the behavior pattern is abandoned or significantly altered or result in reasonably foreseeable takes of marine mammals.

Alternative 1

Vessel noise from both Alternatives 1 and 2 would result in the same potential for effects to marine mammals as the same vessels would be used. In accordance with the ESA, vessel noise may affect, but is not likely to adversely affect, ringed seals and bearded seals. The Proposed Action would not result in the destruction or adverse modification of critical habitat for ringed seals or polar bears. In accordance with E.O. 12114, vessel noise associated with Alternative 1 would not result in significant harm to marine mammals.

Alternative 2 (Preferred Alternative)

Vessel noise from both Alternatives 1 and 2 would result in the same potential for effects to marine mammals as the same vessels would be used. Vessel noise associated with the Proposed Action would not result in reasonably foreseeable takes under the MMPA. In accordance with the ESA, vessel noise may affect, but is not likely to adversely affect, the ESA-listed bowhead whale, bearded seal, and ringed seal. The Proposed Action would not result in the destruction or adverse modification of critical habitat for ringed seals or polar bears. Effects of vessel noise associated with the Proposed Action on polar bears would be discountable in significance; therefore, the Proposed Action would have no effect on the

ESA-listed polar bear. Formal consultation was initiated with NMFS regarding bowhead whale, bearded seal, and ringed seal on April 29, 2022.

In accordance with E.O. 12114, vessel noise associated with Alternative 2 would not result in significant harm to marine mammals.

4.3.2.5.4 Icebreaking (Physical Impacts)

Potential Impacts

Potential Impacts to Marine Mammals

Icebreaking could occur in the Study Area when transiting to deploy, maintain, or retrieve moored or drifting sources, at speeds of 3 to 6 knots. CGC HEALY could be icebreaking between July and October while the ice is at its lowest extent of the year. As discussed in Section 4.3.2.5.2, the noise associated with icebreaking activities is most likely to result in marine mammals swimming away from the icebreaking vessel or avoiding the area for a short period of time. Therefore, it is highly unlikely that icebreaking equipment would strike a marine mammal or cause any physical harm. Pinnipeds that haul out on the ice may be more susceptible to impacts caused by icebreaking, including the potential for habitat fragmentation (Lomac-MacNair et al. 2019a).

Bearded seals are strongly associated with sea ice habitat in the Arctic. In late spring and summer when icebreaking may occur, bearded seals move north as the ice edge recedes into the Chukchi and Beaufort Seas. However, some bearded seals stay near the edge of shorefast ice all winter and do not migrate south. Leads, polynyas, and other openings in the sea ice are important features of bearded seal habitat. Juvenile bearded seals tend to associate with sea ice less than adults and are often found in ice free areas such as bays and estuaries. The distribution of bearded seals appears to be strongly associated with shallow water and high biomass of the benthic prey they feed on. They are limited to feeding depths of less than 492-656 ft (150–200 m). Icebreaking may result in the temporary displacement of primary prey resources of ringed seals or bearded seals, but these species are expected to return to their normal behaviors shortly after the initial disturbance.

In the spring through the fall, areas with thick ice requiring icebreaking are expected to be at a minimum, which would reduce the impact to the ringed seals' proposed critical habitat. The ringed seal subnivean lairs are excavated in drifts over breathing holes in the ice, in which they rest, give birth, and nurse their pups for 5–9 weeks during late winter and spring (Chapskii 1940; McLaren 1958; Smith and Stirling 1975). Most ringed seals are born in early April and about a month after parturition, mating begins in late April and early May. Ringed seals are expected in the Study Area year-round, but during the Arctic summer months, from May to September, pupping will not occur and subnivean lairs will not be occupied. Since icebreaking would occur when sea ice is at its lowest extent icebreaking areas would not likely overlap with subnivean lairs. However, Williams et al. (2006) determined that ringed seals abandoned subnivean lairs in areas where there was high ice deformation. Ringed seals typically construct their lairs in landfast ice (ice securely attached to land) that typically extends 13.5 to 21.6 nm (25 to 40 km) offshore (Kovacs and Mellor 1974; Stringer 1974; Wadhams 2000). Although icebreaking could overlap with ringed seal structures, it is likely that the noise of the icebreaking would alert any seal well before the icebreaker reaches the subnivean lair, and similar to a predator flight response, the seal would abandon the lair. Therefore, it is unlikely that icebreaking would cause injury or mortality to a ringed seal or their pup from the physical presence of the icebreaking. A recent study of pinniped response to an approaching icebreaking vessel found that there were fewer flush responses by seals to the icebreaker at distances greater than 600 m, and no flush responses by seals to the icebreaker at distances greater than 800 (Lomac-MacNair et al. 2019b). During this study, bearded and ringed seals flushed at closer distances (average 410 and 440 m, respectively), which may suggest that these seals

are less sensitive to disturbance than other pinnipeds considered (i.e., hooded seals) (Lomac-MacNair et al. 2019b).

Since bowhead whales do not rely on ice for habitat use, impacts from icebreaking are more difficult to assess (Tynan and DeMaster 1997). There is uncertainty of the effect of sea ice loss on polar marine food webs. The Proposed Action would not be reducing the amount of sea ice, but breaking it apart into smaller pieces. Bowhead whales are capable of inhabiting areas of dense ice cover, although during summer and fall (when icebreaking could be occurring) bowhead whales are found in areas with reduced sea ice cover (less than 40-70 percent) (Moore et al. 2000). In a study by Moore and Laidre (2006), their conceptual model suggested reduced sea ice cover would increase prey availability.

Potential Impacts to Marine Mammal Critical Habitat

As described in Section 3.2.2.5.1, the critical habitat for ringed seals includes the following essential features:

- Snow-covered sea ice habitat suitable for the formation and maintenance of subnivean birth lairs used for sheltering pups during whelping and nursing, which is defined as waters 3 m or more in depth (relative to mean low low water) containing areas of seasonal landfast (shorefast) ice or dense, stable pack ice, that have undergone deformation and contain snowdrifts of sufficient depth to form and maintain birth lairs (typically at least 54 cm deep).
- Sea ice habitat suitable as a platform for basking and molting, which is defined as areas containing sea ice of 15 percent or more concentration in waters 3 m or more in depth (relative to mean low low water).
- Primary prey resources to support Arctic ringed seals, which are defined to be small, often schooling, fishes, in particular Arctic cod, saffron cod, and rainbow smelt; and small crustaceans, in particular, shrimps and amphipods.

Critical habitat for polar bears includes the following essential features, relative to sea ice:

- Sea ice habitat located over the continental shelf at depths of 984 ft (300 m) or less. In spring and summer, this habitat follows the northward progression of the ice edge as it retreats northward. In fall, this sea ice habitat follows the southward progression of the ice edge as it advances southward.
- Sea ice within 1 mi (1.6 km) of the mean high tide line of barrier island habitat. Barrier islands are used as migration corridors. Polar bears can move freely between barrier islands by swimming or walking on ice or sand bars, thereby avoiding human disturbance.

Though no critical habitat is designated for bearded seals within the Study Area, they are also strongly associated with sea ice habitat in the Arctic. In winter, individuals generally move south as the pack ice advances into the Bering Sea. In late spring and summer, bearded seals move north as the ice edge recedes into the Chukchi and Beaufort Seas. However, some bearded seals stay near the edge of shorefast ice all winter and do not migrate south. Leads, polynyas, and other openings in the sea ice are important features of bearded seal habitat. Juvenile bearded seals tend to associate with sea ice less than adults and are often found in ice free areas such as bays and estuaries. The distribution of bearded seals appears to be strongly associated with shallow water and high biomass of the benthic prey they feed on. They are limited to feeding depths of less than 492-656 ft (150–200 m).

Only areas of thick, wide concentrations of sea ice would require icebreaking by CGC HEALY. During the warm season, these areas are expected to be at a minimum, which would reduce the impact to sea ice critical habitat. Since icebreaking would only occur in the deep water area it would most likely be outside of polar bear critical habitat. The 2021 median September ice extent was far outside of polar

bear critical habitat. The 1981-2010 average September ice extent did fall in the outer edge of the polar bear critical habitat. Looking at recent trends in ice extent the past 5-years have been below the average and did not overlap the polar bear critical habitat. Polar bears do not appear to be significantly affected by vessel movement. Some polar bears have been observed walking, running, and swimming away from approaching vessels, but these reactions were brief and localized. Other bears have been observed approaching vessels or having no reaction to vessels (Lomac-MacNair et al. 2019a; Richardson et al. 1995b). Additionally, icebreaking may result in the temporary displacement of primary prey resources of polar bears and ringed seals, but these species are expected to return to their normal behaviors shortly after the initial disturbance.

Alternative 1

Icebreaking from both Alternatives 1 and 2 would result in the same potential for effects to marine mammals, in that the same icebreaking vessel (CGC HEALY) would be utilized for both Alternatives. The use of an icebreaking vessel may result in localized changes to sea ice habitat as larger sheets of floating ice are broken down into smaller sizes. However, icebreaking is not expected to significantly alter overall habitat for use by marine mammals. In accordance with the ESA, physical effects from icebreaking may affect, but are not likely to adversely affect, bowhead whales, bearded seals, ringed seals, ringed seal critical habitat, polar bears, and polar bear critical habitat.

In accordance with E.O. 12114, physical impacts from icebreaking associated with Alternative 1 would not result in significant harm to marine mammals.

Alternative 2 (Preferred Alternative)

Icebreaking from both Alternatives 1 and 2 would result in the same potential for effects to marine mammals, in that the same icebreaking vessel (CGC HEALY) would be utilized for both Alternatives. The use of an icebreaking vessel may result in localized changes to sea ice habitat as larger sheets of floating ice are broken down into smaller sizes. However, icebreaking is not expected to significantly alter overall habitat for use by marine mammals. Physical impacts from icebreaking associated with the Proposed Action would not result in reasonably foreseeable takes under the MMPA.

In accordance with the ESA, physical impacts from icebreaking may affect, but are not likely to adversely affect, the ESA-listed bowhead whale, bearded seal, ringed seal, ringed seal critical habitat, polar bear, and polar bear critical habitat. Formal consultation was initiated with NMFS regarding bowhead whale, bearded seal, ringed seal, and ringed seal critical habitat on April 29, 2022. Informal consultation was initiated with USFWS regarding polar bears and polar bear critical habitat on July 7, 2022, and a letter of concurrence was received by the Navy on August 9, 2022.

In accordance with E.O. 12114, physical impacts from icebreaking associated with Alternative 2 would not result in significant harm to marine mammals.

4.3.2.5.5 Vessel and In-Water Device Strike

Potential Impacts

Interactions between surface vessels and marine mammals have demonstrated that surface vessels represent a source of acute and chronic disturbance for marine mammals (Au et al. 2000; Bejder et al. 2006; Hewitt 1985; Jefferson et al. 2009; Kraus et al. 1986; Magalhães et al. 2002; Nowacek et al. 2004; Richter et al. 2003; Richter et al. 2008; Williams et al. 2009). Studies have established that cetaceans generally engage in avoidance behavior when surface vessels move toward them. In some circumstances, marine mammals respond to vessels with the same behavioral repertoire and tactics they employ when they encounter predators, although it is not clear what environmental cues marine

mammals might respond to—the sound of water being displaced by the ships, the sound of the ships’ engines, or a combination of environmental cues surface vessels produce while they transit.

Vessel collisions are a well-known source of mortality in marine mammals, and can be a significant factor affecting some large whale populations (Knowlton and Kraus 2001; Laist et al. 2001; van Waerebeek et al. 2007). Bowhead whales often begin avoiding vessels from more than 2.2 nm (4 km) away (Richardson et al. 1995b). Avoidance by this species usually entails altered headings, faster swimming speeds, and shorter amounts of time spent surfacing. Bowhead whales are more tolerant of vessels moving slowly or moving in directions other than towards them. In most studies, observers noted bowhead whales exhibiting avoidance within 1,640 ft (500 m) of vessels, though avoidance at further distances was not able to be judged by observers on vessels (Richardson et al. 1995b). In compiling records of vessel strikes to marine species, Schoeman et al. (2020) identified bowhead whale strikes as rare.

During a review of data on the subject, Laist et al. (2001) compiled historical records of ship strikes, which contained 58 anecdotal accounts. It was noted that in the majority of cases, the whale was either not observed or seen too late to maneuver in an attempt to avoid collision. In the 2020 stranding summary report, only one fin whale and two humpback whales were confirmed strandings from a ship strike (out of a total of 239 marine mammal strandings), neither of which are found within the Study Area (Savage 2021). The most vulnerable marine mammals to collision are thought to be those that spend extended periods at the surface or species whose unresponsiveness to vessel sound makes them more susceptible to vessel collisions (Gerstein 2002; Laist and Shaw 2006; Nowacek et al. 2004). Marine mammals such as dolphins, porpoises, and pinnipeds that can move quickly throughout the water column do not appear to be as susceptible to vessel strikes, though the risk of a strike still exists for these species. Schoeman et al. (2020) recently called into question whether the lower numbers of reported small animal strikes, including dolphins and seals, was actually due to fewer strikes or rather reporting bias favoring reporting of large whale strikes. They specifically noted strikes to be noticeable locally (i.e., more than three strikes in a single location but not known as the most common cause of mortality in the location) for beluga and gray whales, although the locations examined had more frequent vessel traffic than the Study Area.

Few authors have specifically described the responses of pinnipeds to vessels, and most of the available information on reactions to boats concerns pinnipeds hauled out on land or ice. Reactions include a wide spectrum of effects from avoidance and alert, to cases where animals in the water are attracted, and cases on land where there is lack of significant reaction suggesting habituation to or tolerance of vessels (Richardson et al. 1995b). No information is available on potential responses to in-water devices. Brueggeman et al. (1992) stated ringed seals hauled out on the ice showed short-term escape reactions when they were within 0.13 to 0.27 nm (0.25 to 0.5 km) of a vessel. A review of seal stranding data from Alaska found that in 2020, within the Arctic region of Alaska, 7 ringed seal, 2 bearded seal, 4 spotted, and 3 unknown pinniped strandings were recorded. Of the 239 marine mammal strandings reported in all regions of Alaska, there were no pinniped strandings caused by vessel collisions (Savage 2021). From the limited data available, it appears that pinnipeds are not as susceptible to vessel strikes as other marine mammal species. This may be due, at least in part, to the large amount of time they spend on ice (especially when resting and breeding) and their high maneuverability in the water.

Polar bears do not appear to be significantly affected by vessel moment. Some polar bears have been observed walking, running, and swimming away from approaching vessels, but these reactions were brief and localized. Other bears have been observed approaching vessels or having no reaction to vessels (Lomac-MacNair et al. 2019a; Richardson et al. 1995b). Strike of a polar bear is not expected.

The speed of the ship is an important factor in predicting the lethality of a strike. Laist et al. (2001) noted that most severe and fatal injuries to marine mammals occurred when the vessel was traveling in excess of 14 knots, and there were no recorded mortalities at speeds less than 10 knots. Although the maximum speed of the vessels associated with the Proposed Action is 12 knots for the R/V Sikuliaq, and 17 knots for CGC HEALY, these vessels are expected to operate at much slower speeds (below 10 knots) during most of the Proposed Action. However, slow speed does not eliminate the chance that a collision would result in fatal injury. Vanderlaan and Taggart (2007) concluded that at speeds below 8 knots, there was still a 20 percent risk of death from blunt trauma.

Alternative 1

Vessel and in-water device strike from both Alternatives 1 and 2 would result in the same potential for effects to marine mammals, in that the same vessels and in-water devices would be utilized for both Alternatives. The probability of a vessel or in-water devices encountering a marine mammal is expected to be low, which decreases the likelihood of vessels striking marine mammals. Any behavioral avoidance displayed, if a marine mammal were to encounter the vessels or in-water device, is expected to be short-term and inconsequential. Behavioral avoidance would not result in any reactions expected to significantly disrupt behavioral patterns such as migration, breathing, nursing, breeding, feeding and sheltering to a point where the behavior pattern is abandoned or significantly altered or result in reasonably foreseeable takes of marine mammals. Direct vessel or in-water device strikes could result in injury or fatal injury to marine mammals. However, vessel and in-water device strikes are unlikely given the slow vessel speeds (under 12 knots for vessels and 0.5 knots for in-water devices), therefore vessel strike associated with the Proposed Action would not result in significant harm to marine mammals. In accordance with the ESA, vessel and in-water device strike associated with the Proposed Action may affect, but is not likely to adversely affect, bowhead whale, bearded seal, and ringed seal. The Proposed Action would not result in the destruction or adverse modification of critical habitat for ringed seals or polar bears. Strike of a polar bear by a vessel or in-water device is unlikely; effects of this stressor would be discountable. Therefore, there would be no effect to polar bears associated with vessel and in-water device strike from the Proposed Action.

In accordance with E.O. 12114, vessel and in-water device strike associated with Alternative 1 would not result in significant harm to marine mammals.

Alternative 2 (Preferred Alternative)

Vessel and in-water device strike from both Alternatives 1 and 2 would result in the same potential for effects to marine mammals, in that the same vessels and in-water devices would be utilized for both Alternatives. The probability of a vessel or in-water devices encountering a marine mammal is expected to be low, which decreases the likelihood of vessels striking marine mammals. Any behavioral avoidance displayed, if marine mammals were to encounter the vessels or in-water device, is expected to be short-term and inconsequential. Behavioral avoidance would not result in any reactions expected to significantly disrupt behavioral patterns such as migration, breathing, nursing, breeding, feeding and sheltering to a point where the behavior pattern is abandoned or significantly altered or result in reasonably foreseeable takes of marine mammals. Direct vessel or in-water device strikes could result in injury or fatal injury to marine mammals. However, vessel and in-water device strikes are unlikely given the slow vessel speeds (under 12 knots for vessels and 0.5 knots for in-water devices), therefore vessel strike associated with the Proposed Action would not result in significant harm to marine mammals.

Additionally, vessel and in-water device strike associated with the Proposed Action would not result in any reasonably foreseeable takes under the MMPA. In accordance with the ESA, vessel and in-water device strike associated with the Proposed Action may affect, but is not likely to adversely affect,

bowhead whale, bearded seal, and ringed seal. The Proposed Action would not result in the destruction or adverse modification of critical habitat for ringed seals or polar bears. Strike of a polar bear by a vessel or in-water device is unlikely; effects of this stressor would be discountable. Therefore, there would be no effect to polar bears associated with vessel and in-water device strike from the Proposed Action. Formal consultation was initiated with NMFS regarding bowhead whale, bearded seal, and ringed seal on April 29, 2022.

In accordance with E.O. 12114, vessel and in-water device strike associated with Alternative 2 would not result in significant harm to marine mammals.

4.3.2.5.6 Entanglement

Potential Impacts

Devices that pose an entanglement risk are those with lines or tethers; devices with a potential for entanglement include moored or ice-tethered sensors, and lowered devices from the R/V Sikuliaq or CGC HEALY. All lines hanging from buoys or ice tethered equipment would be weighted, and therefore would not have any loops or slack.

The final line that could be a threat for entanglement is the use of a device tethered to an unmanned underwater vehicle (depth of 295 ft [91 m]). The tether for this research initiative has a diameter of 8.9 mm, and is made of Kevlar. This tether has a very high breaking strength (1,543 lb force [700 kg force]), but environmental resources should not be at risk due to the small likelihood of any loops or slack developing in this line, since it would be under positive pressure. No mooring lines would be expended during the Proposed Action, so this further limits the chance for entanglement.

The likelihood of a marine mammal encountering and becoming entangled in a line depends on several factors. The amount of time that the line is in the same vicinity as a marine mammal can increase the likelihood of it posing an entanglement risk. The length of the line varies (up to approximately 12,303 ft [3,750 m]) and greater lengths may increase the likelihood that a marine mammal could become entangled. The behavior and feeding strategy of a species can determine whether they may encounter items on the seafloor. Given the water depths, marine mammals would not forage on the seafloor within the Study Area, eliminating the possibility of entanglement with the bottom mounted acoustic sources.

During the deployment and removal of the lines and buoys, marine mammals could become entangled. However, all equipment would be deployed from a shipboard winch system in a slow and controlled manner, which would decrease the potential of entanglement. Additionally, the lines are weighted to help with deployment, this would make the line free of loops and slack for marine mammals to become entangled.

Once the moorings and anchors are in place the potential for entanglement with tethered moored equipment is considered negligible based on the tension in the line, small buoy sizes (51 in [130 cm] diameter), shape depth (approximately 656 ft [200 m] or on the seafloor), and the large spacing between shapes (minimum of 40.5 nm [75 km]). Bearded seals and ribbon seals may dive up to 656 ft (200 m) underwater; however, both species are expected to be closer to shore than the Study Area. Bowhead whales may dive to depths greater than 1,148 ft (350 m) and may encounter expended materials. However, there would be no slack in the mooring tethers which are under approximately 1,190 lb (540 kg) of tension due to the shape buoyancy. The probability of a whale, such as a bowhead, colliding with a moored shape is considered remote. Pinnipeds are highly maneuverable and could easily avoid bottom or tethered shapes and most pinnipeds (bearded seal, ribbon seal, spotted seal) would not be found over 75.6 nm (140 km) from the shore. Moorings will not have a surface expression and the

buoy which keeps the line taught would be approximately 164 ft (50 m) below the surface of the ice, negating the chance for a seal to become entangled while utilizing a breathing hole. Based on the estimated concentration of deployed mooring lines, impacts from lines are extremely unlikely to occur. Although there is a potential for entanglement from an expended material the amount of materials expended would be low. Ringed seals are very mobile within the water column and avoidance of any expended object is expected.

The chance that an individual animal would encounter expended lines is most likely low based on the distribution of both the lines expended, and the depth of the water in the Study Area where these would be expended. In the 2020 NMFS stranding report, 7 reported ringed seal, 6 gray whale, 4 spotted seal, 2 bearded seal, 3 unidentified pinniped, and 5 unidentified cetacean strandings occurred in the Arctic region of Alaska. There were no confirmed bowhead or beluga strandings reported in the region in 2020. Of the 239 total strandings reported throughout Alaska in 2020, only 31 were documented to be from entanglement. Of these 31 entanglement-related strandings, the only ones of a species expected to occur within the Study Area included one gray whale and two ringed seals (Savage 2021). Given the water depths in the Study Area, marine mammals are not expected to be feeding on the seafloor; any materials that settle to the seafloor would therefore not pose an entanglement risk to marine mammals. An animal would have to swim through loops or become twisted within the lines to become entangled. Based on the limited number of expended lines, harm from lines is extremely unlikely to occur. Although there is a potential for entanglement from an expended material the amount of materials expended would be low. Marine mammals are very mobile within the water column and avoidance of any expended object is expected.

The potential for entanglement would be from mooring lines and towed sources. Bowhead whales' average dive depth is 328 ft (100 m), with maximum recorded dive of 1,155 ft (352 m) where it could encounter expended materials (Krutzikowski and Mate 2000). However, all lines extending from the moorings would be retrieved at the completion of the Proposed Action. Any effects to bowhead whales would not be significant and any reactions are not expected to significantly disrupt behavioral patterns such as migration, breathing, nursing, breeding, feeding and sheltering to a point where the behavior pattern is abandoned or significantly altered or result in reasonably foreseeable takes.

Polar bears are normally found in locations of 50 percent ice cover and at water depths of 984 ft (300 m) within the Beaufort Sea and are not expected to occur in the deep waters of the Study Area during summer months when equipment would be deployed and retrieved. Polar bears would not be foraging or diving to the seafloor at either Study Area. Therefore, the potential of a polar bear becoming entangled in expended materials associated with the Proposed Action is considered negligible.

Alternative 1

Under Alternative 1, the potential for entanglement would be from mooring lines. Alternative 1 has fewer mooring lines associated with the Proposed Action due to the exclusion of the VLF sources. All lines extending from the moorings would be retrieved at the completion of the Proposed Action. Any effects to marine mammals would not be significant and any reactions are not expected to significantly disrupt behavioral patterns such as migration, breathing, nursing, breeding, feeding and sheltering to a point where the behavior pattern is abandoned or significantly altered or result in reasonably foreseeable takes of marine mammals. In accordance with the ESA, entanglement associated with Alternative 1 may affect, but is not likely to adversely affect, the ESA-listed bowhead whale or ringed seal. Entanglement would have no effect on polar bears or bearded seals.

Therefore, in accordance with E.O. 12114, entanglement associated with Alternative 1 would not result in significant harm to mammals.

Alternative 2 (Preferred Alternative)

Under Alternative 2, the potential for entanglement would be from mooring lines. Any effects to marine mammals would not be significant and any reactions are not expected to significantly disrupt behavioral patterns such as migration, breathing, nursing, breeding, feeding and sheltering to a point where the behavior pattern is abandoned or significantly altered or result in reasonably foreseeable takes of marine mammals. Therefore, in accordance with E.O. 12114, entanglement associated with Alternative 2 would not result in significant harm to mammals.

Entanglement associated with Alternative 2 would not result in any reasonably foreseeable takes under the MMPA. Entanglement associated with Alternative 2 would have no effect on the ESA-listed polar bear or bearded seal because there would be no overlap of those species with entanglement stressors. Entanglement associated with Alternative 2 may affect, but is not likely to adversely affect, the ESA-listed bowhead whale or ringed seal. Formal consultation was initiated with NMFS regarding bowhead whale and ringed seal on April 29, 2022.

4.3.2.5.7 Ingestion

Potential Impacts

During the Proposed Action, the only expended materials available for ingestion include the on-ice measurement systems. On-ice measurement systems include the autonomous weather station and the ice mass balance buoy. The autonomous weather station would be deployed on a tripod with insulated foot platforms frozen into the ice. While the ice mass balance buoy would be lowered into the water column through a two-inch hole in the ice, there would be a tripod located on the ice. All other expended objects would be expended into the water column and would sink to the seafloor. Ingestion stressors are not anticipated to affect any resources other than polar bears, due to the large size of the material that is expended in the water column or stationed on the sea ice. These objects (e.g., anchors, buoys) would be too large for any marine mammals to eat. The objects deployed and expended within the water column would be too deep to overlap with a swimming polar bear.

Polar bears typically find alternate food sources (e.g., land-based trash collection sites) when their primary prey (ringed seals) are unavailable (Lunn and Stirling 1985). In a study by Gormezano and Rockwell (2013), polar bear scats (i.e., excrement) from five sites were surveyed. Sites included the town of Churchill and dens around inland lakes. In areas where humans and polar bears came in close proximity, a higher percentage of garbage was found in the scats than areas where polar bears and humans were not in close proximity. Polar bears have also been known to bite buoys located on the ice. This behavior could be out of curiosity or to determine if the object is edible. The likelihood of a polar bear encountering the autonomous weather station tripod from the ice mass balance buoy on the ice, and potentially taking a bite of the equipment is low since a small number of on-ice measurement systems would be deployed on ice floes in the Study Area. Although ingestion of large pieces of the autonomous weather station or tripod from the ice mass balance buoy is not anticipated, small bits could be ingested. If a polar bear does ingest pieces of the autonomous weather station or tripod from the ice mass balance buoy, the bear would likely excrete the material without detrimental effects, as studies indicate that bears foraging in land-based trash sites show no reproductive or survival advantage or disadvantage from feeding on these materials (Lunn and Stirling 1985).

Alternative 1

Under both Alternatives 1 and 2, the potential for ingestion would be limited to exploratory bites of the autonomous weather station or tripod from the ice mass balance buoy on the ice by polar bears. In accordance with the ESA, ingestion would have no effect on bowhead whales, bearded seals, or ringed

seals. In accordance with E.O. 12114 ingestion of materials associated with Alternative 1 would not result in significant harm to marine mammals.

Alternative 2 (Preferred Alternative)

Under both Alternatives 1 and 2, the potential for ingestion would be limited to exploratory bites of the autonomous weather station or tripod from the ice mass balance buoy on the ice by polar bears. Any effects to marine mammals would be minimal and temporary, and therefore would not result in reasonably foreseeable takes under the MMPA.

In accordance with the ESA, Alternative 2 would have no effect on ringed seals, bowhead whales, or bearded seals. Polar bears, however, may be attracted to the autonomous weather station or tripod from the ice mass balance buoy; therefore, ingestion associated with the Proposed Action under Alternative 2 may affect, but is not likely to adversely affect, polar bears. Informal consultation was initiated with USFWS regarding polar bears on July 7, 2022, and a letter of concurrence was received by the Navy on August 9, 2022.

In accordance with E.O. 12114 ingestion of materials associated with Alternative 2 would not result in significant harm to marine mammals.

4.4 Summary of Potential Impacts to Resources

A summary of the potential impacts associated with each of the Action Alternatives and the No Action Alternative are presented in Table 4-6.

Table 4-6. Summary of Potential Impacts to Resource Areas

<i>Resource Area</i>	<i>No Action Alternative</i>	<i>Alternative 1</i>	<i>Alternative 2 (Preferred Alternative)</i>
Physical Resources	No change to baseline.	The potential harm would be temporary and localized due to the minimal number of devices and the infrequency of testing activities, and soft sediment is expected to shift back as it would following a disturbance of tidal energy. No long-term increases in turbidity (sediment suspended in water) would be anticipated. The localized disturbances would not alter the function or habitat provided by marine substrates or sea ice. No significant harm due to changes in ambient noise levels would occur as a result of the Proposed Action.	
Invertebrates	No change to baseline.	With standard operating procedures, potential harm from the Proposed Action would be temporary and/or minimal. The Proposed Action is not expected to result in population-level impacts to invertebrates.	
Marine Birds	No change to baseline.	With standard operating procedures, potential harm from the Proposed Action would not be expected. The Proposed Action is not expected to result in population-level impacts to marine birds.	
Fish	No change to baseline.	With standard operating procedures, potential harm from the Proposed Action would be temporary and/or minimal. The Proposed Action is not expected to result in population-level impacts to fish.	
Essential Fish Habitat	No change to baseline.	With standard operating procedures, potential adverse effects from the Proposed Action would be considered minimal.	
Marine Mammals	No change to baseline.	With standard operating procedures and mitigation measures, potential harm from the Proposed Action would be temporary and/or minimal. The Proposed Action is not expected to result in population-level impacts to marine mammals.	

5 Standard Operating Procedures and Mitigation Measures

Both standard operating procedures and mitigation measures would be implemented during the Proposed Action. Standard operating procedures serve the primary purpose of providing safety and mission success, and are implemented regardless of their secondary benefits (e.g., to a resource), while mitigation measures are used to avoid or reduce potential impacts.

Moored/drifted sources are left in place and cannot be turned off until the following year during ice free months. Once they are programmed they will operate at the specified pulse lengths and duty cycles until they are either turned off the following year or there is failure of the battery and the devices are not able to operate. Due to the ice covered nature of the Arctic it is not possible to recover the sources or interfere with their transmit operations when research cruises are not occurring.

While underway the ships (including non-Navy ships operating on behalf of the Navy) utilizing active acoustic sources will have at least one watch person during activities. While underway, watch personnel are alert at all times and have access to binoculars.

5.1 Standard Operating Procedures

- Ships operated by or for the U.S. Navy, including those associated with the Proposed Action, would have personnel assigned to stand watch at all times, day and night, when moving through the water (underway) or using active acoustic sources.
- While on watch, personnel employ visual search techniques, including the use of binoculars, using a scanning method in accordance with the Marine Species Awareness Training (MSAT). A primary duty of watch personnel is to detect and report all objects and disturbances sighted in the water that may be indicative of a threat to the ship and its crew, such as debris, or surface disturbance. Per safety requirements, watch personnel also report any marine mammals sighted that have the potential to be in the direct path of the ship as a standard collision avoidance procedure.
- While in transit, ships shall be alert at all times, use extreme caution, and proceed at a "safe speed" so that the ship can take proper and effective action to avoid a collision with any marine mammal and can be stopped within a distance appropriate to the prevailing circumstances and conditions.

5.2 Mitigation Measures

- Watch personnel undertake MSAT, which trains Navy lookouts in proper methods for visual observation for the presence of marine species within mitigation zones.
- Mitigation zones for active acoustics involve turning off vessel-bound sources when a marine mammal is sighted within 200 yards (yd; 183 m) from the source. Active transmission will recommence if any one of the following conditions are met: (1) the animal is observed exiting the mitigation zone, (2) the animal is thought to have exited the mitigation zone based on its course and speed and relative motion between the animal and the source, (3) the mitigation zone has been clear from any additional sightings for a period of 30 minutes, (4) the vessel has transited more than 400 yd (366 m) beyond the location of the last sighting.
- During mooring deployment visual observation would start 15 minutes prior to and during the deployment within a mitigation zone of 180 ft (55 m) around the deployed mooring. Deployment will stop if a marine mammal is visually detected within the mitigation zone. Deployment will recommence if any one of the following conditions are met: (1) the animal is observed exiting the mitigation zone, (2) the animal is thought to have exited the mitigation zone based on its course and

speed, or (3) the mitigation zone has been clear from any additional sightings for a period of 15 minutes.

- Ships would avoid approaching marine mammals head on and would maneuver to maintain a mitigation zone of 500 yd (457 m) around observed whales, and 200 yd (183 m) around all other marine mammals, providing it is safe to do so during ice free waters.
- These requirements do not apply if a vessel's safety is at risk, such as when a change of course would create an imminent and serious threat to safety, person, vessel, or aircraft, and to the extent vessels are restricted in their ability to maneuver. No further action is necessary if a marine mammal other than a whale continues to close on the vessel after there has already been one maneuver and/or speed change to avoid the animal. Avoidance measures should continue for any observed whale in order to maintain a mitigation zone of 500 yd (457 m).

5.3 Monitoring and Reporting

There are no specific monitoring plans outside of MSAT-trained lookouts aboard the CGC HEALY and R/V Sikuliaq. Due to the harsh conditions in the Arctic Study Area it is not feasible to tag and monitor marine mammals as it would require additional personnel and equipment. Any marine mammal sightings would be communicated to NMFS in the IHA after-action report.

While there is no monitoring specific to the Proposed Action, the ONR Marine Mammal Biology Program has funded research in Alaska on ice seals and whales. Currently ONR has funded a study to work with Native subsistence hunters and government agencies in Alaska (North Slope Borough Department of Wildlife Management) and Canada (Department of Fisheries and Oceans) to deploy satellite tags on ringed seals, spotted seals, bearded seals, bowhead whales, and beluga whales. The research is aimed to document year-round movements of each species and document habitat use relative to oceanographic conditions, ice cover, and human disturbance.

The Navy is committed to documenting and reporting relevant aspects of training and research activities to verify implementation of mitigation, comply with current permits, and improve future environmental assessments. If any injury or death of a marine mammal is observed during the 2022-2025 Arctic Research Activities, the Navy will immediately halt the activity and report the incident consistent with the stranding and reporting protocol in other Navy documents such as the Atlantic Fleet Training and Testing Environmental Impact Statement/Overseas Environmental Impact Statement.

6 Consistency with Other Federal, State, and Local Laws, Plans, Policies, and Regulations

In accordance with 40 CFR section 1502.16(c), analysis of environmental consequences shall include discussion of possible conflicts between the Proposed Action and the objectives of federal, regional, state and local land use plans, policies, and controls. Table 6-1 identifies the principal federal and state laws and regulations that are applicable to the Proposed Action, and describes briefly how compliance with these laws and regulations would be accomplished.

Table 6-1. Principal Federal and State Laws Applicable to the Proposed Action

<i>Federal, State, Local, and Regional Land Use Plans, Policies, and Controls</i>	<i>Status of Compliance</i>
Arctic Research and Policy Act	This OEA has been prepared in compliance with the goals of the Arctic Research Policy Act.
Endangered Species Act (16 U.S.C. section 1531 et seq.)	<p>This OEA considers impacts on species listed as threatened or endangered pursuant to this act.</p> <p>In accordance with the ESA, consultation with NMFS and USFWS was initiated based on the determination that the Proposed Action may affect, but is not likely to adversely affect, bowhead whales (<i>Balaena mysticetus</i>), bearded seals (<i>Erignathus barbatus</i>), and polar bears (<i>Ursus maritimus</i>). The Proposed Action may affect, and is likely to adversely affect, ringed seals (<i>Phoca hispida</i>). The Proposed Action is not likely to adversely affect critical habitat for ringed seals or polar bears. Formal consultation was initiated with NMFS regarding bowhead whales, bearded seals, and ringed seals on April 29, 2022. Informal consultation was initiated with USFWS regarding polar bears on July 7, 2022, and a letter of concurrence was received by the Navy on August 9, 2022.</p>
Marine Mammal Protection Act (16 U.S.C. section 1361 et seq.)	This OEA considers impacts on protected marine mammal species pursuant to this act. Based on the analysis contained within this OEA, the Navy submitted an application for an IHA with NMFS for the taking of beluga whales and ringed seals on March 21, 2022. A Notice of Availability of the draft IHA was published in the Federal Register on July 26, 2022.
Migratory Bird Treaty Act (16 U.S.C. sections 703-712)	This OEA considers impacts on migratory birds under this Act. Implementation of the Proposed Action would have no effect on migratory bird populations.
Magnuson-Stevens Fishery Conservation and Management Reauthorization Act (16 U.S.C. section 1801 et seq.)	This OEA considers impacts on fish and wildlife and essential fish habitat under this act. Based on the analysis contained within this OEA, the Navy submitted an Essential Fish Habitat Assessment with NMFS regarding potential impacts on EFH designated for Arctic cod on July 6, 2022. Concurrence was received by the Navy from NMFS on July 29, 2022.
Executive Order 12114, Environmental Effects Abroad of Major Federal Actions	This OEA has been prepared in accordance with E.O. 12114 and Navy E.O. 12114 procedures.

7 References

- Aerts, L. A., McFarland, A. E., Watts, B. H., Lomac-MacNair, K. S., Seiser, P. E., Wisdom, S. S., Kirk, A. V., & Schudel, C. A. (2013). Marine mammal distribution and abundance in an offshore sub-region of the northeastern Chukchi Sea during the open-water season. *Continental Shelf Research*, 67, 166-126.
- Alaska Fisheries Science Center Marine Mammal Laboratory. (2014). 2014 Aerial surveys of arctic marine mammals Retrieved from https://www.afsc.noaa.gov/nmml/cetacean/bwasp/flights_2014.php as accessed on May 2017.
- Allen, B. M., & Angliss, R. P. (2011). *Alaska marine mammal stock assessments, 2010*. (NOAA Technical Memorandum NMFS-AFSC-223). Seattle, WA. p. 292.
- Allen, B. M., & Angliss, R. P. (2014). *Alaska marine mammal stock assessments, 2013*. (NOAA Technical Memorandum NMFS-AFSC-277). Seattle, WA. p. 294.
- Alliston, W. G. (1980). *The distribution of ringed seals in relation to winter icebreaking activities near McKinley Bay, NWT, January-June 1980*. Toronto, Ont.: Dome Petroleum. p. 52.
- Alliston, W. G. (1981). *The distribution of ringed seals in relation to winter icebreaking activities in Lake Melville, Labrador*. LGL Limited environmental research associates.
- Amoser, S., & Ladich, F. (2003). Diversity in noise-induced temporary hearing loss in otophysine fishes. *Journal of the Acoustical Society of America*, 113(4), 2170-2179.
- Amoser, S., & Ladich, F. (2005). Are hearing sensitivities of freshwater fish adapted to the ambient noise in their habitats? *The Journal of Experimental Biology*, 208, 3533-3542.
- Amstrup, S. C., Durner, G. M., Stirling, I., Lunn, N. J., & Messier, F. (2000). Movements and distribution of polar bears in the Beaufort Sea. *Canadian Journal of Zoology*, 78, 948-966.
- Amstrup, S. C., & Gardner, C. (1994). Polar bear maternity denning in the Beaufort Sea. *The Journal of Wildlife Management*, 58(1), 1-10.
- Ankley, G. T., Di Toro, D. M., Hansen, D. J., & Berry, W. J. (1996). Technical basis and proposal for deriving sediment quality criteria for metals. *Environmental Toxicology and Chemistry*, 15(12), 2056-2066.
- Appeltans, W., Bouchet, P., Boxshall, G. A., Fauchald, K., Gordon, D. P., Hoeksema, B. W., Poore, G. C. B., van Soest, R. W. M., Stöhr, S., Walter, T. C., & Costello, M. J. (2010). *World Register of Marine Species* (Vol. 2010). Retrieved from <http://www.marinespecies.org/index.php>.
- Ashjian, C. J., Braund, S. R., Campbell, R. G., George, J. C., Kruse, J., Maslowski, W., Moore, S. E., Nicolson, C. R., Okkonen, S. R., Sherr, B. F., Sherr, E. B., & Spitz, Y. H. (2010). Climate variability, oceanography, bowhead whale distribution, and Iñupiat subsistence whaling near Barrow, Alaska. 63(2), 179-194.
- Astrup, J. (1999). Ultrasound detection in fish- A parallel to the sonar-mediated detection of bats by ultrasound-sensitive insects. *Comparative Biochemistry and Physiology*, 124, 19-27.
- Astrup, J., & Mohl, B. (1993). Detection of intense ultrasound by the cod (*Gadus morhua*). *The Journal of Experimental Biology*, 182(1), 71-80.
- Au, W. W. L. (1993). *The sonar of dolphins*. New York, NY: Springer.
- Au, W. W. L., Floyd, R. W., Penner, R. H., & Murchison, A. E. (1974). Measurement of echolocation signals of the Atlantic bottlenose dolphin, *Tursiops truncatus* Montagu, in open waters. *Journal of the Acoustical Society of America*, 56(4), 1280-1290.
- Au, W. W. L., Mobley, J., Burgess, W. C., Lammers, M. O., & Nachtigall, P. E. (2000). Seasonal and Diurnal Trends of Chorusing Humpback Whales Wintering in Waters off Western Maui. *Marine Mammal Science*, 16(3), 530-544.

- Ballinger, T. J., Overland, J., Wang, M., Bhatt, U. S., Brettschneider, B., Hanna, E., Hanssen-Bauer, I., Kim, S. J., Thoman, R. L., & Walsh, J. (2021). Surface Air Temperature. In Moon, T. A., Druckenmiller, M. L. & Thoman, R. L. (Eds.), Arctic Report Card 2021. doi: 10.25923/53xd-9k68
- Beason, R. C. (2004). What can birds hear? *USDA National Wildlife Research Center - Staff Publications, Paper 78*, 92-96.
- Becker, A., Whitfield, A., Cowley, K., Järnegren, J., & Næsje, T. F. (2013). Does boat traffic cause displacement of fish in estuaries? *Marine Pollution Bulletin*, 75(1), 168-173.
- Bejder, L., Samuels, A., Whitehead, H., & Gales, N. (2006). Interpreting Short-Term Behavioural Responses to Disturbance within a Longitudinal Perspective. *Animal Behavior*, 72, 1149-1158.
- Belkin, I. M., & Cornillon, P. C. (2007). Fronts in the world ocean's large marine ecosystems. *ICES CM*, 500(130), 21.
- Bengtson, J. L., Boveng, P. L., Hiruki-Raring, L. M., Laidre, K. L., C., P., & Simpkins, M. A. (2000). *Abundance and distribution of ringed seals (Phoca hispida) in the coastal Chukchi Sea*. (AFSC Processed Rep. 2000-11). 7600 Sand Point Way NE, Seattle, WA 98115: Alaska Fisheries Science Center. pp. 149-160.
- Bengtson, J. L., Hiruki-Raring, L. M., Simpkins, M. A., & Boveng, P. L. (2005). Ringed and bearded seal densities in the eastern Chukchi Sea, 1999–2000. *Polar Biology*, 28, 833-845. doi: 10.1007/s00300-005-0009-1.
- Bernasconi, M., Patel, R., & Nøttestad, L. (2012). Behavioral observations of baleen whales in proximity of a modern fishing vessel. In Popper, A. N. & Hawkins, A. D. (Eds.), *The Effects of Noise on Aquatic Life* (pp. 335-338). New York: Springer.
- Bethke, R., Taylor, M., Amstrup, S. C., & Messier, F. (1996). Population delineation of polar bears using satellite collar data. *Ecological Applications*, 6(1), 311-317.
- Beuter, K. J., Weiss, R., & Frankfurt, B. (1986). *Properties of the Auditory System in Birds and the Effectiveness of Acoustic Scaring Signals*. Paper presented at the Bird Strike Committee Europe (BSCE), 18th Meeting Part I, Copenhagen, Denmark.
- Bickel, S. L., Malloy Hammond, J. D., Tang, K. W. B., Samantha L., Malloy Hammond, J. D., & Tang, K. W. (2011). Boat-generated turbulence as a potential source of mortality among copepods. *Journal of Experimental Marine Biology and Ecology*, 401(1-2), 105-109.
- Bigg, M. A. (1981). Harbour Seal *Phoca vitulina* Kinnaeus, 1758 and *Phoca largha* Pallas, 1811. In Ridgway, S. H. & Harrison, R. (Eds.), *Seals* (pp. 345-354). New York: Academic Press.
- BirdLife International. (2012). *Uria lomvia*. The IUCN Red List of Threatened Species. Version 2015.2 Retrieved from <http://www.iucnredlist.org/details/22694847/0> as accessed on 25 June 2015.
- BirdLife International. (2016). Species factsheet: *Pagophilia eburnea* Retrieved from <http://datazone.birdlife.org/species/factsheet/ivory-gull-pagophila-eburnea/text> as accessed on 7 November 2016.
- Bisby, F. A., Roskov, Y. R., Orrell, T. M., Nicolson, D., Paglinawan, L. E., Bailly, N., & Baillargeon, G. (2014). Species 2000 & ITIS catalogue of life: 2014 annual checklist Retrieved from <http://www.catalogueoflife.org/annual-checklist/2014/browse/tree> as accessed on 5 August 2014.
- Bishop, M. J. B., M. J. (2008). Displacement of epifauna from seagrass blades by boat wake. *Journal of Experimental Marine Biology and Ecology*, 354(1), 111-118.
- Blackwell, S. B., Lawson, J. W., & Williams, M. T. (2004). Tolerance by ringed seals (*Phoca hispida*) to impact pipe-driving and construction sounds at an oil production island. *The Journal of the Acoustical Society of America*, 115(5), 2346-2357.
- Bleckmann, H., & Zelick, R. (2009). Lateral line system of fish. *Integrative Zoology*, 4(1), 13-25.
- Bluhm, B. (2008, January 20, 2011). Sea Bottom. *Arctic Ocean Diversity: Species* Retrieved Retrieved January 20, 2011 from <http://www.arcodiv.org/overview.html> as accessed on June 22.

- Bluhm, B., Gradinger, R., & Schnack-Shiel, S. (2010). Sea ice meio and macrofauna. In Thomas, D. & Diechmann, D. (Eds.), *Sea Ice* (pp. 357-393). New York: Wiley-Blackwell.
- Bluhm, B. A., Gebruk, A. V., Gradinger, R., Hopcroft, R. R., Heuttmann, F., Kosobokova, K. N., Sirenko, B. I., & Weslawski, J. M. (2011). Arctic marine biodiversity: An update of species richness and examples of biodiversity change. *Oceanography*, 24(3), 232-248. doi: <http://dx.doi.org/10.5670/oceanog.2011.75>.
- Bluhm, B. A., & Gradinger, R. (2008). Regional Variability in Food Availability for Arctic Marine Mammals. *Ecological Applications*, 18(2), S77-S96.
- Bluhm, B. A., MacDonald, I. R., Debenham, C., & Iken, K. (2005). Macro- and megabenthic communities in the high Arctic Canada Basin: initial findings. *Polar Biology*, 28, 218-231. doi: DOI 10.1007/s00300-004-0675-4.
- Bonner, W. N. (1986). *Marine mammals of the Falkland Islands*. Cambridge, UK: British Antarctic Survey.
- Born, E. W., Teilmann, J., Acquarone, M., & Riget, F. F. (2004). Habitat use of ringed seals (*Phoca hispida*) in the North Water area (North Baffin Bay). *Arctic*, 57(2), 129-142.
- Boveng, P., & Cameron, M. F. (2013). *Pinniped movements and foraging: seasonal movements, habitat selection, foraging and haul-out behavior of adult bearded seals in the Chukchi Sea. Final Report*. Anchorage, AK: Bureau of Ocean Energy Management, Alaska Outer Continental Shelf Region, p. 91.
- Boveng, P. L., Bengtson, J. L., Buckley, T. W., Cameron, M. F., Dahle, S. P., Megrey, B. A., Overland, J. E., & Williamson, N. J. (2008). *Status review of the ribbon seal (Histriophoca fasciata)*. (NOAA Tech. Memo. NMFS-AFSC-191). p. 115.
- Bowles, A. E., Owen, M. A., Denes, S. L., Graves, S. K., & Keating, J. L. (2008). Preliminary results of a behavioral audiometric study of the polar bear. *The Journal of the Acoustical Society of America*, 123(5), 3509-3509.
- Braham, H. W., Burns, J. J., Fedoseev, G. A., & Krogman, B. D. (1981). *Distribution and density of ice-associated pinnipeds in the Bering Sea*. National Marine Mammal Laboratory (NMML).
- Braham, H. W., Burns, J. J., Feoseev, G. A., Krogman, B. D., & Gennadii, A. (1984). *Habitat partitioning by ice-associated pinnipeds: Distribution and density of seals and walruses in the Bering Sea, April 1976*. National Oceanic and Atmospheric Administration (NOAA). pp. 25-48.
- Braham, H. W., Fraker, M. A., & Krogman, B. D. (1980). Spring migration of the western Arctic population of bowhead whales. *Marine Fisheries Review*, 42(9-10), 36-46.
- Brooke, M. d. L. (2001). Seabird Systematics and Distribution: A Review of Current Knowledge. In Schreiber, E. A. & Burger, J. (Eds.), *Biology of Marine Birds* (pp. 57-85). Boca Raton, FL: CRC Press LLC.
- Brueggeman, J., Green, G., Grotefendt, R., Smultea, M., Volsen, D., Rowlett, R., Swanson, C., Malme, C., Mlawski, R., & Burns, J. (1992). Marine Mammal Monitoring Program (Seals and Whales) Crackerjack and Diamond Prospects Chukchi Sea. *Rep. from EBASCO Environmental, Bellevue, WA, for Shell Western E&P Inc. and Chevron USA Inc*, 62.
- Brueggeman, J. J., Volsen, D. P., Grotefendt, R. A., Green, G. A., Burns, J. J., & Ljungblad, D. K. (1991). *1990 Walrus Monitoring Program: The Popcorn, Burger, and Crackerjack Prospects in the Chukchi Sea*. Fairbanks, AK: Living Resources Inc.
- Budelmann, B. U. (1992a). Hearing by crustacea. In *Evolutionary Biology of Hearing* (pp. 131-139). New York: Springer-Verlag.
- Budelmann, B. U. (1992b). Hearing in nonarthropod invertebrates. In *Evolutionary Biology of Hearing* (pp. 16). New York: Springer-Verlag.
- Budelmann, B. U. (2010). *Cephalopoda*. Oxford, UK: Wiley-Blackwell.
- Budge, S. M., Springer, A. M., Iverson, S. J., Sheffield, G., & Rosa, C. (2008). Blubber fatty acid composition of bowhead whales, *Balaena mysticetus*: Implications for diet assessment and

- ecosystem monitoring. *Journal of Experimental Marine Biology and Ecology*, 359, 40-46. doi: doi: 10.1016/j.jembe.2008.02.014.
- Buran, B. N., Deng, X., & Popper, A. N. (2005). Structural variation in the inner ears of four deep-sea elopomorph fishes. *Journal of Morphology*.
- Burns, J. J. (1970). Remarks on the distribution and natural history of pagophilic pinnipeds in the Bering and Chukchi seas. *Journal of Mammalogy*, 51(3), 445-454.
- Burns, J. J. (1981a). Bearded seal, *Erignathus barbatus* (Erxleben, 1777). In *Handbook of Marine Mammals* (Vol. 2: Seals, pp. 145-170). New York: Academic Press.
- Burns, J. J. (1981b). Ribbon seal-- *Phoca fasciata* (Zimmerman, 1783). In. Ridgway, S. H. & Harrison, R. J. (Eds.), *Seals* (pp. 89-109): Academic Press.
- Burns, J. J., & Frost, K. J. (1979). *The natural history and ecology of the bearded seal, Erignathus barbatus*. Fairbanks, AK: Alaska Department of Fish and Game for the Outer Continental Shelf Environmental Assessment Program. p. 392.
- Burns, J. J., Shapiro, L. H., & Fay, F. H. (1981). Ice as marine mammal habitat in the Bering Sea. In. Hood, D. W. & Calder, J. A. (Eds.), *The eastern Bering Sea shelf: oceanography and resources* (Vol. 2). Juneau, Alaska: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Office of Marine Pollution Assessment.
- Cairns, D. K., & Schneider, D. C. (1990). Hot spots in cold water: feeding habitat selection by thick-billed murre. *Studies in Avian Biology*, 14, 52-60.
- Calambokidis, J., Steiger, G. H., Curtice, C., Harrison, J., Ferguson, M. C., Becker, E. A., DeAngelis, M. L., & Van Parijs, S. M. (2015). Biologically Important Areas for Selected Cetaceans Within U.S. Waters--West Coast Region. *Aquatic Mammals*, 41(1), 39-53.
- Cameron, M. F., & Boveng, P. L. (2009). *Habitat use and seasonal movements of adult and sub-adult bearded seals*. *Alaska Fisheries Science Center Quarterly Report* pp. 1-4.
- Cañadas, A., Roberts, J. J., Schick, R. S., & Halpin, P. N. (2020). *Updated Marine Species Density Models for the Arctic Study Area. Draft Report*. Prepared for Fleet Forces Command.: Submitted to Naval Facilities Engineering Command Atlantic, Norfolk, Virginia.
- Carmack, E. C., & Macdonald, R. W. (2002). Oceanography of the Canadian Shelf of the Beaufort Sea: a setting for marine life. *Arctic*, 29-45.
- Carretta, J. V., Forney, K. A., Lowry, M. S., Barlow, J., Baker, J., Johnston, D., Hanson, B., Muto, M. M., Lynch, D., & Carswell, L. (2008). *U.S. Pacific Marine Mammal Stock Assessments: 2008*. (NOAA-TM-NMFS-SWFSC-434). National Oceanic and Atmospheric Administration. p. 340.
- Carretta, J. V., Oleson, E. M., Forney, K. A., Muto, M. M., Weller, D. W., Lang, A. R., Baker, J., Hanson, B., Orr, A. J., Barlow, J., Moore, J. E., & Brownell Jr, R. L. (2021). *U.S. Pacific Marine Mammal Stock Assessments: 2020*.
- Carroll, A., Przeslawski, R., Duncan, A., Gunning, M., & Bruce, B. (2017). A critical review of the potential impacts of marine seismic surveys on fish & invertebrates. *Marine Pollution Bulletin*, 114(1), 9-24.
- Castellote, M., Mooney, T. A., Quakenbush, L., Hobbs, R., Goertz, C., & Gaglione, E. (2014). Baseline hearing abilities and variability in wild beluga whales (*Delphinapterus leucas*). *The Journal of Experimental Biology*, 217, 1682-1691. doi: 10.1242/jeb.093252.
- Celi, M., Filiciotto, F., Vazzana, M., Arizza, V., Maccarrone, V., Ceraulo, M., Mazzola, S., & Buscaino, G. (2014). Shipping noise affecting immune responses of European spiny lobster (*Palinurus elephas*). *Canadian Journal of Zoology*, 93(2), 113-121.
- Chan, F. T., Stanislawczyk, K., Sneekes, A. C., Dvoretzky, A., Gollasch, S., Minchin, D., David, M., Jelmert, A., Albretsen, J., & Bailey, S. A. (2019). Climate change opens new frontiers for marine species in the Arctic: Current trends and future invasion risks. *Global change biology*, 25(1), 25-38.

- Chapskii, K. K. (1940). *The ringed seal of western seas of the Soviet Arctic (The morphological characteristic, biology and hunting production)*. Leningrad, Moscow: Izd. Glavsevmorputi. p. 147.
- Charrier, I., Bulet, A., & Aubin, T. (2010). Social vocal communication in captive Pacific walruses *Odobenus rosmarus divergens*. *Mammalian Biology*, 2010, 6. doi: doi:10.1016/j.mambio.2010.10.006.
- Christian, J. R., Mathieu, A., Thomson, D. H., White, D., & Buchanan, R. A. (2003). *Effect of seismic energy on snow crab (Chionoecetes opilio)*. (Environmental Research Funds Report No. 144). Calgary: Environmental Studies Research Fund. p. 106.
- Christiansen, J. S., & Reist, J. D. (2013). *Fishes*. Akureyri, Iceland: Conservation of Arctic Flora and Fauna (CAFF), pp. 192-245.
- Citta, J. J., Quakenbush, L. T., Okkonen, S. R., Druckenmiller, M. L., Maslowski, W., Clement-Kinney, J., George, J. C., Brower, H., Small, R. J., Ashjian, C. J., Harwood, L. A., & Heide-Jorgensen, M. P. (2015). Ecological characteristics of core-use areas used by Bering–Chukchi–Beaufort (BCB) bowhead whales, 2006–2012. *136*, 201-222.
- Clark, C. W., Ellison, W. T., Southall, B. L., Hatch, L. T., Parijs, S. M. V., Frankel, A. S., & Ponirakis, D. (2009). Acoustic masking in marine ecosystems: intuitions, analysis, and implication. *Marine Ecology Progress Series*, 395, 201-222.
- Clarke, J., Stafford, S. E., Moore, S. E., Rone, B., Aerts, L., & Crance, J. (2013). Subarctic cetaceans in the southern Chukchi Sea: Evidence of recovery or response to a changing ecosystem. *26*(4), 136-149.
- Clarke, J. T., Brower, A. A., Christman, C. L., & Ferguson, M. C. (2014). *Distribution and Relative Abundance of Marine Mammals in the Northeastern Chukchi and Western Beaufort Seas, 2013. Final Report*. (OCS Study BOEM 2014-018). Seattle, WA: National Marine Mammal Laboratory.
- Clarke, J. T., Ferguson, M. C., Curtice, C., & Harrison, J. (2015). 8. Biologically Important Areas for cetaceans within U.S. waters – Arctic region. *41*(1), 94-103.
- Cohen, D. M., Iwamoto, I. T., & Scialabba, N. (1990). *Vol 10. Gadiform fishes of the world (Order Gadiformes). An annotated and illustrated catalogue of cods, hakes, grenadiers, and other gadiform fishes known to date*. Rome, Italy. p. 442
- Colleye, O., Kever, L., Lecchini, D., Bertin, L., & Parmentier, E. (2016). Auditory evoked potential audiograms in post-settlement stage individuals of coral reef fishes.
- Comiso, J. C., Parkinson, C. L., Gersten, R., & Stock, L. (2008). Accelerated decline in the Arctic sea ice cover. *Geophysical research letters*, 35(1).
- Coombs, S., & Montgomery, J. C. (1999). The enigmatic lateral line system. *Comparative Hearing: Fish and Amphibians*.
- Cooper, L. W., Guarinello, M. L., Grebmeier, J. M., Bayard, A., Lovvorn, J. R., North, C. A., & Kolts, J. M. (2019). A video seafloor survey of epibenthic communities in the Pacific Arctic including Distributed Biological Observatory stations in the northern Bering and Chukchi seas. *Deep Sea Research Part II: Topical Studies in Oceanography*, 162, 164-179.
- Crawford, J. A., Frost, K. J., Quakenbush, L. T., & Whiting, A. (2012). Different habitat use strategies by subadult and adult ringed seals (*Phoca hispida*) in the Bering and Chukchi seas. [journal article]. *Polar Biology*, 35(2), 241-255. doi: 10.1007/s00300-011-1067-1.
- Crawford, R. E. (2003). *Forage fish habitat distribution near the Alaskan coastal shelf areas of the Beaufort and Chukchi seas*. (Contract 09-10-03). PWS Science Center. p. 101.
- Croll, D. A., Gaston, A. J., Burger, A. E., & Konnoff, D. (1992). Foraging behavior and physiological adaptation for diving in thick-billed murre. *Ecology*, 73, 344-356.
- Crowell, S. C. (2016). Measuring in-air and underwater hearing in seabirds. *The Effects of Noise on Aquatic Life II*.

- Crowell, S. E., Wells-Berlin, A. M., Carr, C. E., Olsen, G. H., Therrien, R. E., Yannuzzi, S. E., & Ketten, D. R. (2015). A comparison of auditory brainstem responses across diving bird species. *Journal of Comparative Physiology A*.
- Cummings, W. C., & Thompson, P. O. (1971). Gray whales, *Eschrichtius robustus*, avoid the underwater sounds of killer whales, *Orcinus orca*. *Fishery Bulletin*.
- Davidson, J. G., Dong, H., Linné, M., Andersson, M. H., Piper, A., Prystay, T. S., Hvam, E. B., Thorstad, E. B., Whoriskey, F., & Cooke, S. J. (2019). Effects of sound exposure from a seismic airgun on heart rate, acceleration and depth use in free-swimming Atlantic cod and saithe. *Conservation physiology*, 7(1), coz020.
- Deng, X., Wagner, H.-J., & Popper, A. N. (2011). The inner ear and its coupling to the swim bladder in the deep-sea fish *Antimora rostrata* (Teleostei: Moridae). *Deep Sea Res., Part I*, 58, 27-37.
- Deng, X., Wagner, H. J., & Popper, A. N. (2013). Interspecific Variations of Inner Ear Structure in the Deep-Sea Fish Family Melamphaidae. *The Anatomical Record*, 296(7), 1064-1082.
- Department of Fisheries and Oceans. (2000). *Eastern Beaufort Sea beluga whales*. Department of Fisheries and Oceans.
- Derraik, J. G. B. (2002). The pollution of the marine environment by plastic debris: A review. *Marine Pollution Bulletin*, 44, 842-852.
- Dome Petroleum Ltd., Esso Resources Canada Ltd., & Inc., G. C. R. (1982). Environmental Impact Statement for hydrocarbon development in the Beaufort Sea-Mackenzie Delta Region. *Volume 3A: Beaufort Sea-Delta Setting*.
- Dooling, R. J. (2002). *Avian hearing and the avoidance of wind turbines*. Golden, CO. p. 84.
- Dooling, R. J., Lohr, B., & Dent, M. L. (2000). Hearing in birds and reptiles. *Comparative Hearing, Birds and Reptiles*, 13, 308-359.
- Dooling, R. J., & Therrien, S. C. (2012). Hearing in birds: What changes from air to water. In *The Effects of Noise on Aquatic Life* (pp. 77-82): Springer.
- Durner, G. M., Amstrup, S. C., & Ambrosius, K. J. (2001). Remote identification of polar bear maternal den habitat in northern Alaska. *Arctic*, 54(2), 115-121.
- Durner, G. M., Amstrup, S. C., & Ambrosius, K. J. (2006a). Polar bear maternal den habitat in the Arctic national Wildlife refuge, Alaska. *Arctic*, 59(1), 31-36.
- Durner, G. M., Amstrup, S. C., Nielson, R. M., & McDonald, T. L. (2004). *The use of sea ice habitat by female polar bears in the Beaufort Sea*. (OCS study, MMS 2004- 014). Anchorage, AK: Alaska Science Center. p. 41.
- Durner, G. M., Douglas, D. C., Nielson, R. M., & Amstrup, S. C. (2006b). *A model for autumn pelagic distribution of adult female polar bears in the Chukchi Sea, 1987-1994*. (Contract Completion Report 70181-5-N240). Anchorage, AK: USGS Science Center. p. 67.
- Durner, G. M., Douglas, D. C., Nielson, R. M., Amstrup, S. C., McDonald, T. L., Stirling, I., Mauritzen, M., Born, E. W., Wiig, O., DeWeaver, E., Serreze, M. C., Belikov, S. E., Holland, M. M., Maslanik, J., Aars, J., Bailey, D. C., & Derocher, A. E. (2009). Predicting 21st century polar bear habitat distribution from global climate models. *Ecological Monographs*, 79(1), 25-58.
- Durner, G. M., Fischbach, A. S., Amstrup, S. C., & Douglas, D. C. (2010). *Catalogue of polar bear (Ursus maritimus) maternal den locations in the Beaufort Sea and neighboring regions, Alaska, 1910–2010: U.S. Geological Survey Data Series 568*. p. 14.
- Edds-Walton, P. L. (1997). Acoustic communication signals of mysticete whales. *Bioacoustics*, 8(1-2), 47-60.
- Edmonds, N. J., Firmin, C. J., Goldsmith, D., Faulkner, R. C., & Wood, D. T. (2016). A review of crustacean sensitivity to high amplitude underwater noise: Data needs for effective risk assessment in relation to UK commercial species. *Marine Pollution Bulletin*. doi: 10.1016/j.marpolbul.2016.05.006.

- Ellison, W., Southall, B., Clark, C., & Frankel, A. (2012). A new context-based approach to assess marine mammal behavioral responses to anthropogenic sounds. *Conservation Biology*, 26(1), 21-28.
- Elsner, R., Wartzok, D., Sonafrank, N. B., & Kelly, B. P. (1989). Behavioral and physiological reactions of Arctic seals during under-ice pilotage. *Canadian Journal of Zoology*, 67(10), 2506-2513.
- Environmental Sciences Group. (2005). *Canadian Forces Maritime Experimental and Test Ranges (CFMETR) Environmental Assessment Update 2005*. (RMC-CCE-ES-05-21). Kingston, Ontario, Canada: Environmental Sciences Group, Royal Military College. p. 652.
- Erbe, C., & Farmer, D. M. (2000). Zones of Impact Around Icebreakers Affecting Beluga Whales in the Beaufort Sea. *108*(3), 1332-1340.
- Ershova, E. A., Hopcroft, R. R., Kosobokova, K. N., Matsuno, K., Nelson, R. J., Yamaguchi, A., & Eisner, L. B. (2015). Long-term changes in summer zooplankton communities of the western Chukchi Sea, 1945-2012. *28*(3), 100-115.
- Eschmeyer, W. N., & Fong, J. D. (2017). Species by Family/Subfamily in the Catalog of Fishes. *Catalog of Fishes* Retrieved from <http://researcharchive.calacademy.org/research/ichthyology/catalog/speciesbyfamily.asp> as accessed on 31 October 2017.
- Fay, F. H. (1974). *The role of ice in the ecology of marine mammals of the Bering Sea*: University of Alaska, Institute of Marine Science.
- Fay, F. H., Kelly, B. P., Gehrlich, P. H., Sease, J. L., & Hoover, A. A. (1984). *Modern populations, migrations, demography, trophics, and historical status of the Pacific walrus*. (Final Report R.U. #611). Anchorage, AK: NOAA Outer Continental Shelf Environmental Assessment Program. p. 142.
- Fedoseev, G. A. (2000). *Population biology of ice-associated forms of seals and their role in the northern Pacific ecosystems*. Moscow, Russia: Center for Russian Environmental Policy.
- Finneran, J. J. (2015). Noise-induced hearing loss in marine mammals: A review of temporary threshold shift studies from 1996 to 2015. *The Journal of the Acoustical Society of America*, 138(3), 1702-1726.
- Finneran, J. J., Carder, D. A., Schlundt, C. E., & Dear, R. L. (2010). Growth and recovery of temporary threshold shift at 3 kHz in bottlenose dolphins: Experimental data and mathematical models. *Journal of the Acoustical Society of America*, 127(5), 3256–3266.
- Finneran, J. J., Carder, D. A., Schlundt, C. E., & Ridgway, S. H. (2005). Temporary Threshold Shift in Bottlenose Dolphins (*Tursiops truncatus*) Exposed to Mid-frequency Tones. *Journal of the Acoustic Society of America*, 118(4), 2696-2705.
- Finneran, J. J., & Jenkins, A. K. (2012). *Criteria and Thresholds for Navy Acoustic Effects Analysis Technical Report*. SPAWAR Marine Mammal Program.
- Finneran, J. J., & Schlundt, C. E. (2003). *Effects of Intense Pure Tones on the Behavior of Trained Odontocetes*. San Diego, CA: Space and Naval Warfare Systems Center. p. 18.
- Finneran, J. J., & Schlundt, C. E. (2010). Frequency-dependent and longitudinal changes in noise-induced hearing loss in a bottlenose dolphin (*Tursiops truncatus*). *The Journal of the Acoustical Society of America*, 128(2), 567-570.
- Finneran, J. J., & Schlundt, C. E. (2013). Effects of fatiguing tone frequency on temporary threshold shift in bottlenose dolphins (*Tursiops truncatus*). *The Journal of the Acoustical Society of America*, 133(3), 1819-1826.
- Frost, K. J. (1985). The ringed seal (*Phoca hispida*). In Burns, J. J., Frost, K. J. & Lowry, L. F. (Eds.), *Marine Mammals Species Accounts*. Juneau, AK: Alaska Department of Fish and Game.
- Frost, K. J., & Lowry, L. F. (1984). Trophic relationships of vertebrate consumers in the Alaskan Beaufort Sea. In *The Alaskan Beaufort Sea -- Ecosystems and Environments* (pp. 381-401). New York, NY: Academic Press, Inc.

- Frost, K. J., & Lowry, L. F. (1990). Distribution, abundance, and movements of beluga whales, *Delphinapterus leucas*, in coastal waters of western Alaska. *Canadian Bulletin of Fisheries and Aquatic Sciences*, 224, 39-57.
- Frost, K. J., Lowry, L. F., & Carroll, G. M. (1993). Beluga whale and spotted seal use of a coastal lagoon system in the northeastern Chukchi Sea. *Arctic*, 46(1), 8-16.
- Frost, K. J., Lowry, L. F., & Nelson, R. R. (1985). Radiotagging studies of belukha whales (*Delphinapterus leucas*) in Bristol Bay, Alaska. *Marine Mammal Science*, 1(3), 191-202.
- Frost, K. J., Lowry, L. F., Pendleton, G., & Nute, H. R. (2004). Factors affecting the observed densities of ringed seals, *Phoca hispida*, in the Alaskan Beaufort Sea, 1996–99. *Arctic*, 57(2), 115-128.
- Frouin-Mouy, H., Mouy, X., Berchok, C. L., Blackwell, S. B., & Stafford, K. M. (2019). Acoustic occurrence and behavior of ribbon seals (*Histiophoca fasciata*) in the Bering, Chukchi, and Beaufort seas. *Polar Biology*, 42(4), 657-674.
- Garlich-Miller, J., McCracken, J. G., Snyder, J., Meehan, R., Myers, M., Wilder, J. M., Lance, E., & Matz, A. (2011). *Status Review of the Pacific Walrus (Odobenus rosmarus divergens)*. Anchorage, AK: U.S. Fish and Wildlife Service, Marine Mammals Management.
- Gaston, A. J., & Hipfner, J. M. (2000). Thick-billed murre (*Uria lomvia*). In: Poole, A. (Ed.), *The Birds of North America*. Philadelphia, PA: The Birds of North America Inc.
- Gerstein, E. R. (2002). Manatees, bioacoustics and boats: hearing tests, environmental measurements and acoustic phenomena may together explain why boats and animals collide. *American Scientist*, 90(2), 154-163.
- Ghoul, A., & Reichmuth, C. (2014). Hearing in the sea otter (*Enhydra lutris*): auditory profiles for an amphibious marine carnivore. *Journal of Comparative Physiology A*, 200(11), 967-981.
- Godin, O. A. (2006). Anomalous transparency of water-air interface for low-frequency sound. *Physical Review Letters*, 97(16), 164301. doi: 10.1103/PhysRevLett.97.164301.
- Goetz, K. T., Robinson, P. W., Hobbs, R. C., Laidre, K. L., Huckstadt, L. A., & Shelden, K. E. W. (2012). *Movement and dive behavior of beluga whales in Cook Inlet, Alaska*. Seattle, WA: Alaska Fisheries Science Center, National Marine Fisheries Service. p. 48.
- Golda, J. (2015). *Pacific gray whales*. National Park Service. p. 4.
- Goodall, C., Chapman, C., & Neil, D. (1990). *The acoustic response threshold of Norway lobster Nephrops norvegicus (L.) in a free sound field*: Birkhäuser Basel.
- Gormezano, L. J., & Rockwell, R. F. (2013). Dietary composition and spatial patterns of polar bear foraging on land in western Hudson Bay. *BMC ecology*, 13(1), 51.
- Götz, T., Janik, V. M. G., T., & Janik, V. M. (2010). Aversiveness of sounds in phocid seals: psycho-physiological factors, learning processes and motivation. *The Journal of Experimental Biology*, 213, 1536-1548.
- Gradinger, R., Bluhm, B., & Iken, K. (2010). Arctic sea-ice ridges- Safe havens for sea-ice fauna during periods of extreme ice melt? *Deep-Sea Research II*, 57, 86-95.
- Gradinger, R. R., & Bluhm, B. A. (2004). In-situ observations of the distribution and behavior of amphipods and Arctic cod (*Boreogadus saida*) under the sea ice of the High Arctic Canada Basin. *Polar Biology*, 27, 595-603.
- Graduate School of Oceanography. (2021). How do marine invertebrates detect sounds? *Animals and Sound* Retrieved from <https://dosits.org/animals/sound-reception/how-do-marine-invertebrates-detect-sounds/> as accessed on 18 April 2022.
- Grebmeier, J. M., Bluhm, B. A., Cooper, L. W., Denisenko, S. G., Iken, K., Kędra, M., & Serratos, C. (2015). Time-series benthic community composition and biomass and associated environmental characteristics in the Chukchi Sea during the RUSALCA 2004–2012 Program. *Oceanography*, 28(3), 116-133.

- Grebmeier, J. M., Frey, K. E., Cooper, L. W., & Kędra, M. (2018). Trends in benthic macrofaunal populations, seasonal sea ice persistence, and bottom water temperatures in the Bering Strait region. *Oceanography*, 31(2), 136-151.
- Green, D. M. (1994). Sound's effects on marine mammals need investigation. *Eos, Transactions American Geophysical Union*.
- Green, D. M., DeFerrari, H., McFadden, D., Pearse, J., Popper, A., Richardson, W. J., Ridgway, S. H., & Tyack, P. (1994). Low-frequency sound and marine mammals: Current knowledge and research needs.
- Guglielmo, L., Carrada, G., Catalano, G., Dell'Anno, A., Fabiano, M., Lazzara, L., Mangoni, O., Pusceddu, A., & Saggiomo, V. (2000). Structural and functional properties of sympagic communities in the annual sea ice at Terra Nova Bay (Ross Sea, Antarctica). *Polar Biology*, 23(2), 137-146.
- Gurevich, V. S. (1980). Worldwide distribution and migration patterns of the white whale (Beluga), *Delphinapterus leucas*. *Reports of the International Whaling Commission*, 30, 465-480.
- Halvorsen, M. B., Zeddies, D. G., Ellison, W. T., Chicoine, D. R., & Popper, A. N. (2012). Effects of mid-frequency active sonar on hearing in fish. *The Journal of the Acoustical Society of America*, 131(1), 599-607.
- Hammill, M. O. (2008). Ringed seal *Pusa hispida*. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (Second Edition ed., pp. 972-974). San Diego, CA: Academic Press.
- Hanlon, R. T. (1987). Why Cephalods Are Probably Not Deaf. *The American Naturalist*, 129(2), 312 - 317.
- Hansen, K. A., Hernandez, A., Mooney, T. A., Rasmussen, M. H., Sørensen, K., & Wahlberg, M. (2020). The common murre (*Uria aalge*), an auk seabird, reacts to underwater sound. *J. Acoust. Soc. Am*, 147, 4069-4074.
- Hansen, K. A., Maxwell, A., Siebert, U., Larsen, O. N., & Wahlberg, M. (2017). Great cormorants (*Phalacrocorax carbo*) can detect auditory cues while diving. *The Science of Nature*(5-6).
- Harding, G. C. (1966). *Zooplankton distribution in the Arctic Ocean with notes of life cycles*. Masters of Science, McGill University.
- Harris, R. E., Miller, G. W., & Richardson, W. J. (2001). Seal Responses to Airgun Sounds During Summer Seismic Surveys in the Alaskan Beaufort Sea. *Marine Mammal Science*, 17(4), 795-812.
- Harwood, L. A., McLaughlin, F., Allen, R. M., Illasiak Jr., J., & Alikamik, J. (2005). First-ever marine mammal and bird observations in the deep Canada Basin and Beaufort/Chukchi Seas: expeditions during 2002. *Polar Biology*, 28, 250-253.
- Harwood, L. A., Smith, T. G., Auld, J., Melling, H., & Yurkowski, D. J. (2015). Seasonal movements and diving of ringed seals, *Pusa hispida*, in the Western Canadian Arctic, 1999-2001 and 2010-11. *Arctic*, 193-209.
- Harwood, L. A., Smith, T. G., & Auld, J. C. (2012). Fall migration of ringed seals (*Phoca hispida*) through the Beaufort and Chukchi Seas, 2001 - 02. *Arctic*, 65(1), 35-44.
- Hastings, M. C., & Popper, A. N. (2005). *Effects of sound on fish*. Jones & Stokes for the California Department of Transportation.
- Hatch, L. T., Clark, C. W., Van Parijs, S. M., Frankel, A. S., & Ponirakis, D. W. (2012). Quantifying Loss of Acoustic Communication Space for Right Whales in and around a U.S. National Marine Sanctuary. *Conservation Biology*, 26(6), 983-994. doi: 10.1111/j.1523-1739.2012.01908.x.
- Hauser, D. D., Laidre, K. L., Stern, H. L., Moore, S. E., Suydam, R. S., & Richard, P. R. (2017). Habitat selection by two beluga whale populations in the Chukchi and Beaufort seas. *PLoS One*, 12(2), e0172755.
- Hawkins, A. D., Pembroke, A. E., & Popper, A. N. (2015). Information gaps in understanding the effects of noise on fishes and invertebrates. *Reviews in Fish Biology and Fisheries*.

- Hawkins, A. D., & Popper, A. N. (2017). A sound approach to assessing the impact of underwater noise on marine fishes and invertebrates. *ICES Journal of Marine Science*, 74(3), 635-651.
- Hazard, K. (1988). Beluga whale, *Delphinapterus leucas*. In: Lentfer, J. W. (Ed.), *Selected marine mammals of Alaska. Species accounts with research and management recommendations* (pp. 275). Washington D. C.: Marine Mammal Commission.
- Hazel, J., Lawler, I. R., Marsh, H., & Robson, S. (2007). Vessel Speed Increases Collision Risk for the Green Turtle *Chelonia mydas*. *Endangered Species Research*, 3, 105-113.
- Heide-Jørgensen, M., Laidre, K., Wiig, Ø., Jensen, M., Dueck, L., Maiers, L., Schmidt, H., & Hobbs, R. (2003). From Greenland to Canada in ten days: tracks of bowhead whales, *Balaena mysticetus*, across Baffin Bay. *Arctic*, 21-31.
- Helfman, G. S., Collette, B. B., Facey, D. E., & Bowen, B. W. (2009). The Diversity of Fishes: Biology, Evolution, and Ecology, 2nd. In (pp. 528). Malden, MA: Wiley-Blackwell.
- Henderson, D., Bielefeld, E. C., Carney Harris, K., & Hua Hu, B. (2006). The role of oxidative stress in noise-induced hearing loss. *Ear and Hearing*.
- Henninger, H. P., & Watson, W. H. I. (2005). Mechanisms underlying the production of carapace vibrations and associated waterborne sounds in the American lobster, *Homarus americanus*. *The Journal of Experimental Biology*, 208, 3421-3429. doi: 10.1242/jeb.01771.
- Heptner, L. V., Chapskii, K. K., Arsen'ev, V. A., & Sokolov, V. T. (1976). *Bearded seal. Erignathus barbatus (Erleben, 1777)* (Vol. Vol. II, Part 3. Pinnipeds and Toothed Whales, Pinnipedia and Odontoceti). Moscow, Russia: Vysshaya Shkola Publishers.
- Hetherington, T. (2008). Comparative anatomy and function of hearing in aquatic amphibians, reptiles, and birds. *Sensory evolution on the threshold: adaptations in secondarily aquatic vertebrates*, 183-209.
- Hewitt, R. P. (1985). Reaction of Dolphins to a Survey Vessel: Effects on Census Data. *Fishery Bulletin*, 83(2), 187-193.
- Higgs, D. M., & Radford, C. A. (2013). The contribution of the lateral line to 'hearing' in fish. *Journal of Experimental Biology*, 216(8), 1484-1490.
- Hill, P. S. M. (2009). How do animals use substrate-borne vibrations as an information source? *Naturwissenschaften*, 96, 1355-1371. doi: 10.1007/s00114-009-0588-8.
- Hinzman, L. D., Bettez, N. D., Bolton, W. R., Chapin, F. S., Dyrgerov, M. B., Fastie, C. L., Griffith, B., Hollister, R. D., Hope, A., Huntington, H. P., Jensen, A. M., Jia, G. J., Jorgenson, T., Kane, D. L., Klein, D. R., Kofinas, G., Lynch, A. H., Lloyd, A. H., McGuire, A. D., Nelson, F. E., Oechel, W. C., Osterkamp, T. E., Racine, C. H., Romanovsky, V. E., Stone, R. S., Stow, D. A., Sturm, M., Tweedie, C. E., Vourlitis, G. L., Walker, M. D., Walker, D. A., Webber, P. J., Welker, J. M., Winker, K. S., & Yoshikawa, K. (2005). Evidence and implications of recent climate change in northern Alaska and other Arctic regions. *Climate Change*, 72, 251-298.
- Holland-Bartels, L., Pierce, B., & eds. (2011). *An evaluation of the science needs to inform decisions on the outer continental shelf energy development in the Chukchi and Beaufort Seas*. (U.S. Geological Survey Circular 1370).
- Holst, M., Stirling, I., & Hobson, K. A. (2001). Diet of ringed seals (*Phoca hispida*) on the east and west sides of the north water polynya, northern Baffin Bay. *Marine Mammal Science*, 17(4), 888-908.
- Holt, M. M., Noren, D. P., & Emmons, C. K. (2011). Effects of Noise Levels and Call Types on the Source Levels of Killer Whale Calls. *Journal of the Acoustical Society of America*, 130(5), 3100-3106.
- Hong, Q., Wang, Y., Luo, X., Chen, S., Chen, J., Cai, M., & Mai, B. (2012). Occurrence of polychlorinated biphenyls (PCBs) together with sediment properties in the surface sediments of the Bering Sea, Chukchi Sea, and Canada Basin. *Chemosphere*, 88(11), 1340-1345.

- Hopcroft, R., Bluhm, B., & Gradinger, R. (2008). *Arctic Ocean synthesis: Analysis of climate change impacts in the Chukchi and Beaufort Seas with strategies for future research*. Fairbanks, AK. p. 184.
- Hopcroft, R. R., Clarke, C., Nelson, R. J., & Raskoff, K. A. (2005). Zooplankton communities of the Arctic's Canada Basin: the contribution by smaller taxa. *Polar Biology*, 28(3), 198-206.
- Houser, D. S., Martin, S. W., & Finneran, J. J. (2013a). Behavioral responses of California sea lions to mid-frequency (3250-3450 Hz) sonar signals. *Marine Environmental Research*, 92, 268-278.
- Houser, D. S., Martin, S. W., & Finneran, J. J. (2013b). Exposure amplitude and repetition affect bottlenose dolphin behavioral responses to simulated mid-frequency sonar signals. *Journal of Experimental Marine Biology and Ecology*, 443, 123-133.
- Hu, M. Y., Yan, H. Y., Chung, W. S., Shiao, J. C., & Hwang, P. P. (2009). Acoustically evoked potentials in two cephalopods inferred using the auditory brainstem response (ABR) approach. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 153(3), 278-283. doi: 10.1016/j.cbpa.2009.02.040.
- Hudson, D. M., Krumholz, J. S., Pochtar, D. L., Dickenson, N. C., Dossot, G., Phillips, G., Baker, E. P., & Moll, T. E. (2022). Potential impacts from simulated vessel noise and sonar on commercially important invertebrates. *PeerJ*, 10, e12841.
- Hudson, E., Aihoshi, D., Gaines, T., Simard, G., & Mullock, J. (2001). *The Weather of Nunavut and the Arctic*. Ottawa, Canada: NAV Canada.
- Huntington, H. P., Daniel, R., Hartsig, A., Harun, K., Heiman, M., Meehan, R., Noongwook, G., Pearson, L., Prior-Parks, M., Robards, M., & Stetson, G. (2015). Vessels, Risks, and Rules: Planning for Safe Shipping in Bering Strait. *Marine Policy*, 51, 119-127.
- International Union for the Conservation of Nature. (2016). The IUCN red list of threatened species Retrieved from <http://www.iucnredlist.org> as accessed on 7 November 2016.
- Ivanova, S. V., Kessel, S. T., Espinoza, M., McLean, M. F., O'Neill, C., Landry, J., Hussey, N. E., Williams, R., Vagle, S., & Fisk, A. T. (2020). Shipping alters the movement and behavior of Arctic cod (*Boreogadus saida*), a keystone fish in Arctic marine ecosystems. *Ecological Applications*, 30(3), e02050.
- Jackson, J., Carmack, E., McLaughlin, F., Allen, S., & Ingram, R. (2010). Identification, characterization, and change of the near-surface temperature maximum in the Canada Basin. *J. Geophys. Res.*, 115. doi: 10.1029/2009JC005265.
- Jansen, J. K., Boveng, P. L., Dahle, S. P., & Bengtson, J. L. (2010). Reaction of harbor seals to cruise ships. *Journal of Wildlife Management*.
- Jefferson, T. A., Hung, S. K., & Wursig, B. (2009). Protecting Small Cetaceans from Coastal Development: Impact Assessment and Mitigation Experience in Hong Kong. *Marine Policy*, 33, 305-311.
- Jefferson, T. A., Karczmarski, L., Laidre, K. L., O'Corry-Crowe, G. M., Reeves, R. R., Rojas-Bracho, L., Secchi, E. R., Slooten, E., Smith, B. D., Wang, J. Y., & Zhou, K. (2012). *Delphinapterus leucas*, International Union for Conservation of Nature 2010. *International Union for Conservation of Nature Red List of Threatened Species. Version 2010.4* Retrieved from <http://www.iucnredlist.org/apps/redlist/details/6335/0> as accessed on 28 July 2014.
- Jefferson, T. A., Leatherwood, S., & Webber, M. A. (1993). *FAO species identification guide; marine mammals of the world*. Rome: Food and Agriculture Organization (FAO).
- Jefferson, T. A., Webber, M. A., & Pitman, R. L. (2008). *Marine mammals of the world: A comprehensive guide to their identification*. In (pp. 573). London, UK: Elsevier.
- Jeffries, M. O., Richter-Menge, J., & Overland, J. E. (2014). *Arctic Report Card 2014*.
- Johansen, S., Larsen, O. N., Christensen-Dalsgaard, J., Seidelin, L., Huulvej, T., Jensen, K., Lunneryd, S.-G., Boström, M., & Wahlberg, M. (2016). In-air and underwater hearing in the great cormorant (*Phalacrocorax carbo sinensis*). In *The Effects of Noise on Aquatic Life II* (pp. 505-512): Springer.

- Jones, I. T., Peyla, J. F., Clark, H., Song, Z., Stanley, J. A., & Mooney, T. A. (2021). Changes in feeding behavior of longfin squid (*Doryteuthis pealeii*) during laboratory exposure to pile driving noise. *Marine Environmental Research*, *165*, 105250.
- Jones, I. T., Stanley, J. A., & Mooney, T. A. (2020). Impulsive pile driving noise elicits alarm responses in squid (*Doryteuthis pealeii*). *Marine Pollution Bulletin*, *150*, 110792.
- Jones, J. M., Thayre, B. J., Roth, E. H., Mahoney, M., Sia, I., Mercurief, K., Jackson, C., Zeller, C., Clare, M., & Bacon, A. (2014). Ringed, bearded, and ribbon seal vocalizations north of Barrow, Alaska: Seasonal presence and relationship with sea ice. *Arctic*, *67*(2), 203–222.
- Jørgensen, R., Olsen, K. K., Falk-Petersen, I.-B., & Kanapthippilai, P. (2005). *Investigations of Potential Effects of Low Frequency Sonar Signals on Survival, Development and Behaviour of Fish Larvae and Juveniles*. Tromsø, Norway: University of Tromsø. p. 51.
- Josefson, A. B., Mokievsky, V., Bergmann, M., Blicher, M. E., Bluhm, B., Cochrane, S., Denisenko, N. V., Hasemann, C., Jørgensen, L. L., Klages, M., Schewe, I., Sejr, M. K., Soltwedel, T., Marcin We, sławski, J., & Włodarska-Kowalczyk, M. (2013). Marine invertebrates. In Meltofte, H. (Ed.), *Arctic biodiversity assessment* (pp. 225-257). Denmark: Conservation of Arctic Flora and Fauna (CAFF), Arctic Council.
- Kacimi, S., & Kwok, R. (2022). Arctic snow depth, ice thickness and volume from ICESat-2 and CryoSat-2: 2018-2021. *Geophysical Research Letters*, e2021GL097448.
- Kaifu, K., Akamatsu, T., & Segawa, S. (2008). Underwater sound detection by cephalopod statocyst. *Fisheries Science*, *74*, 781-786.
- Kaschner, K., Watson, R., Trites, A. W., & Pauly, D. (2006). Mapping World-Wide Distributions of Marine Mammal Species Using a Relative Environmental Suitability (RES) Model. *Marine Ecology Progress Series*, *316*, 285-310.
- Kaschner, K. J., Rius-Barile, J., Kesner-Reyes, K., Garilao, C., Kullander, S. O., Rees, T., & Froese, R. (2013). AquaMaps: Predicted range maps for aquatic species. Retrieved from <http://www.aquamaps.org/>.
- Kastak, D., Reichmuth, C., Holt, M. M., Mulsow, J., Southall, B. L., & Schusterman, R. J. (2007). Onset, growth, and recovery of in-air temporary threshold shift in a California sea lion (*Zalophus californianus*). *Journal of the Acoustical Society of America*, *122*(5), 2916–2924. doi: 10.1121/1.2783111.
- Kastak, D., & Schusterman, R. J. (1996). Temporary threshold shift in a harbor seal (*Phoca vitulina*). *J. Acoust. Soc. Am*, *100*(3).
- Kastak, D., & Schusterman, R. J. (1999). In-air and underwater hearing sensitivity of a northern elephant seal (*Mirounga angustirostris*). *Canadian Journal of Zoology*, *77*, 1751-1758.
- Kastak, D., Schusterman, R. J., Southall, B. L., & Reichmuth, C. J. (1999). Underwater Temporary Threshold Shift Induced by Octave-Band Noise in Three Species of Pinniped. *Journal of the Acoustical Society of America*, *106*(2), 1142-1148.
- Kastak, D., Southall, B. L., Schusterman, R. J., & Kastak, C. R. (2005). Underwater temporary threshold shift in pinnipeds: Effects of noise level and duration. *Journal of the Acoustical Society of America*, *118*(5), 3154-3163.
- Kastelein, R., Van Ligteneberg, C., Gjert, I., & Verboom, W. (1993). Free field hearing tests on wild Atlantic walrus (*Odobenus rosmarus rosmarus*) in air. *Aquat Mamm*, *19*, 143-148.
- Kastelein, R. A. (2002). Walrus. In Perrin, W. F., Würsig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 1294-1300). San Diego, CA: Academic Press.
- Kastelein, R. A. (2009). Walrus *Odobenus rosmarus*. In Perrin, W. F., Würsig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 1212-1217). Amsterdam, The Netherlands: Academic Press.

- Kastelein, R. A., Dubbeldam, J. L., Bakker, M. A. G. d., & Gerrits, N. M. (1996). The anatomy of the walrus head (*Odobenus rosmarus*). Part 4: The ears and their function in aerial and underwater hearing. *22*(2), 95-125.
- Kastelein, R. A., Gransier, R., Hoek, L., Macleod, A., & Terhune, J. M. (2012). Hearing threshold shifts and recovery in harbor seals (*Phoca vitulina*) after octave-band noise exposure at 4 kHz. *The Journal of the Acoustical Society of America*, *132*, 2745.
- Kastelein, R. A., Wensveen, P. J., Hoek, L., & Terhune, J. M. (2009a). Underwater Hearing Sensitivity of Harbor Seals (*Phoca vitulina*) for Narrow Noise Bands Between 0.2 and 80 kHz. *Journal of the Acoustical Society of America*, *126*(1), 476-483.
- Kastelein, R. A., Wensveen, P. J., Hoek, L., Verboom, W. C., & Terhune, J. M. (2009b). Underwater Detection of Tonal Signals Between 0.125 and 100 kHz by Harbor Seals (*Phoca vitulina*). *Journal of the Acoustical Society of America*, *125*(2), 1222-1229.
- Kastelein, R. A., & Wiepkema, P. R. (1989). A digging trough as occupational therapy for Pacific walruses (*Odobenus rosmarus divergens*) in human care. *Aquatic Mammals*, *15*(1), 9-17.
- Kaufman, K. (1996). *Lives of North American Birds* (Vol. First Edition). Boston: Houghton Mifflin Harcourt.
- Keller, A. A., Fruh, E. L., Johnson, M. M., Simon, V., & McGourty, C. (2010). Distribution and abundance of anthropogenic marine debris along the shelf and slope of the US west coast. *Marine Pollution Bulletin*, *60*, 692-700.
- Kelly, B. P. (1988a). *Locating and characterizing ringed seal lairs and breathing holes in coordination with surveys using forward looking infra-red sensors* Fisheries and Oceans Freshwater Institute Final Report. p. 17.
- Kelly, B. P. (1988b). Ribbon seal, *Phoca fasciata*. In: Lentfer, J. W. (Ed.), *Selected Marine Mammals of Alaska: Species Accounts with Research and Management Recommendations* (pp. 95-106). Washington, D.C.: Marine Mammal Commission.
- Kelly, B. P. (1988c). Ringed Seal, *Phoca hispida*. In: Lentfer, J. W. (Ed.), *Selected Marine Mammals of Alaska: Species Accounts with Research and Management Recommendations* (pp. 57-75). Washington, D.C.: Marine Mammal Commission.
- Kelly, B. P., Badajos, O. H., Kunnasranta, M., Moran, J. R., Martinez-Bakker, M., Wartzok, D., & Boveng, P. L. (2010a). Seasonal home ranges and fidelity to breeding sites among ringed seals. *Polar Biology*, *33*, 1095-1109.
- Kelly, B. P., Bengtson, J. L., Boveng, P. L., Cameron, M. F., Dahle, S. P., Jansen, J. K., Logerwell, E. A., Overland, J. E., Sabine, G. L., Waring, G. T., & Wilder, J. M. (2010b). *Status review of the ringed seal (Phoca hispida)*. (NOAA Technical Memorandum NMFS-AFSC-212). Seattle, WA: U.S. Department of Commerce, NOAA, National Marine Fisheries Service (NMFS), Alaska Fisheries Science Center. p. 250.
- Kelly, B. P., Burns, J. J., & Quakenbush, L. T. (1988). *Responses of ringed seals (Phoca hispida) to noise disturbance*. Paper presented at the Symposium on Noise and Marine Mammals, Fairbanks, Alaska.
- Kelly, B. P., Ponce, M., Tallmon, D. A., Swanson, B. J., & Sell, S. K. (2009). *Genetic diversity of ringed seals sampled at breeding sites; implications for population structure and sensitivity to sea ice loss*. University of Alaska Southeast, North Pacific Research Board 631 Final Report. p. 28.
- Ketten, D. R. (1998). *Marine Mammal Auditory Systems: A Summary of Audiometric and Anatomical Data and its Implications for Underwater Acoustic Impacts*. La Jolla, CA: National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southwest Fisheries Science Center. p. 74.
- Klishin, V. O., Popov, V. V., & Supin, A. Y. (2000). Hearing capabilities of a beluga whale, *Delphinapterus leucas*. *Aquatic Mammals*, *26*(3), 212-228.

- Knowlton, A. R., & Kraus, S. D. (2001). Mortality and Serious Injury of Northern Right Whales (*Eubalaena glacialis*) in the Western North Atlantic Ocean. *Journal of Cetacean Research and Management, Special Issue 2*, 193-208.
- Kohlbach, D., Graeve, M., A. Lange, B., David, C., Peeken, I., & Flores, H. (2016). The importance of ice algae-produced carbon in the central Arctic Ocean ecosystem: Food web relationships revealed by lipid and stable isotope analyses. *Limnology and Oceanography*, 61(6), 2027-2044. doi: 10.1002/lno.10351.
- Kosobokova, K. N., & Hopcroft, R. R. (2010). Diversity and vertical distribution of mesozooplankton in the Arctic's Canada Basin. *Deep-Sea Research II*, 57, 96-110.
- Kosobokova, K. N., Hopcroft, R. R., & Hirche, H.-J. (2011). Patterns of zooplankton diversity through the depths of the Arctic's central basins. *Marine Biodiversity*, 41(1), 29-50.
- Kovacs, A., & Mellor, M. (1974). *Sea ice morphology and ice as a geologic agent in the southern Beaufort Sea*. Arlington, VA: Arctic Institute of North America.
- Kovacs, K. M. (2002). Bearded seal. In Perrin, W. F., Würsig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 84-87). San Diego, CA: Academic Press.
- Kramer, M., Swadling, K. M., Meiners, K. M., Kiko, R., Scheltz, A., Nicolaus, M., & Werner, I. (2011). Antarctic sympagic meiofauna in winter: Comparing diversity, abundance and biomass between perennially and seasonally ice-covered regions. *Deep Sea Research Part II: Topical Studies in Oceanography*, 58(9-10), 1062-1074. doi: <http://dx.doi.org/10.1016/j.dsr2.2010.10.029>.
- Kraus, S. D., Moore, K. E., Price, C. A., Crone, M. J., Watkins, W. A., Winn, H. E., & Prescott, J. H. (1986). The Use of Photographs to Identify Individual North Atlantic Right Whales (*Eubalaena glacialis*). *Reports of the International Whaling Commission*(Special Issue 10), 145-151.
- Krutzikowski, G. K., & Mate, B. R. (2000). Dive and surfacing characteristics of bowhead whales (*Balaena mysticetus*) in the Beaufort and Chukchi Seas. *Canadian Journal of Zoology*, 78, 1182-1198.
- Kuletz, K. J., Ferguson, M. C., Hurley, B., Gall, A. E., Labunski, E. A., & Morgan, T. C. (2015). Seasonal spatial patterns in seabird and marine mammal distribution in the eastern Chukchi and western Beaufort seas: Identifying biologically important pelagic areas. *Progress in Oceanography*, 136, 175-200.
- Kvadsheim, P. H., Sevaldsen, E. M., Folkow, L. P., & Blix, A. S. (2010). Behavioural and physiological responses of hooded seals (*Cystophora cristata*) to 1 to 7 kHz sonar signals. *Aquatic Mammals*, 36(3), 239-247.
- Ladich, F., & Fay, R. R. (2013). Auditory evoked potential audiometry in fish. *Reviews in Fish Biology and Fisheries*.
- Ladich, F., & Popper, A. N. (2004). Parallel evolution in fish hearing organs. In *Evolution of the Vertebrate Auditory System* (Vol. 22, pp. 33). New York, NY: Springer.
- Laist, D. W. (1987). Overview of the Biological Effects of Lost and Discarded Plastic Debris in the Marine Environment. *Marine Pollution Bulletin*, 18(6B), 319-326.
- Laist, D. W., Knowlton, A. R., Mead, J. G., Collet, A. S., & Podesta, M. (2001). Collisions between Ships and Whales. *Marine Mammal Science*, 17(1), 35-75.
- Laist, D. W., & Shaw, C. (2006). Preliminary Evidence that Boat Speed Restrictions Reduce Deaths of Florida Manatees. *Marine Mammal Science*, 22(2), 472-479.
- Larsen, O. N., Wahlberg, M., & Christensen-Dalsgaard, J. (2020). Amphibious hearing in a diving bird, the great cormorant (*Phalacrocorax carbo sinensis*). *Journal of Experimental Biology*, 223(6), jeb217265.
- LeDuc, R. G., Weller, D. W., Hyde, J., Burdin, A. M., Rosel, P. E., Brownell Jr., R. L., Würsig, B., & Dizon, A. E. (2002). Genetic differences between western and eastern North Pacific gray whales (*Eschrichtius robustus*). *Journal of Cetacean Research and Management*, 4(1), 1-5.

- Leet, W. S., Dewees, C. M., Klingbeil, R., & Larson, E. J. (2001). *California's living marine resources: A status report*. California Department of Fish and Game.
- Lentfer, J. W. (1972). *Alaska Polar Bear Research and Management, 1970-1971*. Alaska Department of Fish and Game. pp. 21-39.
- Levinton, J. (2009). *Marine Biology: Function, Biodiversity, Ecology*. In (3rd ed.). New York: Oxford University Press.
- Little, B., & Ray, R. (2002). A perspective on corrosion inhibition by biofilms. *Corrosion*, 58(5), 424-428.
- Logerwell, E., Rand, K., Danielson, S., & Sousa, L. (2018). Environmental drivers of benthic fish distribution in and around Barrow Canyon in the northeastern Chukchi Sea and western Beaufort Sea. *Deep Sea Research Part II: Topical Studies in Oceanography*, 152, 170-181.
- Lomac-MacNair, K., Pedro Andrade, J., & Esteves, E. (2019a). Seal and polar bear behavioral response to an icebreaker vessel in northwest Greenland. *Human-Wildlife Interactions*, 13(2), 13.
- Lomac-MacNair, K. S., Andrade, J. P., & Esteves, E. (2019b). Seal and Polar Bear Behavioral Response to an Icebreaker Vessel in Northwest Greenland. *Human-Wildlife Interactions*, 13(2). doi: 10.26077/pxn3-h858.
- Lombarte, A., Yan, H. Y., Popper, A. N., Chang, J. S., & Platt, C. (1993). Damage and regeneration of hair cell ciliary bundles in a fish ear following treatment with gentamicin. *Hearing Research*, 64(2), 166-174.
- London, J. M., Boveng, P. L., & Cameron, M. F. (2015). *Dive behavior of bearded, ribbon and spotted seals in the Bering and Chukchi Seas*. Seattle, WA: Polar Ecosystems Program, National Marine Mammal Laboratory, Alaska Fisheries Science Center. p. 1.
- Lovell, J. M., Findlay, M. M., Moate, R. M., & Yan, H. Y. (2005). The hearing abilities of the prawn *Palaemon serratus*. 140, 89-100.
- Lovell, J. M., Findlay, M. M., Nedwell, J. R., & Pegg, M. A. (2006). The Hearing Abilities of the Silver Carp (*Hypophthalmichthys molitrix*) and Bighead Carp (*Aristichthys nobilis*). *Comparative Biochemistry and Physiology, Part A*, 143, 286-291.
- Lovvorn, J. R., North, C. A., Kolts, J. M., Grebmeier, J. M., Cooper, L. W., & Cui, X. (2016). Projecting the effects of climate-driven changes in organic matter supply on benthic food webs in the northern Bering Sea. *Marine Ecology Progress Series*, 548, 11-30.
- Lowry, L. F., Burkanov, V. N., Frost, K. J., Simpkins, M. A., Springer, D. P., DeMaster, D. P., & Suydam, R. (2000). Habitat use and habitat selection by spotted seals (*Phoca largha*) in the Bering Sea. *Canadian Journal of Zoology*, 78(1959-1971).
- Lowry, L. F., Frost, K. J., & Burns, J. J. (1980). Variability in the diet of ringed seals, *Phoca hispida*, in Alaska. *Canadian Journal of Zoology*, 37, 2254-2261.
- Lowry, L. F., Frost, K. J., Davis, R. W., DeMaster, D. P., & Suydam, R. S. (1998). Movements and behavior of satellite-tagged spotted seals (*Phoca largha*) in the Bering and Chukchi Seas. *Polar Biology*, 19(4), 221-230.
- Lowry, L. F., Sheffield, G., & George, J. C. (2004). Bowhead whale feeding in the Alaskan Beaufort Sea, based on stomach content analyses. 6(3), 215-223.
- Lunn, N. J., Schliebe, S., & Born, E. W. (2002). *Polar Bears: Proceedings of the 13th working meeting of the IUCN/SSC Polar Bear Specialist Group, 23-28 June 2001, Nuuk, Greenland*. Gland, Switzerland: International Union for the Conservation of Nature (IUCN). p. 153.
- Lunn, N. J., & Stirling, I. (1985). The significance of supplemental food to polar bears during the ice-free period of Hudson Bay. *Canadian Journal of Zoology*, 63(10), 2291-2297.
- Lydersen, C. (1998). Status and biology of ringed seals (*Phoca hispida*) in Svalbard. In. Heide-Jørgensen, M. P. & Lydersen, C. (Eds.), *Ringed Seals in the North Atlantic* (Vol. 1, pp. 46-62). Tromsø, Norway: NAMMCO Scientific Publications.

- Lydersen, C., & Gjertz, I. (1986). Studies of the ringed seal (*Phoca hispida* Schreber 1775) in its breeding habitat in Kongsfjorden, Svalbard. *Polar Research*, 4(1), 57-63.
- Lydersen, C., & Hammill, M. O. (1993). Diving in ringed seal (*Phoca hispida*) pups during the nursing period. *Canadian Journal of Zoology*, 71(5), 991-996.
- Lydersen, C., Jensen, P. M., & Lydersen, E. (1990). A survey of the Van Mijen Fiord, Svalbard, as habitat for ringed seals, *Phoca hispida*. *Ecography*, 13(2), 130-133.
- Lydersen, C., & Ryg, M. (1991). Evaluating breeding habitat and populations of ringed seals *Phoca hispida* in Svalbard fjords. *Polar Record*, 27(162), 223-228.
- MacDonald, I. R., Bluhm, B. A., Iken, K., Gagaev, S., & Strong, S. (2010). Benthic macrofauna and megafauna assemblages in the Arctic deep-sea Canada Basin. *Deep-Sea Research II*, 57, 136-152.
- Macfadyen, G., Huntington, T., Cappell, R. M., G., Huntington, T., & Cappell, R. (2009). *Abandoned, Lost or Otherwise Discarded Fishing Gear*. (UNEP Regional Seas Report and Studies 185, or FAO Fisheries and Aquaculture Technical Paper 523). Rome, Italy: United Nations Environment Programme Food, Food and Agriculture Organization of the United Nations. p. 115.
- MacIntyre, K. Q., Stafford, K. M., Berchok, C. L., & Boveng, P. L. (2013). Year-round acoustic detection of bearded seals (*Erignathus barbatus*) in the Beaufort Sea relative to changing environmental conditions 2008-2010. *Polar Biology*, 36(8), 1161-1173.
- Magalhães, S., Prieto, R., Silva, M. A., Gonçalves, J. M., Afonso-Dias, M., & Santos, R. S. (2002). Short-Term Reactions of Sperm Whales (*Physeter macrocephalus*) to Whale-Watching Vessels in the Azores. *Aquatic Mammals*, 28(3), 267-274.
- Mallett, R. D., Stroeve, J. C., Tsamados, M., Landy, J. C., Willatt, R., Nandan, V., & Liston, G. E. (2021). Faster decline and higher variability in the sea ice thickness of the marginal Arctic seas when accounting for dynamic snow cover. *The Cryosphere*, 15(5), 2429-2450.
- Mann, D. A., Higgs, D. M., Tavalga, W. N., Souza, M. J., & Popper, A. N. (2001). Ultrasound detection by Clupeiform fishes. *Journal of the Acoustical Society of America*.
- Mann, D. A., Lu, Z., Hastings, M. C., & Popper, A. N. (1998). Detection of ultrasonic tones and simulated dolphin echolocation clicks by a teleost fish, the American shad (*Alosa sapidissima*).
- Mann, D. A., Lu, Z., & Popper, A. N. (1997). A clupeid fish can detect ultrasound. *Nature*.
- Martin, B., Zeddies, D. G., Gaudet, B., & Richard, J. (2016). Evaluation of three sensor types for particle motion measurement. In *The Effects of Noise on Aquatic Life II* (pp. 679-686): Springer.
- Maslanik, J., Stroeve, J., Fowler, C., & Emery, W. (2011). Distribution and trends in Arctic sea ice age through spring 2011. *Geophys. Res. Lett.*, 38(13). doi: 10.1029/2011GL047735.
- Maxwell, B. (1997). *Responding to Global Climate Change in Canada's Arctic*. Environment Canada.
- Mayer, L. A., & Armstrong, A. (2012). *U.S. Law of the Sea Cruise to Map and Sample the US Arctic Ocean Margin*. (Paper 1282). Center for Coastal and Ocean Mapping. p. 159.
- McLaren, I. A. (1958). The biology of the ringed seal (*Phoca hispida* Schreber) in the eastern Canadian Arctic. *Fisheries Research Board of Canada*, 118, 97.
- Mecklenburg, C. W., Byrkjedal, I., Christiansen, J. S., Karamushko, O. V., Lynghammar, A., & Moller, P. R. (2013). List of marine fishes of the Arctic Region annotated with common names and zoogeographic characterizations. In (pp. 35). Akureyri, Iceland: Conservation of Arctic Flora and Fauna (CAFF).
- Mecklenburg, C. W., & Mecklenburg, T. (2009). Fishes Retrieved from <http://www.arcodiv.org/Fish.html> as accessed on June 30, 2015.
- Mecklenburg, C. W., Møller, P. R., & Steinke, D. (2011). Biodiversity of arctic marine fishes: taxonomy and zoogeography. *Marine Biodiversity*, 41(1), 109-140.
- Melvin, E. F., Parrish, J. K., & Conquest, L. L. (1999). Novel tools to reduce seabird bycatch in coastal gillnet fisheries. *Conservation Biology*, 13(6), 1386-1397.

- Merchant, N. D., Pirotta, E., Barton, T. R., & Thompson, P. M. (2014). Monitoring Ship Noise to Assess the Impact of Coastal Developments on Marine Mammals. *Marine Pollution Bulletin*, 78, 85-95.
- Michel, C., Nielsen, T. G., Nozais, C., & Gosselin, M. (2002). Significance of sedimentation and grazing by ice micro-and meiofauna for carbon cycling in annual sea ice (northern Baffin Bay). *Aquatic Microbial Ecology*, 30(1), 57-68.
- Miller, G. W., Moulton, V. D., Davis, R. A., Holst, M., Millman, P., MacGillivray, A., & Hannay, D. (2005). Monitoring seismic effects on marine mammals-southeastern Beaufort Sea, 2001-2002. In Armsworthy, S. L., Cranford, P. J. & Lee, K. (Eds.), *Offshore Oil and Gas Environmental Effects Monitoring/Approaches and Technologies* (pp. 511-542). Columbus, OH.
- Miller, P. J. O., Kvadsheim, P. H., Lam, F.-P. A., Wensveen, P. J., Antunes, R., Alves, A. C., Visser, F., Kleivane, L., Tyack, P. L., & Sivle, L. D. (2012). The severity of behavioral changes observed during experimental exposures of killer (Orcinus orca), long-finned pilot (Globicephala melas), and sperm (Physeter macrocephalus) whales to naval sonar. *Aquatic Mammals*, 38(4), 362-401. doi: 10.1578/am.38.4.2012.362.
- Minerals Management Service. (1991). *Chukchi Sea Planning Area Oil and Gas Lease Sale 126. Final Environmental Impact Statement*. (OCS EIS/EA MMS 2002-006). Herndon, VA.
- Misund, O. A. (1997). Underwater acoustics in marine fisheries and fisheries research. *Reviews in Fish Biology and Fisheries*, 7(1), 1-34.
- Møhl, B. (1968a). Auditory Sensitivity on the Common Seal in Air and Water. *Journal of Auditory Research*, 8, 27-38.
- Møhl, B. (1968b). Hearing in Seals *Behavior and Physiology of Pinnipeds*, 172-195.
- Møller, A. R. (2012). *Hearing: anatomy, physiology, and disorders of the auditory system*: Plural Publishing.
- Montgomery, J. C., Jeffs, A., Simpson, S. D., Meekan, M., & Tindle, C. (2006). Sound as an orientation cue for the pelagic larvae of reef fishes and decapod crustaceans. *Advances in Marine Biology*, 51, 143-196.
- Mooney, T. A., Hanlon, R. T., Christensen-Dalsgaard, J., Madsen, P. T., Ketten, D. R., & Nachtigall, P. E. (2010). Sound detection by the longfin squid (Loligo pealeii) studied with auditory evoked potentials: sensitivity to low-frequency particle motion and not pressure. *Journal of Experimental Biology*, 213, 3748-3759.
- Mooney, T. A., Nachtigall, P. E., Breese, M., Vlachos, S., & Au, W. W. L. (2009). Predicting temporary threshold shifts in a bottlenose dolphin (*Tursiops truncatus*): The effects of noise level and duration. *Journal of Acoustical Society of America*, 125(3), 1816-1826. doi: 10.1121/1.3068456.
- Mooney, T. A., Smith, A., Larsen, O. N., Hansen, K. A., Wahlberg, M., & Rasmussen, M. H. (2019). Field-based hearing measurements of two seabird species. *Journal of Experimental Biology*, 222(4), jeb190710.
- Moore, S. E. (2000). Variability of cetacean distribution and habitat selection in the Alaskan Arctic, Autumn 1982-91. *Arctic*, 53(4), 448-460.
- Moore, S. E., DeMaster, D. P., & Dayton, P. K. (2000). Cetacean habitat selection in the Alaskan Arctic during summer and autumn. *Arctic*, 53(4), 432-447.
- Moore, S. E., & Laidre, K. L. (2006). Trends in sea ice cover within habitats used by bowhead whales in the western Arctic. *Arctic*, 59(3), 932-944.
- Moore, S. E., & Reeves, R. R. (1993). Distribution and movement. In Burns, J. J., Montague, J. J. & Cowles, C. J. (Eds.), *The Bowhead Whale* (Vol. 2, pp. 313-386). Lawrence, KS: Society for Marine Mammology.
- Moretti, D., Thomas, L., Marques, T., Harwood, J., Dilley, A., Neals, B., Shaffer, J. A., McCarthy, E., New, L., Jarvis, S., & Morrissey, R. (2014). A risk function for behavioral disruption of Blainville's

- beaked whales (*Mesoplodon densirostris*) from mid-frequency active sonar. *PLoS One*, 9(1), e85064.
- Moulton, V. D., Richardson, W. J., McDonald, T. L., Elliott, R. E., & Williams, M. T. (2002). Factors influencing local abundance and haulout behaviour of ringed seals (*Phoca hispida*) on landfast ice of the Alaskan Beaufort Sea. *Canadian Journal of Zoology*, 80, 1900-1917.
- Murphy, K. A., Davies, H., Shafer, H., Cox, K., Nikolich, K., & Juanes, F. (2019). *Impacts of noise on the behavior and physiology of marine invertebrates: A meta-analysis*. Paper presented at the Proceedings of Meetings on Acoustics 5ENAL.
- Murphy, D. (2010). U.S. Coast Guard Cutter Healy Retrieved from <http://www.who.edu/vanishingarctic/page.do?pid=47775> as accessed on 03 September 2015.
- Muto, M. M., Helker, V. T., Angliss, R. P., Allen, B. A., Boveng, P. L., Breiwick, J. M., Cameron, M. F., Clapham, P. J., Dahle, S. P., Dahlheim, M. E., Fadely, B. S., Ferguson, M. C., Fritz, L. W., Hobbs, R. C., Ivashchenko, Y. V., Kennedy, A. S., London, J. M., Mizroch, S. A., Ream, R. R., Richmond, E. L., Shelden, K. E. W., Towell, R. G., Wade, P. R., Waite, J. M., & Zerbini, A. N. (2017). *Alaska marine mammal stock assessments, 2016*. (NOAA Technical Memorandum NMFS-AFSC-355). U.S. Department of Commerce, National Oceanic and Atmospheric Administration (NOAA), National Marine Fisheries Service (NMFS), Alaska Fisheries Science Center. p. 366.
- Muto, M. M., Helker, V. T., Angliss, R. P., Allen, B. A., Boveng, P. L., Breiwick, J. M., Cameron, M. F., Clapham, P. J., Dahle, S. P., Dahlheim, M. E., Fadley, B. E., Ferguson, M. C., Fritz, L. W., Hobbs, R. C., Ivashchenko, Y. V., Kennedy, A. S., London, J. M., Mizroch, S. A., Ream, R. R., Richmond, E. L., Shelden, K. E. W., Towell, R. G., Wade, P. R., Waite, J. M., & Zerbini, A. N. (2016). *Alaska marine mammal stock assessments, 2015*. (NOAA Technical Memorandum NMFS-AFSC-323). Seattle, WA. p. 300.
- Muto, M. M., Helker, V. T., Delean, B., Young, N., Freed, J., Angliss, R., Friday, N., Boveng, P., Breiwick, J., & Brost, B. (2021). *Alaska marine mammal stock assessments, 2020*.
- Myrberg, A. A. (1981). Sound Communication and Interception in Fishes. In: Tavolga, W. N., Popper, A. N. & Fay, R. R. (Eds.), *Hearing and Sound Communication in Fishes* (pp. 395-452). New York: Springer-Verlag.
- Myrberg, A. A., Jr. (1990). The Effects of Man-Made Noise on the Behavior of Marine Animals. *Environmental International*, 16, 575-586.
- Nachtigall, P. E., Supin, A. Y., Amundin, M., Röken, B., Møller, T., Mooney, T. A., Taylor, K. A., & Yuen, M. M. L. (2007). Polar bear *Ursus maritimus* hearing measured with auditory evoked potentials. *The Journal of Experimental Biology*, 210, 1116-1122.
- National Audubon Society. (2015). *Arctic Marine Synthesis: Atlas of the Chukchi and Beaufort Seas* Retrieved from <http://ak.audubon.org/arctic-marine-synthesis-atlas-chukchi-and-beaufort-seas> as accessed on 24 August 2015.
- National Marine Fisheries Service. (2016). *Technical Guidance for Assessing the Effects of Anthropogenic Sound on Marine Mammal Hearing: Underwater Acoustic Thresholds for Onset of Permanent and Temporary Threshold Shifts*. p. 178 p.
- National Research Council. (2003). *Ocean Noise and Marine Mammals*. Washington, DC: National Academics Press. p. 203.
- National Snow and Ice Data Center. (2007). *National Snow and Ice Data Center World Data Center for Glaciology, Boulder Annual Report 2007*. Boulder, CO. p. 56.
- National Snow and Ice Data Center. (2012). Are icebreakers changing the climate? Retrieved from <https://nsidc.org/cryosphere/icelights/2012/04/are-icebreakers-changing-climate>
- National Snow and Ice Data Center. (2022). *Charctic Interactive Sea Ice Graph*. *Arctic Sea Ice News & Analysis* Retrieved from <https://nsidc.org/arcticseaicenews/charctic-interactive-sea-ice-graph/> as accessed on 16 February, 2022.

- Naval Oceanographic Office. (2014). *Oceanographic and atmospheric master library (OAML) summary document*. (OAML-SUM-21W). p. 148.
- Naval Sea Systems Command. (2015). *Speed Series Starboard Aspect One-third Octave Signature*.
- Nedelec, S. L., Campbell, J., Radford, A. N., Simpson, S. D., & Merchant, N. D. (2016). Particle motion: the missing link in underwater acoustic ecology.
- Nedelec, S. L., Radford, A. N., Simpson, S. D., Nedelec, B., Lecchini, D., & Mills, S. C. (2014a). Anthropogenic noise playback impairs embryonic development and increases mortality in a marine invertebrate. *Scientific Reports*, 4(5891), 4. doi: 10.1038/srep05891.
- Nedelec, S. L., Radford, A. N., Simpson, S. D., Nedelec, B., Lecchini, D., & Mills, S. C. (2014b). Anthropogenic noise playback impairs embryonic development and increases mortality in a marine invertebrate. *Scientific Reports*, 4(1), 1-4.
- Nedelec, S. L., Simpson, S. D., Morley, E. L., Nedelec, B., & Radford, A. N. (2015). Impacts of regular and random noise on the behaviour, growth and development of larval Atlantic cod (*Gadus morhua*). *Proceedings of the Royal Society B: Biological Sciences*, 282(1817), 20151943.
- Nedwell, J. R., Edwards, B., Turnpenny, A. W. H., & Gordon, J. (2004). *Fish and Marine Mammal Audiograms: A Summary of Available Information*. (Subacoustech Report ref: 534R0214). Hampshire, England: Subacoustech Ltd. p. 278.
- Nelson, R. R., Burns, J. J., & Frost, K. J. (1984). The bearded seal (*Erignathus barbatus*). *Marine Mammal Species Accounts, Wildlife Technical Bulletin* 7, 1-6.
- NOAA National Centers for Environmental Information. (2021). State of the Climate: Monthly Global Snow and Ice Report for June 2021 Retrieved from <https://www.ncei.noaa.gov/access/monitoring/monthly-report/global-snow/202106>.
- North Atlantic Marine Mammal Commission. (2004). *The ringed seal*. Tromsø, Norway: North Atlantic Marine Mammal Commission (NAMMCO).
- North Pacific Fishery Management Council. (2009). *Arctic Fishery Management Plan for fish resources of the Arctic Management Area*. Anchorage, AK: North Pacific Fishery Management Council (NPFMC).
- Notz, D., & Community, S. (2020). Arctic sea ice in CMIP6. *Geophysical Research Letters*, 47(10), e2019GL086749.
- Nowacek, D. P., Johnson, M. P., & Tyack, P. L. (2004). North Atlantic Right Whales (*Eubalaena glacialis*) Ignore Ships but Respond to Alerting Stimuli. *Proceedings of the Royal Society of London B*, 271, 227-231.
- Nowacek, D. P., Thorne, L. H., Johnston, D. W., & Tyack, P. L. (2007). Responses of Cetaceans to Anthropogenic Noise. *Mammal Review*, 37(2), 81-115.
- Nummela, S. (2008a). Hearing. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (Second Edition ed., pp. 553-561). Burlington, MA: Academic Press.
- Nummela, S. (2008b). Hearing in aquatic mammals. In Thewissen, J. G. M. & Nummela, S. (Eds.), *Sensory Evolution on the Threshold* (pp. 211-224). Berkeley, CA: University of California Press.
- Nuttall, M. (2005). *Encyclopedia of the Arctic*: Routledge.
- O'Corry-Crowe, G., Mahoney, A. R., Suydam, R., Quakenbush, L., Whiting, A., Lowry, L., & Harwood, L. (2016). Genetic profiling links changing sea-ice to shifting beluga whale migration patterns. *12*, 20160404.
- O'Brien, M. C., Melling, H., Pedersen, T. F., & Macdonald, R. W. (2013). The role of eddies on particle flux in the Canada Basin of the Arctic Ocean. *Deep Sea Research Part I: Oceanographic Research Papers*, 71, 1-20. doi: <http://dx.doi.org/10.1016/j.dsr.2012.10.004>.
- Ocean Conservancy. (2010). *Trash Travels: From Our Hands to the Sea, Around the Globe, and Through Time*. The Ocean conservancy. p. 60.

- Office of Environment Alaska OCS Region. (2012). *ION Geophysical 2012 Seismic Survey: Beaufort Sea and Chukchi Sea, Alaska*. Bureau of Ocean Energy Management. p. 102.
- Offutt, G. C. (1970). Acoustic Stimulus Perception by the American Lobster *Homarus americanus* (Decapoda). *Experientia*, 26, 1276-1278.
- Okkonen, S. R., Ashjian, C. J., Campbell, R. G., Clarke, J., Moore, S. E., & Taylor, K. D. (2011). Satellite observations of circulation features associated with a bowhead whale feeding 'hotspot' near Barrow, Alaska. *115*, 2168-2174.
- Øren, K., Kovacs, K. M., Yoccoz, N. G., & Lydersen, C. (2018). Assessing site-use and sources of disturbance at walrus haul-outs using monitoring cameras. *Polar Biology*, 41(9), 1737-1750.
- Overland, J., Hanna, E., Hanssen, B., I., Kim, S., Walsh, J., Wang, M., & Bhatt, U. S. (2014). *Air Temperature*
- Overland, J. E., & Wang, M. (2013). When will the summer Arctic be nearly sea ice free? *Geophysical Research Letters*, 40(10), 2097-2101. doi: 10.1002/grl.50316.
- Owen, M. A., & Bowles, A. E. (2011). In-air auditory psychophysics and the management of a threatened carnivore, the polar bear (*Ursus maritimus*). *International Journal of Comparative Psychology*, 24(3), 244-254.
- Owl Ridge Natural Resource Consultants Inc. (2017). *Application for the Incidental Harassment Authorization for the Taking of Pacific Walrus and Polar Bears in Conjunction with the Quintillion Subsea Operations Cable Project, 2017. Prepared for: Quintillion Subsea Operations, LLC*. Anchorage, AK.
- Packard, A., Karlsen, H. E., & Sand, O. (1990). Low Frequency Hearing in Cephalopods. *Journal of Comparative Physiology A*, 166, 501-505.
- Palo, J. U. (2003). *Genetic diversity and phylogeography of landlocked seals*. Helsinki, Finland: University of Helsinki.
- Palo, J. U., Makinen, H. S., Helle, E., Stenman, O., & Vainola, R. (2001). Microsatellite variation in ringed seals (*Phoca hispida*): Genetic structure and history of the Baltic Sea population. *Journal of Heredity*, 86, 609-617.
- Parks, S. E., Johnson, M., Nowacek, D., & Tyack, P. L. (2011). Individual Right Whales Call Louder in Increased Environmental Noise. *Biology Letters*, 7, 33-35.
- Patek, S. N., & Caldwell, R. L. (2006). The stomatopod rumble: Low frequency sound production in *Hemisquilla californiensis*. *Marine and Freshwater Behaviour and Physiology*, 39(2), 99-111.
- Paxton, J. R., & Eshmeyer, W. N. (1998). *Encyclopedia of Fishes* (2nd ed.). San Diego, CA: Academic Press.
- Payne, J. F., Andrews, C. A., Fancey, L. L., Cook, A. L., & Christian, J. R. (2007). *Pilot Study on the Effects of Seismic Air Gun Noise on Lobster (Homarus Americanus)*. (Environmental Studies Research Funds Report No. 171).
- Perovich, D., Gerland, S., Hendricks, S., Meier, W., Nicolaus, M., Richter-Menge, J., & Tschudi, M. (2013). Sea ice Retrieved from http://www.arctic.noaa.gov/reportcard/sea_ice.html
- Pine, M. K., Nikolich, K., Martin, B., Morris, C., & Juanes, F. (2020). Assessing auditory masking for management of underwater anthropogenic noise. *The Journal of the Acoustical Society of America*, 147(5), 3408-3417.
- Pine, M. K., Wilson, L., Jeffs, A. G., McWhinnie, L., Juanes, F., Scuderi, A., & Radford, C. A. (2021). A Gulf in lockdown: How an enforced ban on recreational vessels increased dolphin and fish communication ranges. *Global Change Biology*, 27(19), 4839-4848.
- Pirotta, E., Merchant, N. D., Thompson, P. M., Barton, T. R., & Lusseau, D. (2015). Quantifying the Effect of Boat Disturbance on Bottlenose Dolphin Foraging Activity. *Biological Conservation*, 181, 82-89.
- Plueddemann, A., Krishfield, R., Takizawa, T., Hatakeyama, K., & Honjo, S. (1998). Upper ocean velocities in the Beaufort Gyre. *Geophysical research letters*, 25(2), 183-186.

- Polar Bears International. (2015). Home Range. *About Polar Bears* Retrieved from <http://www.polarbearsinternational.org/habits-and-behavior/home-range> as accessed on 8 July 2015.
- Popov, V. V., Supin, A. Y., Wang, D., Wang, K., Dong, L., & Wang, S. (2011). Noise-induced temporary threshold shift and recovery in Yangtze finless porpoises *Neophocaena phocaenoides asiaeorientalis*. *The Journal of the Acoustical Society of America*, 130(1), 574-584.
- Popper, A. N. (2003). Effects of anthropogenic sounds on fishes. *Fisheries Research*, 28(10), 24-31.
- Popper, A. N. (2008). *Effects of mid- and high-frequency sonars on fish*. Newport, RI: Department of the Navy (DoN). p. 52.
- Popper, A. N. (2014). *Classification of fish and sea turtles with respect to sound exposure*. Technical report prepared for ANSI-Accredited. Standards Committee. S3/SC1.
- Popper, A. N., & Fay, R. R. (1994). *Comparative Hearing: Mammals*: Springer-Verlag.
- Popper, A. N., & Fay, R. R. (2010). Rethinking sound detection by fishes. *Hearing Research*, 273, 1-12.
- Popper, A. N., & Hastings, M. C. (2009). The effects of human-generated sound on fish. *Integrative Zoology*.
- Popper, A. N., & Hawkins, A. D. (2018). The importance of particle motion to fishes and invertebrates. *The Journal of the Acoustical Society of America*, 143(1), 470-488.
- Popper, A. N., Plachta, D. T. T., Mann, D. A., & Higgs, D. (2004). Response of Clupeid Fish to Ultrasound: A Review. *ICES Journal of Marine Science*, 61, 1057-1061.
- Popper, A. N., Salmon, M., & Horch, K. W. (2001). Acoustic detection and communication by decapod crustaceans. *Journal of Comparative Physiology A*, 187(2), 83-89.
- Popper, A. N., & Schilt, C. R. (2008). Hearing and acoustic behavior: Basic and applied considerations. In *Fish Bioacoustics* (pp. 17-48). New York, NY: Springer.
- Porsild, A. E. (1945). Mammals of the Mackenzie Delta. *Canadian Field-Naturalist*, 59, 4-22.
- Quakenbush, L. (2008). Bowhead Whale. In Game, A. D. o. F. a. (Ed.). Retrieved from http://www.adfg.alaska.gov/static/education/wns/bowhead_whale.pdf.
- Quakenbush, L. T. (1988). Spotted seal, *Phoca largha*. In: Lentfer, J. W. (Ed.), *Selected Marine Mammals of Alaska: Species Accounts with Research and Management Recommendations* (pp. 107-124). Washington, D.C.: Marine Mammal Commission.
- Quakenbush, L. T., Small, R. J., & Citta, J. J. (2010). *Satellite tracking of western Arctic bowhead whales*. Anchorage, AK. p. 65.
- Radford, C. A., Montgomery, J. C., Caiger, P., & Higgs, D. M. (2012). Pressure and particle motion detection thresholds in fish: a re-examination of salient auditory cues in teleosts. *Journal of Experimental Biology*, 215(19), 3429-3435.
- Rainville, L., Lee, C. M., & Woodgate, R. A. (2011). Impact of wind-driven mixing in the Arctic Ocean. *Oceanography*, 24(3), 136-145. doi: 10.5670/oceanog.2011.65.
- Rand, K. M., & Logerwell, E. A. (2010). The first demersal trawl survey of benthic fish and invertebrates in the Beaufort Sea since the late 1970s. *Polar Biology*, 34, 475-488. doi: 10.1007/s00300-010-0900-2.
- Ravelo, A. M., Bluhm, B. A., Foster, N., & Iken, K. (2020). Biogeography of epibenthic assemblages in the central Beaufort Sea. *Marine Biodiversity*, 50(1), 1-19.
- Reeder, D. M., & Kramer, K. M. (2005). Stress in free-ranging mammals: Integrating physiology, ecology, and natural history.
- Reeves, R. R., Stewart, B. S., Clapham, P. J., & Powell, J. A. (2002). *National Audubon Society Guide to Marine Mammals of the World*. In (pp. 527). New York, NY: Alfred A. Knopf.
- Regehr, E. V., Amstrup, S. C., & Stirling, I. (2006). *Polar bear population status in the southern Beaufort Sea*. (2006-1337). p. 20.

- Reichmuth, C. (2008). Hearing in Marine Carnivores. *Bioacoustics: The International Journal of Animal Sound and its Recording*, 17(1-3), 89-92. doi: 10.1080/09524622.2008.9753777.
- Reichmuth, C. J., Holt, M. M., Mulsow, J., Sills, J. M., & Southall, B. L. (2013). Comparative assessment of amphibious hearing in pinnipeds. *Journal of Comparative Physiology A*.
- Reilly, S. B., Bannister, J. L., Best, P. B., Brown, M. W., Brownell Jr, R. L., Butterworth, D. S., Clapham, P. J., Cooke, J., Donovan, G. P., Urban R., J., & Zerbini, A. N. (2008). *Eschrichtius robustus*. *The IUCN Red List of Threatened Species. Version 2014.2* Retrieved from www.iucnredlist.org as accessed on 4 August 2014.
- Renaud, P. E., Sejr, M. K., Bluhm, B. A., Sirenko, B., & Ellingsen, I. H. (2015). The future of Arctic benthos: expansion, invasion, and biodiversity. *Progress in Oceanography*, 139, 244-257.
- Rice, D. W. (1981). Status of the eastern Pacific (California) stock of the gray whale. In *Mammals in the Seas, General Paper and Large Cetaceans* (Vol. 3, pp. 181-187). Rome, Italy: FAO.
- Rice, D. W. (1998). *Marine mammals of the world: systematics and distribution*. (Special Publication Number 4). Lawrence, KS: Society for Marine Mammology, p. 231.
- Rice, D. W., & Wolman, A. A. (1971). *The life history and ecology of the gray whale (Eschrichtius robustus)* (Vol. 3). Seattle, WA: The American Society of Mammalogists.
- Rice, D. W., Wolman, A. A., & Braham, H. W. (1984). The Life History and Ecology of the Gray Whale (*Eschrichtius robustus*). *Marine Fisheries Review*, 46(4), 7-14.
- Rice, D. W., Wolman, A. A., Withrow, D. E., & Fleischer, L. A. (1981). Gray whales on the winter grounds in Baja California. *Reports of the International Whaling Commission*, 31, 477-493.
- Richard, P. R., Heide-Jørgensen, M. P., Orr, J. R., Dietz, R., & Smith, T. G. (2001). Summer and autumn movements and habitat use by belugas in the Canadian high Arctic and adjacent areas. *Arctic*, 54(3), 207-222.
- Richardson, W. J., Greene Jr., C. R., Hanna, J. S., Koski, W. R., Miller, G. W., Patenaude, N. J., & Smultea, M. A. (1995a). *Acoustic effects of oil production activities on bowhead and white whales visible during spring migrations near Point Barrow, Alaska - 1991 and 1994 phases: sound propagation and whale responses to playbacks of icebreaker noise*. (14-12-001-30412). Anchorage, AK: U.S. Department of the Interior. p. 570.
- Richardson, W. J., Greene Jr., C. R., Malme, C. I., & Thomson, D. H. (1995b). *Marine Mammals and Noise*. San Diego, CA: Academic Press.
- Richter, C. F., Dawson, S. M., & Slooten, E. (2003). *Sperm Whale Watching off Kaikoura, New Zealand: Effects of Current Activities on Surfacing and Vocalisation Patterns*. (Science for Conservation 219). Wellington, New Zealand: Department of Conservation. p. 78.
- Richter, C. F., Gordon, J. C. D., Jaquet, N., & Würsig, B. (2008). Social Structure of Sperm Whales in the Northern Gulf of Mexico. *Gulf of Mexico Science*, 26(2), 118-123.
- Roberts, L., & Breithaupt, T. (2016). Sensitivity of Crustaceans to Substrate-Borne Vibration. In *The Effects of Noise on Aquatic Life II* (pp. 925-931): Springer.
- Roberts, L., & Laidre, M. E. (2019). *Noise alters chemically-mediated search behavior in a marine hermit crab: Studying cross-modal effects on behavior*. Paper presented at the Proceedings of Meetings on Acoustics 5ENAL.
- Rosowski, J. J. (1994). Outer and Middle Ears. In Fay, R. R. & Popper, A. N. (Eds.), *Comparative Hearing: Mammals*: Springer-Verlag.
- Roth, E. H., Schmidt, V., Hildebrand, J. A., & Wiggins, S. M. (2013). Underwater radiated noise levels of a research icebreaker in the central Arctic Ocean. *Journal of Acoustical Society of America*, 133(4), 1971-1980.
- Rudels, B., Jones, E., Erson, L., & Kattner, G. (1994). On the intermediate depth waters of the Arctic Ocean. *AGU Physical Monograph*, 85, 33-46.

- Rudels, B., Muench, R., Gunn, J., Schauer, U., & Friedrich, H. (2000). Evolution of the Arctic Ocean boundary current north of the Siberian shelves. *J. Mar. Sys.*, 25(1), 77-99.
- Rugh, D. J., DeMaster, D. P., Rooney, A., Breiwick, J. M., Shelden, K. E. W., & Moore, S. (2003). A review of bowhead whale (*Balaena mysticetus*) stock identity. *Journal of Cetacean Research and Management*, 5(3), 267-280.
- Rugh, D. J., & Shelden, K. E. W. (2009). Bowhead whale *Balaena mysticetus*. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (2nd ed., pp. 131-133). Amsterdam, The Netherlands: Academic Press.
- Rugh, D. J., Shelden, K. E. W., & Schulman-Janiger, A. (2001). Timing of the gray whale southbound migration. *Journal of Cetacean Research and Management*, 3(1), 31-39.
- Salmon, M. (1971). Signal characteristics and acoustic detection by the fiddler crabs, *Uca rapax* and *Uca pugilator*. 44, 210-224.
- Sand, O., & Karlsen, H. E. (1986). Detection of Infrasound by the Atlantic Cod. *Journal of Experimental Biology*, 125, 197-204.
- Savage, K. (2021). *2020 Alaska Region Marine Mammal Stranding Summary*. Juneau, AK: National Marine Fisheries Service.
- Schack, H. B., Malte, H., & Madsen, P. T. (2008). The responses of Atlantic cod (*Gadus morhua* L.) to ultrasound-emitting predators: stress, behavioural changes or debilitation? *The Journal of Experimental Biology*, 211, 2079-2086.
- Schlundt, C. E., Finneran, J. J., Carder, D. A., & Ridgway, S. H. (2000). Temporary Shift in Masked Hearing Thresholds of Bottlenose Dolphins, *Tursiops truncatus*, and White Whales, *Delphinapterus leucas*, after Exposure to Intense Tones. *Journal of the Acoustical Society of America*, 107(6), 3496-3508.
- Schneider, D. C., Harrison, N. M., & Hunt Jr., G. L. (1990). Seabird diet at a front near the Pribilof Islands, Alaska. *Studies in Avian Biology*, 14, 61-66.
- Schoeman, R. P., Patterson-Abrolat, C., & Plön, S. (2020). A global review of vessel collisions with marine animals. *Frontiers in Marine Science*, 7, 292.
- Scholik, A. R., & Yan, H. Y. (2001). Effects of Underwater Noise on Auditory Sensitivity of a Cyprinid Fish. *Hearing Research*, 152, 17-24.
- Serreze, M. C., Maslanik, J. A., Scambos, T. A., Fetterer, F., Stroeve, J., Knowles, K., Fowler, C., Drobot, S., Barry, R. G., & Haran, T. M. (2003). A Record Minimum Arctic Sea Ice Extent and Area in 2002. *Geophysical Research Letters*, 30(3), 10/1-10/4.
- Shaughnessy, P. D., & Fay, F. H. (1977). A review of the taxonomy and nomenclature of North Pacific harbour seals. *Journal of Zoology, London*, 182, 385-419.
- Sheffield-Guy, L., Duffy-Anderson, J., Matarese, A. C., Mordy, C. W., Napp, J. M., & Stabeno, P. J. (2014). Understanding climate control of fisheries recruitment in the eastern Bering Sea: Long-term measurements and process studies. 27(4), 90-103.
- Shelden, K. E. W. (1994). *Beluga whales (Delphinapterus leucas) in Cook Inlet - A review*. 1335 East-West Highway, Silver Spring, MD 20910: Office of Protected Resources, National Marine Fisheries Service, National Oceanic and Atmospheric Administration.
- Shimada, K., Carmack, E., Hatakeyama, K., & Takizawa, T. (2001). Varieties of shallow temperature maximum waters in the western Canadian Basin of the Arctic Ocean. *Geophys. Res. Lett.*, 28(18), 3441-3444.
- Sills, J. M., Southall, B. L., & Reichmuth, C. (2015). Amphibious hearing in ringed seals (*Pusa hispida*): underwater audiograms, aerial audiograms and critical ratio measurements. *Journal of Experimental Biology*. doi: 10.1242/jeb.120972.

- Simpkins, M. A., Hiruki-Raring, L. M., Sheffield, G., Grebmeier, J. M., & Bengston, J. L. (2003). Habitat selection by ice-associated pinnipeds near St. Lawrence Island, Alaska in March 2001. *Polar Biology*, 26, 577-586.
- Simpson, S. D., Radford, A. N., Nedelec, S. L., Ferrari, M. C. O., Chivers, D. P., McCormick, M. I., & Meekan, M. G. (2016). Anthropogenic noise increases fish mortality by predation. 7.
- Simpson, S. D., Radford, A. N., Tickle, E. J., Meekan, M. G., & Jeffs, A. G. (2011). Adaptive avoidance of reef noise. *PLoS One*, 6(2), 1-5.
- Sirenko, B. I. (2001). List of species of free-living invertebrates of Eurasian Arctic seas and adjacent deep waters. *Explorations of the Fauna of the Seas*, 51(59).
- Sirenko, B. I., Clarke, C., Hopcroft, R. R., Huettmann, F., Bluhm, B. A., & Gradinger, R. (2010). The Arctic Register of Marine Species (ARMS) compiled by the Arctic Ocean Diversity (ArcOD) Retrieved from <http://www.marinespecies.org/arms> as accessed on 2 September 2014.
- Širović, A., Hildebrand, J. A., & Wiggins, S. M. (2007). Blue and fin whale call source levels and propagation range in the Southern Ocean. *Journal of the Acoustical Society of America*.
- Sivle, L. D., Kvadsheim, P. H., Fahlman, A., Lam, F. P. A., Tyack, P. L., & Miller, P. J. O. (2012). Changes in dive behavior during naval sonar exposure in killer whales, long-finned pilot whales, and sperm whales. *Frontiers in Physiology*, 3(Article 400), 1-11. doi: 10.3389/fphys.2012.00400.
- Smith, J., Ellis, K., & Boyd, T. (1999). Circulation features in the central Arctic Ocean revealed by nuclear fuel reprocessing tracers from Scientific Ice Expeditions 1995 and 1996. *J. Geophys. Res.*, 104(C12), 29663-29677.
- Smith, M., Goldman, M., Knight, E., & Warrenchuk, J. (2017). Ecological Atlas of the Bering. *Chukchi, and Beaufort Seas. 2nd ed Anchorage, AK: Audubon Alaska*.
- Smith, M. E., Coffin, A. B., Miller, D. L., & Popper, A. N. (2006). Anatomical and Functional Recovery of the Goldfish (*Carassius auratus*) Ear following Noise Exposure. *Journal of Experimental Biology*, 209, 4193-4202.
- Smith, M. E., Kane, A. S., & Popper, A. N. (2004a). Acoustical Stress and Hearing Sensitivity in Fishes: Does the Linear Threshold Shift Hypothesis Hold Water? *The Journal of Experimental Biology*, 207, 3591-3602.
- Smith, M. E., Kane, A. S., & Popper, A. N. (2004b). Noise-Induced Stress Response and Hearing Loss in Goldfish (*Carassius auratus*). *The Journal of Experimental Biology*, 207, 427-435.
- Smith, T. G. (1987). *The ringed seal, Phoca hispida, of the Canadian western Arctic*. Bulletin Fisheries Research Board of Canada. p. 81.
- Smith, T. G., & Hammill, M. O. (1981). Ecology of the ringed seal, *Phoca hispida*, in its fast ice breeding habitat. *Canadian Journal of Zoology*, 59, 966-981.
- Smith, T. G., Hammill, M. O., & Taugbøl, G. (1991). A review of the developmental, behavioural and physiological adaptations of the ringed seal, *Phoca hispida*, to life in the Arctic winter. *Arctic*, 44(2), 124-131.
- Smith, T. G., & Lydersen, C. (1991). Availability of suitable land-fast ice and predation as factors limiting ringed seal populations, *Phoca hispida*, in Svalbard. *Polar Research*, 10(2), 585-594.
- Smith, T. G., & Stirling, I. (1975). The Breeding Habitat of the Ringed Seal (*Phoca hispida*). The Birth Lair and Associated Structures. *Canadian Journal of Zoology*, 53, 1297-1305.
- Smultea, M. A., Brueggeman, J., Robertson, F., Fertl, D., Bacon, C., Rowlett, R. A., & Green, G. A. (2016). Polar Bear (*Ursus maritimus*) Behavior near Icebreaker Operations in the Chukchi Sea, 1991. 69(2).
- Smultea, M. A., & Mobley, J. R. (2009). Appendix A - Aerial Survey Monitoring for Marine Mammals and Sea Turtles in Conjunction with SCC OPS Navy Exercises off Kauai, 18-21 August 2008, Final Report, May 2009. p. 32.

- Solé, M., Lenoir, M., Fontuño, J. M., Durfort, M., Van der Schaar, M., & André, M. (2016). Evidence of Cnidarians sensitivity to sound after exposure to low frequency noise underwater sources. *Scientific reports*, 6(1), 1-18.
- Solé, M., Sigray, P., Lenoir, M., van der Schaar, M., Lalander, E., & André, M. (2017). Offshore exposure experiments on cuttlefish indicate received sound pressure and particle motion levels associated with acoustic trauma. *Scientific reports*, 7(1), 1-13.
- Sørensen, K., Neumann, C., Dähne, M., Hansen, K. A., & Wahlberg, M. (2020). Gentoo penguins (*Pygoscelis papua*) react to underwater sounds. *Royal Society open science*, 7(2), 191988.
- Soudijn, F. H., van Kooten, T., Slabbekoorn, H., & de Roos, A. M. (2020). Population-level effects of acoustic disturbance in Atlantic cod: a size-structured analysis based on energy budgets. *Proceedings of the Royal Society B*, 287(1929), 20200490.
- Southall, B. L., Bowles, A. E., Ellison, W. T., Finneran, J. J., Gentry, R. L., Greene Jr., C. R., Kastak, D., Ketten, D. R., Miller, J. H., Nachtigall, P. E., Richardson, W. J., Thomas, J. A., & Tyack, P. L. (2007). Marine mammal noise exposure criteria: initial scientific recommendations. *Aquatic Mammals*, 33(4), 411-521.
- Southall, B. L., Finneran, J. J., Reichmuth, C., Nachtigall, P. E., Ketten, D. R., Bowles, A. E., Ellison, W. T., Nowacek, D. P., & Tyack, P. L. (2019). Marine mammal noise exposure criteria: updated scientific recommendations for residual hearing effects. *Aquatic Mammals*, 45(2).
- St Aubin, D., & Dierauf, L. (2001). Stress and marine mammals. *CRC handbook of marine mammal medicine*, 2, 253-269.
- Staaterman, E. R., Clark, C. W., Gallagher, A. J., deVries, M. S., Claverie, T., & Patek, S. N. (2011). Rumbling in the benthos: acoustic ecology of the California mantis shrimp *Hemisquilla californiensis*. *Aquatic Biology*, 13, 97-105.
- Staaterman, E. R., Clark, C. W., Gallagher, A. J., deVries, M. S., Claverie, T., & Patek, S. N. (2016). Rumbling in the benthos: Acoustic ecology of the California mantis shrimp *Hemisquilla californiensis*. *Aquatic Biology*, 13, 97-105. doi: 10.3354/ab00361.
- Stanley, J. A., Van Parijs, S. M., & Hatch, L. T. (2017). Underwater sound from vessel traffic reduces the effective communication range in Atlantic cod and haddock. *Scientific reports*, 7(1), 1-12.
- Stewart, W. J., Nair, A., Jiang, H., & McHenry, M. J. (2014). Prey fish escape by sending the bow wave of a predator. *Journal of Experimental Biology*, 217, 4328-4336. doi: 10.1242/jeb.111773.
- Stimpert, A. K., Wiley, D. N., Au, W. W. L., Johnson, M. P., & Arsenault, R. (2007). 'Megapclicks': Acoustic click trains and buzzes produced during night-time foraging of humpback whales (*Megaptera novaeangliae*). *Biology Letters*.
- Stirling, I. (1997). The importance of polynas, ice edges, and leads to marine mammals and birds. *Journal of Marine Systems*, 10(1), 9-21.
- Stirling, I., Lunn, N. J., & Iacozza, J. (1999). Long-term trends in the population ecology of polar bears in western Hudson Bay in relation to climatic change. *Arctic*, 52(3), 294-306.
- Stringer, W. J. (1974). *Morphology of the Beaufort Sea Shorefast Ice*. Arlington, VA: Arctic Institute of North America.
- Suydam, R. S. (2009). *Age, growth, reproduction, and movements of beluga whales (Delphinapterus leucas) from the eastern Chukchi Sea*. University of Washington, School of Aquatic and Fishery Sciences.
- Suydam, R. S., Lowry, L. F., Frost, K. J., O'Corry-Crowe, G. M., & Pikok Jr., D. (2001). Satellite tracking of eastern Chukchi Sea beluga whales in to the Arctic Ocean. *Arctic*, 54(3), 237-243.
- Swartz, S. L., Taylor, B. L., & Rugh, D. J. (2006). Gray whale *Eschrichtius robustus* population and stock identity. *Mammal Review*, 36, 66-84.
- Symon, C., Arris, L., & Heal, B. (Eds.). (2005). *Arctic Climate Impact Assessment*. New York, NY: Cambridge University Press.

- Terhune, J. M., & Ronald, K. (1971). The Harp Seal, *Pagophilus groenlandicus* (Erxleben, 1777). X. The Air Audiogram. *Canadian Journal of Zoology*, 49(3), 385-390.
- Terhune, J. M., & Ronald, K. (1972). The Harp Seal, *Pagophilus groenlandicus* (Erxleben, 1777). III. The Underwater Audiogram. *Canadian Journal of Zoology*, 50(5), 565-569.
- Thiebault, A., Charrier, I., Aubin, T., Green, D. B., & Pistorius, P. A. (2019). First evidence of underwater vocalisations in hunting penguins. *PeerJ*, 7, e8240.
- Thiessen, G. J. (1958). Threshold of hearing of a ring-billed gull. *Journal of the Acoustical Society of America*, 30(11).
- Thistle, D. (2003). The Deep-Sea Floor: An Overview. In: Tyler, P. A. (Ed.), *Ecosystems of the Deep Ocean* (pp. 5-37). Amsterdam, Netherlands: Elsevier.
- Tidau, S., & Briffa, M. (2016). *Review on behavioral impacts of aquatic noise on crustaceans*. Paper presented at the Proceedings of Meetings on Acoustics 4ENAL.
- Timmermans, M. L., & Proshutinsky, A. (2014). Arctic Ocean Sea Surface Temperature [in Arctic Report Card 2014].
- Tynan, C. T., & DeMaster, D. P. (1997). Observations and Predictions of Arctic Climatic Change: Potential Effects on Marine Mammals. 50(4), 308-322.
- U.S. Department of the Navy. (2011). *Appendix E – Information on Sightings Recorded by U.S. Navy MMOs on Vessels during Sonar Test Events in the NSWC PCD Study Area*.
- U.S. Department of the Navy. (2014). *Marine Species Monitoring Report for the U.S. Navy's Atlantic Fleet Active Sonar Training (AFAST) and Virginia Capes, Cherry Point, Jacksonville, and Gulf of Mexico Range Complexes - Annual Report 2013*. U.S. Fleet Forces Command. p. 150.
- U.S. Department of the Navy. (2017a). *Criteria and Thresholds for U.S. Navy Acoustic and Explosive Effects Analysis (Phase III)*. p. 180.
- U.S. Department of the Navy. (2017b). *Dive distribution and group size parameters for marine species occurring in the U.S. Navy's Atlantic and Hawaii-Southern California training and testing Study Areas*.
- U.S. Department of the Navy. (2017c). *Final Overseas Environmental Assessment for Office of Naval Research Arctic Activities in the Beaufort Sea 2018-2021*.
- U.S. Department of the Navy. (2017d). *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing*.
- U.S. Department of the Navy. (2018a). *Final Overseas Environmental Assessment for Office of Naval Research Arctic Activities in the Beaufort Sea 2018-2021*.
- U.S. Department of the Navy. (2018b). *Hawaii-Southern California Training and Testing Environmental Impact Statement/Overseas Environmental Impact Statement*. Pearl Harbor, Hawaii: Naval Facilities Engineering Command,.
- U.S. Department of the Navy. (2021a). *A Blue Arctic: A Strategic Blueprint for the Arctic*.
- U.S. Department of the Navy. (2021b). *Final Supplemental Overseas Environmental Assessment for Arctic Research Activities in the Beaufort and Chukchi Seas*.
- U.S. Fish and Wildlife Service. (2006). *Short-Tailed Shearwater*. p. 2.
- U.S. Fish and Wildlife Service. (2021). Polar Bear Stock Assessment Report Retrieved from <https://www.fws.gov/media/polar-bear-southern-beaufort-sea-stock-assessment-report> as accessed on 02 May 2022.
- United Nations Environment Programme. (2012). *Scientific synthesis on the impacts of underwater noise on marine and coastal biodiversity and habitats*. p. 93.
- United States Coast Guard. (2013). CGC Healy Ship's Characteristics Retrieved from <http://www.uscg.mil/pacarea/cgchealy/ship.asp> as accessed on 03 September 2015.
- University of Alaska Fairbanks. (2014). Sikuliaq Specifications: R/V Sikuliaq Characteristics Retrieved from <https://www.sikuliaq.alaska.edu/ops/?q=node/19> as accessed on 03 September 2015.

- van Waerebeek, K., Baker, A. N., Felix, F., Gedamke, J., Iniguez, M., Sanino, G. P., Secchi, E., Sutaria, D., van Helden, A., & Wang, Y. (2007). Vessel Collisions with Small Cetaceans Worldwide and with Large Whales in the Southern Hemisphere, an Initial Assessment. *Latin American Journal of Aquatic Mammals*, 6(1), 43-69.
- Vancoppenolle, M., Meiners, K. M., Michel, C., Bopp, L., Brabant, F., Carnat, G., Delille, B., Lannuzel, D., Madec, G., & Moreau, S. (2013). Role of sea ice in global biogeochemical cycles: emerging views and challenges. *Quaternary science reviews*, 79, 207-230.
- Vanderlaan, A. S. M., & Taggart, C. T. (2007). Vessel Collisions with Whales: The Probability of Lethal Injury Based on Vessel Speed. *Marine Mammal Science*, 23(1), 144-156.
- Virketis, M. A. (1957). Some data on the zooplankton from the central part of the Arctic Basin. In *Materials of scientific observations of the drift stations "North Pole 3" and "North Pole 4" 1954/1955* (pp. 238-311). Moscow, Russia.
- Wadhams, P. (2000). Ice in the Ocean. In (pp. 351). United Kingdom: Gordon and Breach Science Publishers.
- Waga, H., Hirawake, T., & Grebmeier, J. M. (2020). Recent change in benthic macrofaunal community composition in relation to physical forcing in the Pacific Arctic. *Polar Biology*, 43(4), 285-294.
- Wale, M. A., Briers, R. A., Hartl, M. G., Bryson, D., & Diele, K. (2019). From DNA to ecological performance: Effects of anthropogenic noise on a reef-building mussel. *Science of The Total Environment*, 689, 126-132.
- Wale, M. A., Simpson, S. D., & Radford, A. N. (2013). Noise negatively affects foraging and antipredator behaviour in shore crabs. *Animal Behaviour*, 86, 111-118.
- Walther, G. R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., Fromentin, J., Hoegh-Guldberg, O., & Bairlein, F. (2002). Ecological Responses to Recent Climate Change. *Nature*, 416, 389-395.
- Ward, W. D. (1997). Effects of high-intensity sound. In Crocker, M. J. (Ed.), *Encyclopedia of Acoustics* (pp. 1497-1507). New York, NY: Wiley.
- Wartzok, D., Elsner, R., Stone, H., Kelly, B. P., & Davis, R. W. (1992a). Under-ice movements and the sensory basis of hole finding by ringed and Weddell seals. *Canadian Journal of Zoology*, 70(9), 1712-1722.
- Wartzok, D., & Ketten, D. R. (1999). *Marine mammal sensory systems*. Washington, DC: Smithsonian Institution Press.
- Wartzok, D., Popper, A. N., Gordon, J., & Merrill, J. (2003). Factors affecting the responses of marine mammals to acoustic disturbance. *Marine Technology Society Journal*, 37(4), 6-15.
- Wartzok, D., Sayegh, S., Stone, H., Barchak, J., & Barnes, W. (1992b). Acoustic tracking system for monitoring under-ice movements of polar seals. *Journal of the Acoustical Society of America*, 92, 682-687.
- Watkins, W. A. (1986). Whale Reactions to Human Activities in Cape Cod Waters. *Marine Mammal Science*, 2(4), 251-262.
- Watwood, S., Fagan, M., D'Amico, A., & Jefferson, T. (2012). *Cruise Report, Marine Species Monitoring & Lookout Effectiveness Study Koa Kai, November 2011, Hawaii Range Complex*. p. 12.
- Webb, J. F., Montgomery, J. C., & Mogdans, J. (2008). Bioacoustics and the Lateral Line of Fishes. *Fish Bioacoustics*.
- Wei, C. L., Cusson, M., Archambault, P., Belley, R., Brown, T., Burd, B. J., Edinger, E., Kenchington, E., Gilkinson, K., & Lawton, P. (2020). Seafloor biodiversity of Canada's three oceans: Patterns, hotspots and potential drivers. *Diversity and Distributions*, 26(2), 226-241.
- Weinberg, H., & Keenan, R. (2008). *CASS V4.2 Software Requirements Specification(SRS), Software Design Document(SDD) and Software Test Description(STD)*. Alion Science and Technology Corporation.

- Weingartner, T., Shimada, K., McLaughlin, F., & Proshutinsky, A. (2008). *Physical Oceanography*. Fairbanks, AK: Institute of Marine Sciences, University of Alaska. pp. 6-17.
- Welch, H. E., Crawford, R. E., & Hop, H. (1993). Occurrence of Arctic cod (*Boreogadus saida*) schools and their vulnerability to predation in the Canadian high Arctic. *Arctic*, 46(4), 331-339.
- Weller, D. W., Bettridge, S., Brownell Jr, R. L., Laake, J. L., Moore, J. E., Rosel, P. E., Taylor, B. L., & Wade, P. R. (2013). *Report of the National Marine Fisheries Service gray whale stock identification workshop*. (NOAA Technical Memo. NOAA-TM-NMFS-SWFSC-507).
- Wever, E. G., Herman, P. N., Simmons, J. A., & Hertzler, D. R. (1969). Hearing in the blackfooted penguin, *Spheniscus demersus*, as represented by the cochlear potentials. *Proceedings of the National Academy of Sciences USA*, 63(3), 676-680.
- Whitledge, T., Mathis, J., Sapozhnikov, V., & Moran, B. (2008). *Chemical Oceanography*. Fairbanks, AK: Institute of Marine Sciences, University of Alaska.
- Wiig, O., Bachmann, L., Janik, V. M., Kovacs, K. M., & Lydersen, C. (2007). Spitsbergen bowhead whales revisited. *Marine Mammal Science*, 23(3), 688-693. doi: 10.1111/j.1748-7692.2007.02373.x.
- Wilkinson, T. A., Wiken, E., Creel, J. B., Hourigan, T. F., & Agardy, T. (2009). *Marine Ecoregions of North America*: Instituto Nacional de Ecologia.
- Williams, R., Erbe, C., Ashe, E., Beerman, A., & Smith, J. (2014). Severity of killer whale behavioral responses to ship noise: A dose-response study. 79, 254-260.
- Williams, R., Lusseau, D., & Hammond, P. S. (2006). Estimating relative energetic costs of human disturbance to killer whales (*Orcinus orca*). 133, 301-311.
- Williams, R., Lusseau, D., & Hammond, P. S. (2009). The Role of Social Aggregations and Protected Areas in Killer Whale Conservation: The Mixed Blessing of Critical Habitat. *Biological Conservation*, 142, 709-719.
- Winfree, M. (2005). *Preliminary aerial reconnaissance surveys of eelgrass beds on Togiak National Wildlife Refuge, Alaska, 2004*. Togiak National Wildlife Refuge. p. 10.
- Woodgate, R. (2012). Arctic Ocean Circulation: Going Around At the Top Of the World. *Nature Education Knowledge*, 4(8), 8.
- Woodgate, R., Aagaard, K., Muench, R., Gunn, J., Bjork, G., Rudels, B., Roach, A., & Schauer, U. (2001). The Arctic Ocean boundary current along the Eurasian slope and the adjacent Lomonosov Ridge: Water mass properties, transports and transformations from moored instruments. *Deep Sea Res., Part I*, 48(8), 1757-1792.
- Woodgate, R., Aagaard, K., & Weingartner, T. J. (2005). Monthly temperature, salinity, and transport variability of the Bering Strait through flow. *Geophys. Res. Lett.*, 32(4). doi: 10.1029/2004GL021880.
- Wyllie-Echeverria, S., & Ackerman, J. D. (2003). *The Pacific Coast of North America*. Berkeley, CA: University of California Press.
- Wysocki, L. E., Dittami, J. P., & Ladich, F. (2006). Ship noise and cortisol secretion in European freshwater fishes. *Biological Conservation*, 128(4), 501-508.

This page intentionally left blank

8 List of Preparers

This OEA was prepared collaboratively between the Navy and contractor preparers.

U.S. Department of the Navy

Emily Robinson (Naval Undersea Warfare Center, Division Newport)
Masters of Environmental Science and Management
B.S. Integrated Science and Technology
Years of Experience: 8 years Environmental Planning
Responsible for: Project Lead, Project Coordination, Document Development

Cassandra DePietro (Naval Undersea Warfare Center, Division Newport)
Masters of Applied Math
B.S. Mathematics
Years of Experience: 5 year Modeling and Prototype
Responsible for: Marine Mammal Modeling and Prototyper

Raymond Soukup (Office of Naval Research)
M.A. Mathematical Statistics
B.S. Physics
Years of Experience: 32 years Ocean Acoustics Research and Program Management
Responsible for: Program Officer, Acoustician, and Environmental Planner

Contractors

Jessica Greene (McLaughlin Research Corporation)
Masters of Environmental Science and Management
B.S. Wildlife Conservation
Years of Experience: 8 years GIS Analyst
Responsible for: GIS

Katherine Griswold (McLaughlin Research Corporation)
M.S. Marine Biological Resources
B.S. Marine Science
Years of Experience: 1 Year Environmental Planning
Responsible for: Document Development

David Loiselle (McLaughlin Research Corporation)
Masters of Environmental Science and Management
B.S. Environmental Science and Management
Years of Experience: 3 years Environmental Planning
Responsible for: Document Development

This page intentionally left blank

Appendix A Stressor Matrices

Ten categories of stressors were identified and analyzed within this OEA. Stressors applicable to each activity and resource are provided in Appendix Table A-1 and

Appendix Table A-2. A description of each stressor, including the platforms that contribute to the stressor, is provided below.

- **Non-Impulsive Acoustic Sources:** Includes only those active sources that may harm a resource from acoustics that are not considered *de minimis* and require quantitative analysis.
- **Icebreaking Noise:** Includes noise from CGC HEALY when icebreaking.
- **Vessel Noise:** Includes the noise generated by the R/V Sikuliaq and CGC HEALY. This does not include the sound CGC HEALY generates when icebreaking.
- **Icebreaking (Physical Impacts):** Includes the potential for harm to resources from ice breaking apart, due to CGC HEALY breaking ice as it moves through the Study Area.
- **Vessel and In-Water Device Strike:** Includes the potential for vessels (i.e., surface ships) and in-water devices (e.g., gliders) to come into direct contact with a resource.
- **Bottom Disturbance:** Includes the potential for the material to strike a resource as it sinks and settles on the seafloor. Expended material is also analyzed for potential disturbance to the seafloor.
- **Entanglement:** Includes the potential for a resource to become entangled in a temporarily-deployed device and those materials that will be expended.
- **Ingestion:** Includes the possibility of ingesting complete objects as well as small pieces of objects to determine if they are edible.

Appendix Table A-1. Stressors by Activity

<i>Action</i>	<i>Acoustic Stressors</i>			<i>Physical Stressors</i>			<i>Expended Material</i>	
	<i>Non-Impulsive Acoustic Sources</i>	<i>Icebreaking Noise</i>	<i>Vessel Noise</i>	<i>Icebreaking (Physical Impacts)</i>	<i>Vessel and In-Water Device Strike</i>	<i>Bottom Disturbance</i>	<i>Entanglement</i>	<i>Ingestion</i>
Glider Surveys					X			
Research Vessel Activities		X	X	X	X		X	
Ship-Deployed Active Acoustic Sources	X				X		X	
Moored Acoustic Sources	X				X	X	X	
De minimis Sources					X	X		
Drifting Oceanographic Sensors						X	X	
Moored Oceanographic Sensors					X	X	X	
On-Ice Measurement Systems								X

Appendix Table A-2. Stressors by Resource

<i>Resource</i>	<i>Acoustic Stressors</i>			<i>Physical Stressors</i>			<i>Expended Material</i>	
	<i>Non-Impulsive Acoustic Sources</i>	<i>Icebreaking Noise</i>	<i>Vessel Noise</i>	<i>Icebreaking (Physical Impacts)</i>	<i>Vessel and In- Water Device Strike</i>	<i>Bottom Disturbance</i>	<i>Entanglement</i>	<i>Ingestion</i>
Ice				X				
Bottom Substrate						X		
Marine Mammals	X	X	X	X	X		X	X
Marine Birds	X	X	X					
Invertebrates	X	X	X	X	X	X	X	
Fish	X	X	X	X	X	X	X	
Essential Fish Habitat	X	X		X				

Appendix B Non-Impulsive and Impulsive Source Modeling

B.1. Introduction

The marine mammal acoustics effects analysis was conducted in accordance with current Navy sonar policy, as advised by the Chief of Naval Operations Environmental Readiness Division. Accordingly, ensonified areas and exposure estimates for marine mammals were reported based on SEL and SPL thresholds. PTS is the criterion used to establish the onset of non-recoverable physiological effects. TTS is the criterion used to establish the onset of recoverable physiological effects, and a BRF is used to determine non-physiological behavioral effects. Environmental parameters were collected and archived, and propagation modeling was performed with the Naval Oceanographic Office's Oceanographic and Atmospheric Master Library CASS/GRAB model (Weinberg and Keenan 2008). The acoustics effects modeling utilized the databases and tools collectively referred to as NAEMO (U.S. Department of the Navy 2017d). Results were then computed for the defined operational scenario. This section provides a brief discussion of several key components of the acoustics effects modeling process, specifically: environmental inputs, acoustic sources, propagation modeling, and the NAEMO modeling software suite.

B.2. Source Characteristics and Scenario Description

The parameters for the acoustic and impulsive transmissions associated with research activities can be found in Table 2-1 above, the parameters for icebreaking can be found in Table 4-3 and Table 4-4 above.

B.3. Environmental Characteristics

Data for four environmental characteristics (bathymetry, sound speed profile, sediment characteristics, and wind speed) were obtained for both the cold and warm seasons to support the acoustic and impulsive analysis. The databases used to obtain these data and the resulting parameters are provided in Appendix Table B-1. All of the databases are maintained by the Oceanographic and Atmospheric Master Library.

Appendix Table B-1. Environmental Parameters for ARA

<i>Model / Parameter</i>	<i>Data Input</i>	<i>Database</i>
Propagation Model	Specific data are not applicable for this parameter.	Comprehensive Acoustic System Simulation Version 4.3b
Absorption Model	Specific data are not applicable for this parameter.	Francois-Garrison (the CASS/GRAB default)
Analysis Locations	Study Area	Database not used for this parameter
Analysis Specifics	Acoustic sources: 18 radials => 1 radial per 20 degrees Impulsive sources: 9 radials => 1 radial per 40 degrees Range increment: 50 meters* Depth increment: 25 meters*	Database not used for this parameter
Bathymetry	Data was obtained from a location centered around 72° 53'N, 146° 28'W. Resolution was at five hundredths (0.5) of a degree.	Digital Bathymetric Data Base Variable Resolution (DBDB-V) Version 6.2
Sound Speed Profiles	Sound speed profiles were extracted at the highest database resolution of 0.25 degree.	Generalized Digital Environmental Model Variable (GDEM-V) Version 3.0
Wind Speed	Wind speed was extracted at the highest database resolution of one (1) degree. Average wind speed: N/A for the cold season since the Study Area is ice covered	Surface Marine Gridded Climatology (SMGC) Version 2.0
Geo-Acoustic Parameters	Sediment type of sand was determined for the Study Area.	High Frequency Environmental Acoustics Version 2 HFEVA
Surface Reflection Coefficient Model	Specific data are not applicable for this parameter.	Navy Standard Forward Surface Loss Model

*Range and depth increments for impulsive source modeling are not uniform. The steps are small when close to the source and spread out when moving away from the source. Increments shown are largest steps.

B.4. Marine Mammal Density Estimates

Marine mammal densities utilized in the acoustic analysis were based on the best available science for the Study Area. Density data estimates were obtained from Kaschner et al. (2006) and Cañadas et al. (2020). The Navy used a Seasonal Relative Environmental Suitability model (Kaschner et al. 2006), based on seasonal habitat preferences and requirements of known occurrences, such as temperature, bathymetry, and distance to land data and literature review, because occurrence information for marine mammals in the Study Area is not well known. Empirical data is coupled with Relative Environmental Suitability modeling data to generate predictions of density data for locations where no survey data exist. Densities derived from survey data were used when available, primarily in the most southern portions of the Study Area during the warm season (Cañadas et al. 2020).

B.5. Criteria and Thresholds

Harassment criteria for marine mammals are evaluated based on thresholds developed from observations of trained cetaceans exposed to intense underwater sound under controlled conditions (Finneran and Schlundt 2003; Kastak and Schusterman 1996; Kastak and Schusterman 1999; Kastak et al. 2005; Kastelein et al. 2012). These data are the most applicable because they are based on controlled, tonal sound exposures within the typical sonar frequency ranges and because the species studied are

closely related to the animals expected in the Study Area. Studies have reported behavioral alterations, or deviations from a subject’s normal trained behavior, and exposure levels above which animals were observed to exhibit behavioral deviations (Finneran and Schlundt 2003; Schlundt et al. 2000).

Criteria and thresholds used for determining the potential effects from the Proposed Action are from NMFS technical guidance on acoustic and impulsive thresholds for PTS/TTS. The behavioral criteria was developed in coordination with NMFS to support Phase III environmental analyses and MMPA Letter of Authorization renewals (U.S. Department of the Navy 2017a). Appendix Table B-2 below provides the criteria and thresholds used in this analysis for estimating quantitative acoustic and impulsive exposures of marine mammals from the Proposed Action, respectively. Weighted criteria are shown in the table below. Frequency-weighting functions are used to adjust the received sound level based on the sensitivity of the animal to the frequency of the sound. For weighting function derivation, the most critical data required are TTS onset exposure levels as a function of exposure frequency. These values can be estimated from published literature by examining TTS as a function of SEL for various frequencies.

Appendix Table B-2. Acoustic Injury (PTS) and Disturbance (TTS, Behavioral) Thresholds for Underwater Sounds¹

Group	Species	Behavioral Criteria		Physiological Criteria	
		Non-Impulsive Acoustic Sources	Icebreaking Sources	Onset TTS	Onset PTS
Low Frequency Cetaceans	Gray whale, bowhead whale	Low-Frequency BRF dose response function ³	120 dB re 1 μPa step function	179 dB SEL cumulative	199 dB SEL cumulative
Mid Frequency Cetaceans	Beluga whale	Mid-Frequency BRF dose response function ³	120 dB re 1 μPa step function	178 dB SEL cumulative	198 dB SEL cumulative
Phocidae (in water)	Bearded seal, pacific walrus, ribbon seal, spotted seal, ringed seal	Pinniped Dose Response Function ³	120 dB re 1 μPa step function	181 dB SEL cumulative	201 dB SEL cumulative
Otariidae (in water) and other non-phocid marine carnivores	Polar bear	Pinniped Dose Response Function ³	120 dB re 1 μPa step function	199 dB SEL cumulative	219 dB SEL cumulative

¹ The threshold values provided are assumed for when the source is within the animal’s best hearing sensitivity. The exact threshold varies based on the overlap of the source and the frequency weighting.

² BRF = Behavioral Response Function

³ See Appendix Figure B-1

To estimate TTS onset values, only TTS data from behavioral hearing tests were used. To determine TTS onset for each subject, the amount of TTS observed after exposures with different SPLs and durations were combined to create a single TTS growth curve as a function of SEL. The use of (cumulative) SEL is a simplifying assumption to accommodate sounds of various SPLs, durations, and duty cycles. This is referred to as an “equal energy” approach, since SEL is related to the energy of the sound and this approach assumes exposures with equal SEL result in equal effects, regardless of the duration or duty cycle of the sound. It is well-known that the equal energy rule will over-estimate the effects of intermittent noise, since the quiet periods between noise exposures will allow some recovery of hearing compared to noise that is continuously present with the same total SEL (Ward 1997). For continuous

exposures with the same SEL but different durations, the exposure with the longer duration will also tend to produce more TTS (Finneran et al. 2010; Kastak et al. 2007; Mooney et al. 2009).

As in previous acoustic effects analysis (Finneran and Jenkins 2012; Southall et al. 2007), the shape of the PTS exposure function for each species group is assumed to be identical to the TTS exposure function for each group. A difference of 20 dB between TTS onset and PTS onset is used for all marine mammals including pinnipeds. This is based on estimates of exposure levels actually required for PTS (i.e. 40 dB of TTS) from the marine mammal TTS growth curves, which show differences of 13 to 37 dB between TTS and PTS onset in marine mammals. Details regarding these criteria and thresholds can be found in National Marine Fisheries Service (2016).

B.5.1. Behavioral Reactions or Responses

Behavioral criteria for both acoustic and impulsive sources are described below.


B.5.1.1. Acoustic Criteria

The response of a marine mammal to an anthropogenic sound will depend on the frequency, duration, temporal pattern and amplitude of the sound as well as the animal's prior experience with the sound and the context in which the sound is encountered (i.e., what the animal is doing at the time of the exposure). The distance from the sound source and whether it is perceived as approaching or moving away can also affect the way an animal responds to a sound (Wartzok et al. 2003). For marine mammals, a review of responses to anthropogenic sound was first conducted by Richardson *et al.* (1995b). Reviews by Nowacek *et al.* (2007) and Southall *et al.* (2007) address studies conducted since 1995 and focus on observations where the received sound level of the exposed marine mammal(s) was known or could be estimated. Multi-year research efforts have conducted sonar exposure studies for odontocetes and mysticetes (Miller et al. 2012; Sivle et al. 2012). Several studies with captive animals have provided data under controlled circumstances for odontocetes and pinnipeds (Houser et al. 2013a; Houser et al. 2013b). Moretti *et al.* (2014) published a beaked whale dose-response curve based on passive acoustic monitoring of beaked whales during U.S. Navy training activity at Atlantic Underwater Test and Evaluation Center during actual Anti-Submarine Warfare exercises. This new information has necessitated the update of the Navy's behavioral response criteria.

Southall *et al.* (2007) synthesized data from many past behavioral studies and observations to determine the likelihood of behavioral reactions at specific sound levels. While in general, the louder the sound source the more intense the behavioral response, it was clear that the proximity of a sound source and the animal's experience, motivation, and conditioning were also critical factors influencing the response (Southall et al. 2007). After examining all of the available data, the authors felt that the derivation of thresholds for behavioral response based solely on exposure level was not supported because context of the animal at the time of sound exposure was an important factor in estimating response. Nonetheless, in some conditions, consistent avoidance reactions were noted at higher sound levels depending on the marine mammal species or group allowing conclusions to be drawn. Phocid seals showed avoidance reactions at or below 190 dB re 1 μ Pa at 1m; thus, seals may actually receive levels adequate to produce TTS before avoiding the source.

The Phase III pinniped behavioral criteria was updated based on controlled exposure experiments on the following captive animals: hooded seal, gray seal, and California sea lion (Götz et al. 2010; Houser et al. 2013a; Kvadsheim et al. 2010). Overall exposure levels were 110-170 dB re 1 μ Pa for hooded seals, 140-180 dB re 1 μ Pa for gray seals and 125-185 dB re 1 μ Pa for California sea lions; responses occurred at received levels ranging from 125 to 185 dB re 1 μ Pa. However, the means of the response data were between 159 and 170 dB re 1 μ Pa. Hooded seals were exposed to increasing levels of sonar until an

avoidance response was observed, while the gray seals were exposed first to a single received level multiple times, then an increasing received level. Each individual California sea lion was exposed to the same received level ten times, these exposure sessions were combined into a single response value, with an overall response assumed if an animal responded in any single session. Because these data represent a dose-response type relationship between received level and a response, and because the means were all tightly clustered, the Bayesian biphasic Behavioral Response Function for pinnipeds most closely resembles a traditional sigmoidal dose-response function at the upper received levels (Appendix Figure B-1), and has a 50% probability of response at 166 dB re 1 μ Pa. Additionally, to account for proximity to the source discussed above and based on the best scientific information, a conservative distance of 5.4 nautical miles (10 km) is used beyond which exposures would not constitute a take under the military readiness definition.



Appendix Figure B-1. A) The Bayesian biphasic dose-response BRF for Odontocetes. B) The Bayesian biphasic dose-response BRF for Pinnipeds C) The Bayesian biphasic dose-response BRF for Mysticetes. The blue solid line represents the Bayesian Posterior median values, the green dashed line represents the biphasic fit, and the grey represents the variance. [X-Axis: Received Level (dB re 1 μ Pa), Y-Axis: Probability of Response]

B.5.1.2. NAEMO Software

The Navy performed a quantitative analysis to estimate the number of mammals that could be harassed by the underwater acoustic (non-impulsive and impulsive) sources during the Proposed Action. Inputs to the quantitative analysis included marine mammal density estimates obtained from the NMSDD, marine mammal depth occurrence distributions (U.S. Department of the Navy 2017b), oceanographic and environmental data, marine mammal hearing data, and criteria and thresholds for levels of potential effects. The quantitative analysis consists of computer modeled estimates and a post-model analysis to determine the number of potential animal exposures. The model calculates sound energy propagation from the proposed sonars, the sound received by animat (virtual animal) dosimeters representing

marine mammals distributed in the area around the modeled activity, and whether the sound received by a marine mammal exceeds the thresholds for effects.

The Navy developed a set of software tools and compiled data for estimating acoustic effects on marine mammals without consideration of behavioral avoidance or Navy's standard mitigations. These databases and tools collectively form NAEMO. In NAEMO, animats are distributed nonuniformly based on species-specific density, depth distribution, and group size information. Animats record energy received at their location in the water column. A fully three-dimensional environment is used for calculating sound propagation and animat exposure in NAEMO. Site-specific bathymetry, sound speed profiles, wind speed, and bottom properties are incorporated into the propagation modeling process. NAEMO calculates the likely propagation for various levels of energy (sound or pressure) resulting from each source used during the testing event.

NAEMO then records the energy received by each animat within the energy footprint of the event and calculates the number of animats having received levels of energy exposures that fall within defined impact thresholds. Predicted effects on the animats within a scenario are then tallied and the highest order effect (based on severity of criteria; e.g., PTS over TTS) predicted for a given animat is assumed. Each scenario or each 24-hour period for scenarios lasting greater than 24 hours is independent of all others, and therefore, the same individual marine animal could be impacted during each independent scenario or 24-hour period. In few instances, although the activities themselves all occur within the Study Area, sound may propagate beyond the boundary of the Study Area. Any exposures occurring outside the boundary of the Study Area are counted as if they occurred within the Study Area boundary. NAEMO provides the initial estimated impacts on marine species with a static horizontal distribution.

There are limitations to the data used in the acoustic effects model, and the results must be interpreted within these context. While the most accurate data and input assumptions have been used in the modeling, when there is a lack of definitive data to support an aspect of the modeling, modeling assumptions believed to overestimate the number of exposures have been chosen:

- Animats are modeled as being underwater, stationary, and facing the source and therefore always predicted to receive the maximum sound level (i.e., no porpoising or pinnipeds' heads above water).
- Animats do not move horizontally (but change their position vertically within the water column), which may overestimate physiological effects such as hearing loss, especially for slow moving or stationary sound sources in the model.
- Animats are stationary horizontally and therefore do not avoid the sound source, unlike in the wild where animals would most often avoid exposures at higher sound levels, especially those exposures that may result in PTS.
- Multiple exposures within any 24-hour period are considered one continuous exposure for the purposes of calculating the temporary or permanent hearing loss, because there are not sufficient data to estimate a hearing recovery function for the time between exposures.
- Mitigation measures that are implemented were not considered in the model. In reality, sound-producing activities would be reduced, stopped, or delayed if marine mammals are detected within the mitigation zones around sound sources.

Because of these inherent model limitations and simplifications, model-estimated results must be further analyzed, considering such factors as the range to specific effects, avoidance, and the likelihood of successfully implementing mitigation measures. This analysis uses a number of factors in addition to the acoustic model results to predict acoustic effects on marine mammals.

For non-impulsive acoustic sources, NAEMO calculates the SPL and SEL for each active emission during an event. This is done by taking the following factors into account over the propagation paths: bathymetric relief and bottom types, sound speed, and attenuation contributors such as absorption, bottom loss and surface loss. Platforms such as a ship using one or more sound sources are modeled in accordance with relevant vehicle dynamics and time durations by moving them across an area whose size is representative of the testing event's operational area. For each modeled iteration, the slow moving platform in this experiment was programmed to move along straight line tracks from a randomly selected initial location with a randomly selected course. Specular reflection was employed at the boundaries to contain the vehicle within the Study Area.

NAEMO records the SPL and SEL received by each animat within the ensonified area of the event and evaluates them in accordance with the species-specific threshold criteria. For each animat, predicted SEL effects are accumulated over the course of the event and the highest order SPL effect is determined. Each 24-hour period is independent of all others, and therefore, the same individual animat could be exposed during each independent scenario or 24-hour period. Initially, NAEMO provides the overpredicted exposures to marine species because predictions used in the model include: all animats facing the source, not accounting for horizontal avoidance and mitigation is not implemented. After the modeling results are complete they are further analyzed to produce final estimates of potential marine mammal exposures.

B.6. Results

For non-impulsive acoustic sources, NAEMO calculates maximum received SPL and accumulated SEL over the entire duration of the event for each animat based on the received sound levels. These data are then processed using a bootstrapping routine to compute the number of animats exposed to SPL and SEL in 1 dB bins across all track iterations and population draws. SEL is checked during this process to ensure that all animats are grouped in either an SPL or SEL category. Additional detail on the bootstrapping process is included in Section B.7.1.

A mean number of SPL and SEL exposures are computed for each 1 dB bin. The mean value is based on the number of animats exposed at that dB level from each track iteration and population draw. The behavioral risk function curve is applied to each 1 dB bin to compute the number of behaviorally exposed animats per bin. The number of behaviorally exposed animats per bin is summed to produce the total number of behavior exposures.

Mean 1 dB bin SEL exposures are then summed to determine the number of PTS and TTS exposures. PTS exposures represent the cumulative number of animats exposed at or above the PTS threshold. The number of TTS exposures represents the cumulative number of animats exposed at or above the TTS threshold and below the PTS threshold. Animats exposed below the TTS threshold were grouped in the SPL category.

B.7.1. Bootstrap Approach

Estimation of exposures in NAEMO is accomplished through the use of a simple random sampling with replacement by way of statistical bootstrapping. This sampling approach was chosen due to the fact that the number of individuals of a species expected within an area over which a given Navy activity occurs is often too small to offer a statistically significant sampling of the geographical area. Additionally, NAEMO depends on the fact that individual animats move vertically in the water column at a specified displacement frequency for sufficient sampling of the depth dimension. By overpopulating at the time of animat distribution and drawing samples from this overpopulation with replacement, NAEMO is able to provide sufficient sampling in the horizontal dimensions for statistical confidence. Sampling with

replacement also produces statistically independent samples, which allows for the calculation of metrics such as standard error and confidence intervals for the underlying Monte Carlo process.

For each scenario and each species, the number of samples equating to the overpopulation factor is drawn from the raw data. Each sample size consists of the true population size of the species evaluated. Exposure data is then computed for each sample using 1 dB exposure bins. The average number of exposures across the sample and scenario iteration is then computed.

For example, assuming that an overpopulation factor of 10 was defined for a given species and that 15 ship track iterations were completed. The bootstrap Monte Carlo process would have generated statistics for 10 draws on each of the 15 raw animat data files generated by the 15 ship tracks evaluated for this scenario, thereby yielding 150 independent sets of exposure estimates. Samples drawn from the overpopulated population are replaced for the next draw, allowing for the re-sampling of animals. The resultant 150 sets of exposures were then combined to yield a mean number of exposures and a 95 percent confidence interval per species for the scenario. In addition to the mean, the statistics included the upper and lower bounds of all samples.

B.7.2. Estimated Exposures

Based on the methodology contained herein, Appendix Table B-4 provides the annual modeled marine mammal exposures associated with the thresholds defined in Section B.5 for 2022-2025.

Appendix Table B-3. Predicted Annual Marine Mammal Exposures All Events (Acoustic and Icebreaking)

<i>Species</i>	<i>Alternative 1</i>			<i>Alternative 2 (Preferred Alternative)</i>		
	<i>Behavioral</i>	<i>TTS</i>	<i>PTS</i>	<i>Behavioral</i>	<i>TTS</i>	<i>PTS</i>
Beluga whale	289	0	0	289	0	0
Bowhead whale ¹	0	0	0	0	0	0
Gray whale	0	0	0	0	0	0
Polar bear ¹	0	0	0	0	0	0
Bearded seal ¹	0	0	0	0	0	0
Ribbon seal	0	0	0	0	0	0
Ringed seal ¹	3,377	1	0	3,377	1	0
Spotted seal	0	0	0	0	0	0
Pacific walrus	0	0	0	0	0	0

¹ESA-listed species