

## **SOWERBY’S BEAKED WHALE (*Mesoplodon bidens*): Western North Atlantic Stock**

### **STOCK DEFINITION AND GEOGRAPHIC RANGE**

Within the genus *Mesoplodon*, there are four species of beaked whales that reside in the northwest Atlantic. These include True’s beaked whale, *M. mirus*; Gervais’ beaked whale, *M. europaeus*; Blainville’s beaked whale, *M. densirostris*; and Sowerby’s beaked whale, *M. bidens* (Mead 1989). These species are difficult to identify to the species level at sea; therefore, much of the available characterization for beaked whales is to genus level only. Stock structure for each species is unknown. Thus, it is plausible the stock could actually contain multiple demographically independent populations since the current stock spans multiple eco-regions (Longhurst 1998; Spalding et al. 2007).

The distributions of *Mesoplodon* spp. in the northwest Atlantic are known principally from stranding records (Mead 1989; Nawojchik 1994; Mignucci-Giannoni et al. 1999; MacLeod et al. 2006). Off the U.S. Atlantic coast, beaked whale (*Mesoplodon* spp.) sightings have occurred principally along the shelf-edge and deeper oceanic waters (Figure 1; CETAP 1982; Waring et al. 1992; Tove 1995; Waring et al. 2001; Hamazaki 2002; Palka 2006). Most sightings occurred in late spring and summer, corresponding to survey effort. The distributions of Sowerby’s beaked whales are also known from acoustical surveys (Cholewiak et al. 2013, Stanistreet et al. 2018) and bycatch confirmed genetically to be *M. bidens* (Wenzel et al. 2013).

Sowerby’s beaked whales represent a transboundary stock reported to occur from New England waters north to the ice pack (e.g., Davis Strait), and individuals are seen along the Newfoundland coast in summer (Leatherwood et al. 1976; Mead 1989; MacLeod et al. 2006; Jefferson et al. 2008) (Figure 1). Furthermore, a single stranding occurred off the Florida west coast (Mead 1989). This species is considered rare in Canadian waters (Lien et al. 1990) and has been designated as “Special Concern” by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC). Whitehead (2013) reports that in the 23 years of cetacean observations in the Gully Marine Protected Area, on the edge of the Scotian Shelf, Nova Scotia, Canada, they have observed a significant increase in sightings of Sowerby’s.

### **POPULATION SIZE**

The best abundance estimate for Sowerby’s beaked whales is the sum of the 2021 survey estimates—492 (CV=0.50). This estimate, derived from shipboard surveys, covers most of this stock’s known range. In the 2021 survey, improvements to field protocols for both visual observers and passive acoustic monitoring of *Mesoplodon* spp. facilitated differentiation of species during encounters. This enabled abundance estimates to be calculated for each species individually rather than grouping together at the genus level. .

### **Recent Surveys and Abundance Estimates**

Abundance estimates of 6,760 (CV=0.37) and 3,347 (CV=0.29) *Mesoplodon* spp. beaked whales (not including *Ziphius*) were generated from vessel surveys conducted in U.S. waters of the western North Atlantic during the summer of 2016 (Table 1; Garrison 2020; Palka 2020). One survey was conducted from 27 June to 25 August in waters north of 38°N latitude and consisted of 5,354 km of on-effort trackline along the shelf break and offshore to the outer limit of the U.S. EEZ (NEFSC and SEFSC 2018). The second vessel survey covered waters from Central Florida to approximately 38°N latitude between the 100-m isobath and the outer limit of the U.S. EEZ during 30 June–19 August. A total of 4,399 km of trackline was covered on effort (NEFSC and SEFSC 2018). Both surveys utilized two visual teams and an independent observer approach to estimate detection probability on the trackline (Laake and Borchers 2004). Mark-recapture distance sampling was used to estimate abundance. Estimates from the two surveys were combined and CVs pooled to produce an abundance estimate for the stock area, yielding a combined total of 10,107 *Mesoplodon* beaked whales (CV=0.27). These estimates are known to be biased low due to the fact that unidentified Ziphiidae abundance was estimated at 3,755 (CV=0.42) in the NE and at 2,812 (CV=0.43) in the SE. These estimates likely include an unknown number of *Mesoplodon* beaked whales.

A more recent abundance estimate of 492 (CV=0.50) Sowerby’s beaked whales was generated from vessel surveys conducted in U.S. waters of the western North Atlantic during the summer of 2021 (Table 1; Garrison and Dias 2023; Palka 2023). One survey was conducted from 16 June to 23 August in waters north of 36°N latitude and consisted of 5,871 km of on-effort trackline along the shelf break and offshore to the outer edge of the U.S. EEZ (NEFSC and SEFSC 2022). The second vessel survey covered waters from central Florida (25°N latitude) to

approximately 38°N latitude between the 200-m isobaths and the outer edge of the U.S. EEZ during 12 June–31 August. A total of 5,659 km of trackline was covered on effort (NEFSC and SEFSC 2022). Both surveys utilized two visual teams and an independent observer approach to estimate detection probability on the trackline (Laake and Borchers 2004). Mark-recapture distance sampling was used to estimate abundance. Estimates from the two surveys were combined and CVs pooled to produce a species abundance estimate for the stock area.

**Table 1. Summary of abundance estimates for the Mesoplodon beaked whales (2016 surveys) and Sowerby’s beaked whales (2021 surveys), month, year, and area covered during each abundance survey, and resulting abundance estimate ( $N_{best}$ ) and coefficient of variation (CV).**

Month/Year	Area	$N_{best}$	CV
Jun–Sep 2016	Central Virginia to lower Bay of Fundy ( <i>Mesoplodon</i> spp.)	6,760	0.37
Jun–Aug 2016	Central Florida to Central Virginia ( <i>Mesoplodon</i> spp.)	3,347	0.29
Jun–Sep 2016	Central Florida to lower Bay of Fundy (COMBINED, <i>Mesoplodon</i> spp.)	10,107	0.27
Jun–Aug 2021	New Jersey to lower Bay of Fundy ( <i>M. bidens</i> only)	492	0.50
Jun–Aug 2021	Central Florida to New Jersey ( <i>M. bidens</i> only)	0	0
<b>Jun–Aug 2021</b>	<b>Central Florida to lower Bay of Fundy (COMBINED, <i>M. bidens</i> only)</b>	<b>492</b>	<b>0.50</b>

### Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for Sowerby’s beaked whales is 492 (CV=0.50). The minimum population estimate for Sowerby’s beaked whales in the western North Atlantic is 340.

### Current Population Trend

There are insufficient data to determine population trends for this stock. The statistical power to detect a trend in abundance for this stock is poor due to the relatively imprecise abundance estimates and long survey interval. For example, the power to detect a precipitous decline in abundance (i.e., 50% decrease in 15 years) with estimates of low precision (e.g., CV > 0.30) remains below 80% (alpha = 0.30) unless surveys are conducted on an annual basis (Taylor et al. 2007). There is current work to standardize the strata-specific previous abundance estimates to consistently represent the same regions and include appropriate corrections for perception and availability bias. These standardized abundance estimates will be used in state-space trend models that incorporate environmental factors that could potentially influence the process and observational errors for each stratum.

### CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. *Mesoplodon bidens* life history parameters that could be used to estimate net productivity include: length at birth of up to 2.4 m and maximum length of 5 m for females and 5.5 m for males (Mead 1984).

For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow et al. 1995).

### POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a recovery factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size for Sowerby’s beaked whales is 492. The maximum productivity rate is 0.04, the default value for cetaceans. The recovery factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP) is assumed to be 0.5. PBR for Sowerby’s beaked whales in the western North Atlantic is 3.4.

**Table 2. Best and minimum abundance estimates for Sowerby’s beaked whales (*Mesoplodon bidens*) of the Western North Atlantic with Maximum Productivity Rate ( $R_{max}$ ), Recovery Factor ( $F_r$ ) and PBR.**

$N_{est}$	CV $N_{est}$	$N_{min}$	$F_r$	$R_{max}$	PBR
492	0.50	340	0.5	0.04	3.4

### ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

The 2017–2021 total average estimated annual mortality of Sowerby’s beaked whales in observed fisheries in the U.S. Atlantic EEZ is zero.

#### Fishery Information

Total fishery-related mortality and serious injury cannot be estimated separately for each beaked whale species because of the uncertainty in species identification by fishery observers. The Atlantic Scientific Review Group advised adopting the risk-averse strategy of assuming that any beaked whale stock which occurred in the U.S. Atlantic EEZ might have been subject to the observed fishery-related mortality and serious injury.

In 2017–2021, estimated annual average fishery-related mortality or serious injury of this stock in U.S. fisheries was 0 for all beaked whales. Detailed fishery information is reported in Appendix III.

#### Earlier Interactions

See Appendix V for more information on historical takes.

#### Other Mortality

During 2017–2021 1 Sowerby’s beaked whale stranded along the U.S. Atlantic coast without evidence of human interaction (Table 3; NOAA National Marine Mammal Health and Stranding Response Database, accessed 15 October 2022).

**Table 3. Sowerby’s beaked whale (*Mesoplodon bidens*) strandings along the U.S. Atlantic coast.**

State	2017	2018	2019	2020	2021	Total
Maine	0	0	0	1	0	1
Massachusetts	0	0	0	0	0	0
<b>Total</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>1</b>

### STATUS OF STOCK

While Sowerby’s beaked whales are not listed as threatened or endangered under the Endangered Species Act, they have been listed as a species of Special Concern by both COSEWIC and SARA (the Species at Risk Act) in Canada due to concerns about potential effects on this species from widespread seismic operations as well as occasional military sonar use off of Canada’s East Coast (COSEWIC 2006). The western North Atlantic stock of Sowerby’s beaked whale is not considered strategic under the Marine Mammal Protection Act. There are insufficient data to determine the population size or trends. The permanent closure of the pelagic drift gillnet fishery has eliminated the principal known source of incidental fishery mortality, and no fishery-related mortality and serious injury has been observed during the recent 5-year (2013–2017) period. Therefore, the total U.S. fishery mortality and serious injury rate can be considered to be insignificant and approaching zero. The status of Sowerby’s beaked whales relative to OSP is unknown.

### OTHER FACTORS THAT MAY BE AFFECTING THE STOCK

#### Habitat Issues

The chronic impacts of contaminants (polychlorinated biphenyls [PCBs] and chlorinated pesticides [DDT, DDE, dieldrin, etc.]) on marine mammal reproduction and health are of concern (e.g., Pierce et al. 2008; Jepson et al. 2016; Hall et al. 2018; Murphy et al. 2018), but research on contaminant levels for the western north Atlantic beaked whales is lacking.

Anthropogenic sound in the world’s oceans has been shown to affect marine mammals, with vessel traffic, seismic

surveys, and active naval sonars being the main anthropogenic contributors to low- and mid-frequency noise in oceanic waters (e.g., Nowacek et al. 2015; Gomez et al. 2016; NMFS 2018). Questions have been raised regarding potential effects of human-made sounds on deep-diving cetacean species such as Sowerby's beaked whales (Richardson et al. 1995). The long-term and population consequences of these impacts are less well-documented and likely vary by species and other factors. Impacts on marine mammal prey from sound are also possible (Carroll et al. 2017), but the duration and severity of any such prey effects on marine mammals are unknown.

Several unusual mass strandings of beaked whales throughout their worldwide range have been associated with naval activities (D'Amico et al. 2009; Filadelfo et al. 2009). During the mid- to late 1980s multiple mass strandings of Cuvier's beaked whales (4 to about 20 per event) and small numbers of Gervais' beaked whale and Blainville's beaked whales occurred in the Canary Islands (Simmonds and Lopez-Jurado 1991). Twelve Cuvier's beaked whales that live stranded and subsequently died in the Mediterranean Sea on 12-13 May 1996 were associated with low frequency acoustic sonar tests conducted by the North Atlantic Treaty Organization (Frantzis 1998; D'Amico et al. 2009; Filadelfo et al. 2009). In March 2000, 14 beaked whales live stranded in the Bahamas; 6 beaked whales (5 Cuvier's and 1 Blainville's) died (Balcomb and Claridge 2001; NMFS 2001; Cox et al. 2006). Four Cuvier's, 2 Blainville's, and 2 unidentified beaked whales were returned to sea. The fate of the animals returned to sea is unknown, since none of the whales have been resighted. Necropsy of 6 dead beaked whales revealed evidence of tissue trauma associated with an acoustic or impulse injury that caused the animals to strand. Subsequently, the animals died due to extreme physiologic stress associated with the physical stranding (i.e., hyperthermia, high endogenous catecholamine release) (Cox et al. 2006).

Fourteen beaked whales (mostly Cuvier's beaked whales but also including Gervais' and Blainville's beaked whales) stranded in the Canary Islands in 2002 (Cox et al. 2006, Fernandez et al. 2005; Martin et al. 2004). Gas bubble-associated lesions and fat embolism were found in necropsied animals from this event, leading researchers to link nitrogen supersaturation with sonar exposure (Fernandez et al. 2005).

Climate-related changes in spatial distribution and abundance, including poleward and depth shifts, have been documented in or predicted for plankton species and commercially important fish stocks (Nye et al. 2009; Head et al. 2010; Pinsky et al. 2013; Poloczanska et al. 2013; Hare et al. 2016; Grieve et al. 2017; Morley et al. 2018) and cetacean species (e.g., MacLeod 2009; Sousa et al. 2019). Chavez-Rosales et al. (2022) documented an overall 178 km northeastward spatial distribution shift of the seasonal core habitat of Northwest Atlantic cetaceans that was related to changing habitat/climatic factors. Results varied by season and species. This study used sightings data collected during seasonal aerial and shipboard line transect abundance surveys during 2010 to 2017. There is uncertainty in how, if at all, the changes in distribution and population size of cetacean species may interact with changes in distribution of prey species and how the ecological shifts will affect human impacts to the species.

## REFERENCES CITED

- Balcomb, K.C.I. and D.E. Claridge. 2001. A mass stranding of cetaceans caused by naval sonar in the Bahamas. *Bahamas J. Sci.* 2: 2-12.
- Barlow, J., M.C. Ferguson, W.F. Perrin, L. Balance, T. Gerrodette, G. Joyce, C.D. MacLeod, K. Mullin, D.L. Palka and G. Waring. 2006. Abundance and densities of beaked and bottlenose whales (family *Ziphiidae*). *J. Cetacean Res. Manage.* 7: 263-270.
- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade. 1995. U.S. Marine Mammal Stock Assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73 pp.
- Buckland, S.T., D.R. Anderson, K.P. Burnham, J.L. Laake, D.L. Borchers and L. Thomas. 2001. Introduction to distance sampling: estimating abundance of biological populations. Oxford University Press. 432 pp.
- Carroll, A.G., R. Przeslawski, A. Duncan, M. Gunning, B. Bruce. 2017. A critical review of the potential impacts of marine seismic surveys on fish & invertebrates. *Mar. Pollut. Bull.* 114:9-24.
- CETAP. 1982. A characterization of marine mammals and turtles in the mid- and north Atlantic areas of the U.S. outer continental shelf, final report, Cetacean and Turtle Assessment Program, University of Rhode Island. Washington, DC, Bureau of Land Management. #AA551-CT8-48: 576.
- Chavez-Rosales S., E. Josephson, D. Palka and L. Garrison. 2022. Detection of habitat shifts of cetacean species: a comparison between 2010 and 2017 habitat suitability conditions in the northwest Atlantic Ocean. *Front. Mar. Sci.* 9:877580. doi: 10.3389/fmars.2022.877580
- Cholewiak, D., S. Baumann-Pickering and S. Van Parijs. 2013. Description of sounds associated with Sowerby's beaked whales (*Mesoplodon bidens*) in the western North Atlantic Ocean. *J. Acoust. Soc. Am.* 134 (5):3905-3912.

- COSEWIC. 2006. COSEWIC Assessment and update status on the Sowerby's beaked whale in Canada, Committee on the Status of Endangered Wildlife in Canada. Ottawa, vi+20 p. [http://publications.gc.ca/collections/collection\\_2007/ec/CW69-14-502-2007E.pdf](http://publications.gc.ca/collections/collection_2007/ec/CW69-14-502-2007E.pdf)
- Cox, T. M., T. J. Ragen, A. J. Read, E. Vos, R. W. Baird, K. Balcomb, J. Barlow, J. Caldwell, T. Cranford, L. Crum, A. D'Amico, G. D. Spain, A. Fernandez, J. Finneran, R. Gentry, W. Gerth, F. Gulland, J. Hildebrand, D. Houser, T. Hullar, P. D. Jepson, D. Ketten, C. D. MacLeod, P. Miller, S. Moore, D. Mountain, D. Palka, P. Ponganis, S. Rommel, T. Rowles, B. Taylor, P. Tyack, D. Wartzok, R. Gisiner, J. Mead and L. Benner. 2006. Understanding the impacts of anthropogenic sound on beaked whales. *J. Cetacean Res. Manage.* 7(3): 177-187.
- D'Amico, A., R.C. Gisiner, D.R. Ketten, J.A. Hammock, C. Johnson, P.L. Tyack and J. Mead. 2009. Beaked whale strandings and naval exercises. *Aq. Mamm.* 35(4) 452-472.
- Fernandez, A., J.F. Edwards, F. Rodriguez, A.E. de los Monteros, P. Herraiez, P. Castro, J.R. Jaber, V. Martin and M. Arbelo. 2005. "Gas and Fat Embolic Syndrome" involving a mass stranding of beaked whales (Family *Ziphiidae*) exposed to anthropogenic sonar signals. *Veterinary Pathology* 42(4):446-457.
- Filadelfo, R., J. Mintz, E. Michlovich, A. D'Amico, P.L. Tyack and D.R. Ketten. 2009. Correlating military sonar use with beaked whale mass strandings: What do the historical data show? *Aq. Mamm.* 35(4) 435-444.
- Frantzis, A. 1998. Does acoustic testing strand whales? *Nature* 392: 29.
- Garrison, L.P. 2020. Abundance of cetaceans along the southeast U.S. east coast from a summer 2016 vessel survey. Southeast Fisheries Science Center, Protected Resources and Biodiversity Division, 75 Virginia Beach Dr., Miami, FL 33140. PRD Contribution # PRD-2020-04, 17 pp.
- Garrison, L.P. and L.A. Dias. 2023. Abundance of marine mammals in waters of the southeastern U.S. Atlantic during summer 2021. SEFSC MMTD Contribution: #MMTD-2023-01. 23 pp. <https://repository.library.noaa.gov/view/noaa/49152>
- Grieve, B.D., J.A. Hare and V.S. Saba. 2017. Projecting the effects of climate change on *Calanus finmarchicus* distribution within the US Northeast continental shelf. *Sci. Rep.* 7:6264.
- Gomez, C., J.W. Lawson, A.J. Wright, A.D. Buren, D. Tollit and V. Lesage. 2016. A systematic review on the behavioural responses of wild marine mammals to noise: The disparity between science and policy. *Can. J. Zool.* 94:801-819.
- Hare, J.A., W.E. Morrison, M.W. Nelson, M.M. Stachura, E.J. Teeters, R.B. Griffis, M.A. Alexander, J.D. Scott, L. Alade, R.J. Bell, A.S. Chute, K.L. Curti, T.H. Curtis, D. Kurcheis, J.F. Kocik, S.M. Lucey, C.T. McCandless, L.M. Milke, D.E. Richardson, E. Robillard, H.J. Walsh, M.C. McManus, K.E. Maranick and C.A. Griswold. 2016. A vulnerability assessment of fish and invertebrates to climate change on the Northeast U.S. continental shelf, *PLoS ONE* 11:e0146756. <https://doi.org/10.1371/journal.pone.0146756.s014>.
- Hall, A.J., B.J. McConnell, L.J. Schwacke, G.M. Ylitalo, R. Williams and T.K. Rowles. 2018. Predicting the effects of polychlorinated biphenyls on cetacean populations through impacts on immunity and calf survival. *Environ. Poll.* 233:407-418.
- Head, E.J.H., and P. Pepin. 2010. Spatial and inter-decadal variability in plankton abundance and composition in the Northwest Atlantic (1958-2006). *J. Plankton Res.* 32:1633-1648.
- Hamazaki, T. 2002. Spatiotemporal prediction models of cetacean habitats in the mid-western North Atlantic Ocean (from Cape Hatteras, No. Carolina, USA to Nova Scotia, Canada). *Mar. Mamm. Sci.* 18(4): 920-939.
- Jefferson, T.A., M.A. Webber, and R.L. Pitman 2008. *Marine Mammals of the World*. Elsevier, Amsterdam, Netherlands. 573 pp.
- Jepson, P.D., R. Deaville, J.L. Barber, A. Aguilar, A. Borrell, S. Murphy, J. Barry, A. Brownlow, J. Barnett, S. Berrow and A.A. Cunningham. 2016. PCB pollution continues to impact populations of orcas and other dolphins in European waters. *Sci. Rep.-U.K.* 6:18573.
- Laake, J. L., and D. L. Borchers. 2004. Methods for incomplete detection at distance zero. Pages 108-189 in: S.T. Buckland, D.R. Andersen, K.P. Burnham, J.L. Laake and L. Thomas (eds.), *Advanced distance sampling*. Oxford University Press, New York, New York. Leatherwood, S., D. K. Caldwell and H. E. Winn. 1976. Whales, dolphins, and porpoises of the western North Atlantic. A guide to their identification. NOAA Tech. Rep. NMFS Circ. 396. 176 pp.
- Lien, J., F. Barry, K. Breeck and U. Zuschlag. 1990. Status of Sowerby's Beaked Whale, *Mesoplodon bidens*, in Canada. *Can. Field-Nat.* 104(1): 125-130.
- Longhurst, A.R. 1998. *Ecological geography of the sea*, Second Edition., Elsevier Academic Press. 560 pp.
- Lucas, Z. N. and S. K. Hooker. 2000. Cetacean strandings on Sable Island, Nova Scotia, 1970-1998. *Can. Field-Nat.* 114(1): 46-61.

- MacLeod, C., W.F. Perrin, R. Pitman, J. Barlow, L. Ballance, A. D'Amico, T. Gerrodette, G. Joyce, K.D. Mullin, D.L. Palka and G.T. Waring. 2006. Known and inferred distributions of beaked whale species (Cetacea: Ziphiidae). *J. Cetacean Res. Manage.* 7(3): 271–286.
- MacLeod, C.D. 2009. Global climate change, range changes and potential implications for the conservation of marine cetaceans: a review and synthesis. *Endang. Species Res.* 7:125–136.
- Martín, V., A. Servidio and S. García 2004. Mass strandings of beaked whales in the Canary Islands. *ECS Newsletter* 42:33-6.
- Mead, J. G. 1984. Survey of reproductive data for the beaked whales (*Ziphiidae*). *Rep. Int. Whal. Comm.* (Special Issue) 6: 91-96.
- Mead, J. G. 1989. Beaked whales of the genus *Mesoplodon*. Pages 349-430 in: S.H. Ridgway and R. Harrison, (eds.) *Handbook of marine mammals*, Vol. 4: River Dolphins and toothed whales. Academic press, San Diego.
- Mignucci-Giannoni, A.A., B. Pinto-Rodríguez, M. Velasco-Escudero, R.A. Montoya-Ospina, N.M. Jiménez, M.A. Rodríguez-López, J.E.H. Williams and D.K. Odell 1999. Cetacean strandings in Puerto Rico and the Virgin Islands. *J. Cetacean Res. Manage.* 1: 191-198.
- Morley, J.W., R.L. Selden, R.J. Latour, T.L. Frolicher, R.J. Seagraves and M.L. Pinsky. 2018. Projecting shifts in thermal habitat for 686 species on the North American continental shelf. *PLoS ONE* 13(5):e0196127.
- Murphy, S., R.J. Law, R. Deaville, J. Barnett, M.W. Perkins, A. Brownlow, R. Penrose, N.J. Davison, J.L. Barber P.D. Jepson. 2018. Organochlorine contaminants and reproductive implication in cetaceans: A case study of the common dolphin. Pages 3–38 in M.C. Fossi and C. Panti, (eds.) *Marine mammal ecotoxicology: Impacts of multiple stressors on population health*. Academic Press, New York, New York.
- Nawojchik, R. 1994. First record of *Mesoplodon densirostris* (Cetacea: Ziphiidae) from Rhode Island. *Mar. Mamm. Sci.* 10: 477-480.
- NEFSC [Northeast Fisheries Science Center] and Southeast Fisheries Science Center [SEFSC]. 2022. 2021 Annual report of a comprehensive assessment of marine mammal, marine turtle, and seabird abundance and spatial distribution in US waters of the Western North Atlantic Ocean – AMAPPS III. 125 pp. <https://repository.library.noaa.gov/view/noaa/41734>
- NMFS 2001. Joint interim report on the Bahamas marine mammal stranding event of 15-16 March 2000 (December 2001). NOAA unpublished report 55 pp. <https://repository.library.noaa.gov/view/noaa/16198>
- NMFS [National Marine Fisheries Service]. 2018. 2018 Revisions to: Technical guidance for assessing the effects of anthropogenic sound on marine mammal hearing (Version 2.0): Underwater thresholds for onset of permanent and temporary threshold shifts. NOAA Tech. Memo. NMFS-OPR-59, 167 pp. Available from: <https://repository.library.noaa.gov/view/noaa/17892>
- Nowacek, D.P., C.W. Clark, D. Mann, P.J.O. Miller, H.C. Rosenbaum, J.S. Golden, M. Jasny, J. Kraska and B.L. Southall. 2015. Marine seismic surveys and ocean noise: time for coordinated and prudent planning. *Front. Ecol. Environ.* 13:378–386.
- Nye, J., J. Link, J. Hare and W. Overholtz. 2009. Changing spatial distribution of fish stocks in relation to climate and population size on the Northeast United States continental shelf. *Mar. Ecol. Prog. Ser.* 393:111–129.
- Palka, D.L. 2006. Summer abundance estimates of cetaceans in US North Atlantic Navy Operating Areas. *Northeast Fish. Sci. Cent. Ref. Doc.* 06-03. 41 pp. <https://repository.library.noaa.gov/view/noaa/5258>
- Palka, D. 2020. Cetacean abundance estimates in US northwestern Atlantic Ocean waters from summer 2016 line transect surveys conducted by the Northeast Fisheries Science Center. *Northeast Fish. Sci. Cent. Ref. Doc.* 20-05.
- Palka, D.L. 2012. Cetacean abundance estimates in US northwestern Atlantic Ocean waters from summer 2011 line transect survey. *Northeast Fish. Sci. Cent. Ref. Doc.* 12-29. 37 pp. <https://repository.library.noaa.gov/view/noaa/4312>
- Palka, DL. 2023. Cetacean abundance in the US Northwestern Atlantic Ocean, Summer 2021. *Northeast Fish. Sci. Cent. Ref. Doc.* 23-08.
- Pierce, G.J. M.B. Santos, S. Murphy, J.A. Learmonth, A.F. Zuur, E. Rogan, P. Bustamante, F. Caurant, V. Lahaye, V. Ridoux, B.N. Zegers, A. Mets, M. Addink, C. Smeenk, T. Jauniaux, R.J. Law, W. Dabin, A. López, J.M. Alonso Farré, A.F. González, A. Guerra, M. García-Hartmann, R.J. Reid, C.F. Moffat, C. Lockyer, J.P. Boon. 2008. Bioaccumulation of persistent organic pollutants in female common dolphins (*Delphinus delphis*) and harbour porpoises (*Phocoena phocoena*) from western European seas: Geographical trends, causal factors and effects on reproduction and mortality. *Environmental Pollution.* 153:401–415.
- Pinsky, M.L., B. Worm, M.J. Fogarty, J.L. Sarmiento and S.A. Levin. 2013. Marine taxa track local climate velocities. *Science* 341:1239–1242.
- Poloczanska, E.S., C.J. Brown, W.J. Sydeman, W. Kiessling, D.S. Schoeman, P.J. Moore, K. Brander, J.F. Bruno, L.B. Buckley, M.T. Burrows, C.M. Duarte, B.S. Halpern, J. Holding, C.V. Kappel, M.I. O'Connor, J.M.

- Pandolfi, C. Parmesan, F. Schwing, S.A. Thompson and A.J. Richardson. 2013. Global imprint of climate change on marine life. *Nat. Clim. Change* 3:919–925.
- Richardson, W.J., C.R. Greene Jr, C.I. Malme, and D.H. Thomson 1995 *Marine Mammals and Noise*, Academic Press, San Diego, CA. 576 pp.
- Simmonds, M.P. and L.F. Lopez-Jurado. 1991. Whales and the military. *Nature*: 351:448.
- Sousa, A., F. Alves, A. Dinis, J. Bentz, M.J. Cruz and J.P. Nunes. 2019. How vulnerable are cetaceans to climate change? Developing and testing a new index. *Ecol. Indic.* 98:9–18.
- Spalding, M.D., H.E. Fox, G.R. Allen, N. Davidson, Z.A. Ferdaña, M. Finlayson, B.S. Halpern, M.A. Jorge, A. Lombana, S.A. Lourie, K.D. Martin, E. McManus, J. Molnar, C.A. Recchia and J. Robertson. 2007. Marine ecoregions of the world: a bioregionalization of coastal and shelf areas. *BioScience* 57(7):573-583.
- Taylor, B.L., M. Martinez, T. Gerrodette, J. Barlow and Y.N. Hrovat. 2007. Lessons from monitoring trends in abundance in marine mammals. *Mar. Mamm. Sci.* 23(1): 157-175.
- Thomas L, J.L. Laake, E. Rexstad, S. Strindberg, F.F.C. Marques, S.T. Buckland, D.L. Borchers, D.R. Anderson, K.P. Burnham, M.L. Burt, S.L. Hedley, J.H. Pollard, J.R.B. Bishop and T.A. Marques. 2009. Distance 6.0. Release 2. [Internet]. University of St. Andrews (UK): Research Unit for Wildlife Population Assessment. Available from: <http://distancesampling.org/Distance/>.
- Tove, M. 1995. Live sighting of *Mesoplodon CF. M. Mirus*, True's Beaked Whale. *Mar. Mamm. Sci.* 11(1): 80-85.
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp. Available from: <https://repository.library.noaa.gov/view/noaa/15963>
- Waring, G.T., C.P. Fairfield, C.M. Ruhsam and M. Sano. 1992. Cetaceans associated with Gulf Stream Features off the Northeastern USA Shelf. *ICES [Int. Counc. Explor. Sea] C.M.* 1992/N:12.
- Waring, G.T., T. Hamazaki, D. Sheehan, G. Wood and S. Baker. 2001. Characterization of beaked whale (*Ziphiidae*) and sperm whale (*Physeter macrocephalus*) summer habitat in shelf-edge and deeper waters off the northeast U.S. *Mar. Mamm. Sci.* 17(4): 703-717.
- Wenzel, F.W., P.T. Polloni, J.E. Craddock, D.P. Gannon, J.R. Nicolas, A.J. Read, P.E. Rosel. 2013. Food habits of Sowerby's beaked whales, *Mesoplodon bidens*, taken in the western North Atlantic pelagic drift gillnet fishery. *Fish. Bul.* 111(4):381-389.
- Whitehead, H. 2013. Trends in cetacean abundance in the Gully submarine canyon, 1988–2011, highlight a 21% per year increase in Sowerby's beaked whales (*Mesoplodon bidens*). *Can. J. Zool.*, 2013, 91(3):141-148.